

UNIVERSIDAD NACIONAL DE CÓRDOBA
FACULTAD DE MATEMÁTICA, ASTRONOMÍA Y FÍSICA

SERIE “A”

TRABAJOS DE MATEMÁTICA

Nº 108/2011

ON NICHOLS ALGEBRAS OF DIAGONAL TYPE

Iván E. Angiono



Editores: Jorge G. Adrover – Gastón García
CIUDAD UNIVERSITARIA – 5000 CÓRDOBA
REPÚBLICA ARGENTINA

ON NICHOLS ALGEBRAS OF DIAGONAL TYPE

IVÁN ANGIOÑO

ABSTRACT. We give an explicit and essentially minimal list of defining relations of a Nichols algebra of diagonal type with finite root system. This list contains the well-known quantum Serre relations but also many new variations. A conjecture by Andruskiewitsch and Schneider states that any finite-dimensional pointed Hopf algebra over an algebraically closed field of characteristic zero is generated as an algebra by its group-like and skew-primitive elements. As an application of our main result, we prove the conjecture when the group of group-like elements is abelian.

INTRODUCTION

1. Let \mathbf{k} be an algebraically closed field of characteristic zero and let θ be a natural number. Let $\mathbf{q} = (q_{ij})_{1 \leq i, j \leq \theta}$ be a matrix with invertible entries on \mathbf{k} and let V be a vector space of dimension θ . The Nichols algebra associated to \mathbf{q} is a graded connected algebra $\mathcal{B}(V) = \bigoplus_{n \geq 0} \mathcal{B}^n(V)$ with many favourable properties. It plays a fundamental role in the classification of finite-dimensional (or finite growth) pointed Hopf algebras. Precisely, a basic question in the classification Program [AS1] is the following:

Question 1. [An, Question 5.9]: *Given (V, \mathbf{q}) , determine if the associated Nichols algebra $\mathcal{B}(V)$ is finite-dimensional. In such case, compute the dimension of $\mathcal{B}(V)$ and give a presentation by generators and relations.*

The first part of this question has been answered by Heckenberger [H3], who obtained the list of all matrices \mathbf{q} whose associated Nichols algebra has a finite root system. Roughly, this list contains three classes of matrices:

- Standard matrices [AA]: they are associated with finite Weyl groups. Their root systems coincide with root systems of finite Cartan matrices. This family includes properly the so-called braidings of Cartan type, in particular the matrices related with the positive part of the small quantum groups.
- Matrices of super type [AAY], related with the positive part of quantized enveloping algebras of contragredient Lie superalgebras. Their root systems become from the corresponding Lie superalgebras.
- A finite list of exceptional matrices, whose associated diagram has connected components with at most 7 vertices, and the scalars defining these braidings are roots of unity of low order.

There are several answers to the second part of Question 1 under particular assumptions:

- ▷ [L] for the positive part of quantized enveloping algebras of semisimple Lie algebras and small quantum groups, using the full representation theory of quantum groups;
- ▷ [AS2] for braidings of Cartan type;
- ▷ [A1] for braidings of standard type;
- ▷ [Y] for the positive part of quantized enveloping algebras of contragredient Lie superalgebras;

2010 *Mathematics Subject Classification.* 16W30.

The work was partially supported by CONICET, FONCyT-ANPCyT, Secyt (UNC), Mincyt (Córdoba).

- ▷ [AA Y] for braidings of super type;
- ▷ [H1], giving a general form of relations for matrices of rank two;
- ▷ [He] for some examples of rank two matrices, but giving explicit relations.

In [A2] we gave general formulae for defining relations of Nichols algebras of diagonal type, see Theorem 1.25 below. The expression of those relations and the proof that they generate the defining ideal are independent of Heckenberger's classification; they rely in Kharchenko's and Rosso's PBW bases [Kh, R] and a detailed study of convex orders in generalized root systems [A2], through the classification of coideal subalgebras [HS]. In this paper we refine the main result of [A2] and prove:

Theorem 1. *A minimal set of relations of $\mathcal{B}(V)$ is obtained by considering relations of the following type:*

- (1) *Quantum Serre relations, and powers of generators x_i corresponding to non-Cartan vertices; they are needed to introduce Lusztig's isomorphisms at the level of doubles of tensor algebras.*
- (2) *Relations in the image of the previous ones by the Lusztig's isomorphisms, and correspond to relations (23) in Theorem 1.25.*
- (3) *Relations that guarantee that the ideal generated by the previous relations is a braided biideal: they appear in the coproduct of relations of the item (2) in the tensor algebra $T(V)$.*
- (4) *Powers of root vectors (generators of the PBW basis) corresponding to roots in the orbit of Cartan vertices.*

See Theorem 3.1 for a complete and explicit set of relations. In this set we distinguish relations appearing in [A2] for standard braidings, and relations in [Y] related with braidings of super type. There exists also a large list of new relations, related with the set of exceptional braidings or with braidings of super type evaluated in roots of unity of small order.

2. The knowledge of the explicit relations of a Nichols algebra has several potential applications to the theory of pointed Hopf algebras, that we discuss now:

- One of the basic question in the Lifting Method [AS1, AS3] for the classification of Hopf algebras is the following:

Conjecture 1. [AS2, Conjecture 1.4] *Let Γ be a finite group and \mathbf{k} an algebraically closed field of characteristic 0. If H is a finite-dimensional pointed Hopf algebra over \mathbf{k} such that $G(H) = \Gamma$, then H is generated as an algebra by Γ and its skew-primitive elements.*

This question was answered in [AS4] for braidings of Cartan type under some mild conditions. This result was extended to the case of standard braidings in [AGI]. In Section 4 we obtain as a consequence of Theorem 3.1:

Theorem 2. *Let H be a finite dimensional pointed Hopf algebra over an abelian group Γ . Then H is generated as an algebra by Γ and its skew primitive elements.*

That is, we answer positively Conjecture 1 in a general context: when $G(H)$ is any abelian group. This Theorem is also applied to the known cases of finite-dimensional Nichols algebras over non-abelian groups

- Another crucial step of the Lifting Method is to obtain all deformations of the pointed Hopf algebras $\mathcal{B}(V)\#\mathbf{k}\Gamma$; that is, all the pointed Hopf algebras such that their associated coradically graded algebras are isomorphic to $\mathcal{B}(V)\#\mathbf{k}\Gamma$.

This problem was solved for Γ abelian in [AS4] – under the restriction that the order is not divisible by 2,3,5,7. We believe that the explicit presentation in this paper would be substancial to solve the question for any abelian group.

- The explicit relations would be also useful in the study of various elements in the representation theory of pointed Hopf algebras. In this direction, the theory of Nichols algebras of diagonal type provides an uniform approach to the study of quantum groups and quantum supergroups.

3. The plan of this paper is the following. We introduce the notion of Nichols algebras in Section 1. We give a PBW basis of any Nichols algebra and some properties of this basis following [Kh, R]. Next we recall the notions of Weyl groupoid and its associated root system following [CH2, HS], and make a connection with the theory of Nichols algebras of diagonal type. We present the needed material from [A2], in particular Theorem 1.25, a key result for our purposes.

Section 2 is devoted to Lusztig's isomorphisms in the general context of braidings of diagonal type [H4], extending analogous isomorphisms from [L].

In Section 3 we give the mentioned presentation by generators and relations, based in the classification of braidings of diagonal type with finite root system [H3]. The strategy of proof consists first to define Lusztig isomorphisms for the Drinfeld doubles of the braided Hopf algebras U^+ obtained by quotient by the relations in Theorem 1, except the group in (4). This quotient is analogous to the algebra $U_q^+(\mathfrak{g})$; the Drinfeld double $u_q(\mathfrak{g})$ of the Nichols algebra is a quotient of the previous algebra, as it was considered by Lusztig and Andruskiewitsch-Schneider. We denote these two algebras by U^+ and u^+ , respectively, so $u^+ = \mathcal{B}(V)$. The existence of the Lusztig's isomorphisms prove that the PBW generators corresponding to the algebras U^+ and their quotients u^+ are the same, but the heights of some generators are not the order of the associated scalar in U^+ . Therefore we obtain u^+ after to quotient U^+ by some powers of root vectors as in (4).

Theorem 3.1 extends the presentation obtained in [A1] for standard braidings, and in [AAY] for braidings of super type, and gives a new proof in the case of braidings of Cartan type, in particular quantized enveloping algebras $U_q(\mathfrak{g})$ and small quantum groups $u_q(\mathfrak{g})$.

Finally, Section 4 is devoted to the proof of Theorem 2. We prove first that any finite dimensional braided graded Hopf algebra of diagonal type

$$S = \bigoplus_{n \geq 0} S_n, \quad S_0 = \mathbf{k}1, \quad S_1 \cong V,$$

generated as an algebra by V is isomorphic to the Nichols algebra $\mathcal{B}(V)$; this result extends [AS4, Thm. 5.5], [AGI, Thm. 2.5], but the proof follows the same scheme.

Acknowledges. This work is part of the author's PhD Thesis. I want to thank specially to my advisor Nicolás Andruskiewitsch for his inspiring guidance, patience and supervision during these years. I want to thank also to my family for all their support, and to Antonela for all her love.

1. PRELIMINARIES

In this Section we recall results from different works needed in the sequel. First we consider the existence of PBW bases for Nichols algebras of diagonal type [Kh, R], and the rich combinatoric related to them. Next we recall the definitions of Weyl groupoid, the associated root systems and some properties thereof [HS, HY]. We close this Section stating a general presentation of Nichols algebras coming from [A2].

1.1. Lyndon words and PBW bases for Nichols algebras of diagonal type. To begin with, we recall the definition of a Nichols algebras and show a characterization in the case of a diagonal braiding.

Definition 1.1. [AS3] Given $V \in {}^H_H \mathcal{YD}$, the tensor algebra $T(V)$ admits a unique structure of graded braided Hopf algebra in ${}^H_H \mathcal{YD}$ such that $V \subseteq \mathcal{P}(V)$. Consider the family \mathfrak{S} of all the homogeneous Hopf ideals of $I \subseteq T(V)$ such that

- I is generated by homogeneous elements of degree ≥ 2 ,
- I is a Yetter-Drinfeld submodule of $T(V)$.

The *Nichols algebra* $\mathcal{B}(V)$ associated to V is the quotient of $T(V)$ by the biggest ideal $I(V)$ of \mathfrak{S} .

Let (V, c) be a braided vector space of diagonal type such that $q_{ij} = q_{ji}$ for any i, j . Let $\Gamma = \mathbb{Z}^\theta$, and $\alpha_1, \dots, \alpha_\theta$ be the canonical basis. We set the characters $\chi_1, \dots, \chi_\theta$ of Γ given by $\chi_j(\alpha_i) = q_{ij}$, $1 \leq i, j \leq \theta$.

Consider V as a Yetter-Drinfeld module over $\mathbf{k}\Gamma$ such that $x_i \in V_{\alpha_i}^{\chi_i}$. In this context we can characterize the Nichols algebra as a quotient that admits a certain non-degenerate bilinear form.

Proposition 1.2. [L, Prop. 1.2.3], [AS3, Prop. 2.10] *There exists a unique bilinear form $(\cdot|\cdot) : T(V) \times T(V) \rightarrow \mathbf{k}$ such that $(1|1) = 1$, and:*

- (1) $(x_i|x_j) = \delta_{ij}$, for any i, j ;
- (2) $(x|yy') = (x_1|y)(x_2|y')$, for any $x, y, y' \in T(V)$;
- (3) $(xx'|y) = (x|y_1)(x'|y_2)$, for any $x, x', y \in T(V)$.

This is a symmetric form, for which we have:

- (4) $(x|y) = 0$, for any $x \in T(V)_g$, $y \in T(V)_h$, $g, h \in \Gamma$, $g \neq h$.

The radical of this form $\{x \in T(V) : (x|y) = 0, \forall y \in T(V)\}$ coincides with $I(V)$, so $(\cdot|\cdot)$ induces a non-degenerate bilinear form on $\mathcal{B}(V) = T(V)/I(V)$, denoted also by $(\cdot|\cdot)$. \square

Therefore $I(V)$ is a \mathbb{Z}^θ -homogeneous ideal, and then $\mathcal{B}(V)$ is \mathbb{Z}^θ -graded.

Let A be an algebra, $P, S \subset A$ and $h : S \mapsto \mathbb{N} \cup \{\infty\}$. We fix a linear order $<$ on S . $B(P, S, <, h)$ will denote the set

$$\{p s_1^{e_1} \dots s_t^{e_t} : t \in \mathbb{N}_0, s_1 > \dots > s_t, s_i \in S, 0 < e_i < h(s_i), p \in P\}.$$

If $B(P, S, <, h)$ is a \mathbf{k} -linear basis, we say that $(P, S, <, h)$ is a set of *PBW generators*, whose *height* is h , and $B(P, S, <, h)$ is a *PBW basis* of A .

We will describe a particular PBW basis for any graded braided Hopf algebra $\mathcal{B} = \bigoplus_{n \in \mathbb{N}} \mathcal{B}^n$ generated by $\mathcal{B}^1 \cong V$ as an algebra, where V is a braided vector space; we will follow the results in [Kh].

Fix $\theta \in \mathbb{N}$, and a set $X = \{x_1, \dots, x_\theta\}$. Let \mathbb{X} be the set of words with letters in X and consider the lexicographical order on \mathbb{X} .

Definition 1.3. An element $u \in \mathbb{X}$, $u \neq 1$ is a *Lyndon word* if for any decomposition $u = vw$, $v, w \in \mathbb{X} - \{1\}$, we have $u < w$. We will denote the set of all Lyndon words by L .

Remark 1.4. • Each Lyndon word begin with its smaller letter.

- Each $u \in \mathbb{X} - X$ is a Lyndon word if and only if for each decomposition $u = u_1 u_2$ with $u_1, u_2 \in \mathbb{X} \setminus 1$, we have $u_1 u_2 = u < u_2 u_1$.
- If $u_1, u_2 \in L$ and $u_1 < u_2$, then $u_1 u_2 \in L$.

A basic Lyndon's result says that any word $u \in \mathbb{X}$ admits a unique decomposition as non-increasing product of Lyndon words:

$$(5) \quad u = l_1 l_2 \dots l_r, \quad l_i \in L, l_r \leq \dots \leq l_1.$$

It is called the *Lyndon decomposition* of $u \in \mathbb{X}$, and the $l_i \in L$ in (5) are called the *Lyndon letters* of u .

Another characterization of Lyndon words is the following:

Lemma 1.5. [Kh, p.6] *Let $u \in \mathbb{X} - X$. Then $u \in L$ if and only if there exist $u_1, u_2 \in L$ such that $u_1 < u_2$ and $u = u_1 u_2$. \square*

Definition 1.6. For each $u \in L - X$, the *Shirshov decomposition* of u is the decomposition $u = u_1 u_2$, $u_1, u_2 \in L$, such that u_2 is the smallest end of u between all the possible decompositions with these conditions.

Given a finite-dimensional vector space V , fix a basis $X = \{x_1, \dots, x_\theta\}$ de V ; we can identify $\mathbf{k}\mathbb{X}$ with $T(V)$. In what follows we consider two graduations for the algebra $T(V)$: the usual \mathbb{N}_0 -graduation $T(V) = \bigoplus_{n \geq 0} T^n(V)$, and \mathbb{Z}^θ -graduation of $T(V)$, determined by the condition $\deg x_i = \alpha_i$, $1 \leq i \leq \theta$, where $\{\alpha_1, \dots, \alpha_\theta\}$ is the canonical basis of \mathbb{Z}^θ .

Consider a braiding c for V . The *braided bracket* of $x, y \in T(V)$ is defined by

$$(6) \quad [x, y]_c := \text{multiplication} \circ (\text{id} - c)(x \otimes y).$$

Assume that (V, c) is of diagonal type, and let $\chi : \mathbb{Z}^\theta \times \mathbb{Z}^\theta \rightarrow \mathbf{k}^\times$ by the bicharacter determined by the condition

$$(7) \quad \chi(\alpha_i, \alpha_j) = q_{ij}, \quad \text{for each pair } 1 \leq i, j \leq \theta.$$

Then, for each pair of \mathbb{Z}^θ -homogeneous elements $u, v \in \mathbb{X}$,

$$(8) \quad c(u \otimes v) = q_{u,v} v \otimes u, \quad q_{u,v} = \chi(\deg u, \deg v) \in \mathbf{k}^\times.$$

In such case, the braided vector satisfies a "braided Jacobi identity" and determines skew-derivations as follows:

$$(9) \quad [[u, v]_c, w]_c = [u, [v, w]_c]_c - \chi(\alpha, \beta) v [u, w]_c + \chi(\beta, \gamma) [u, w]_c v,$$

$$(10) \quad [u, v w]_c = [u, v]_c w + \chi(\alpha, \beta) v [u, w]_c,$$

$$(11) \quad [u v, w]_c = \chi(\beta, \gamma) [u, w]_c v + u [v, w]_c,$$

where $u, v, w \in T(V)$ are homogeneous of degree $\alpha, \beta, \gamma \in \mathbb{N}^\theta$, respectively.

Using the previous decompositions, we can define the \mathbf{k} -linear endomorphism $[-]_c$ of $\mathbf{k}\mathbb{X}$ as follows:

$$[u]_c := \begin{cases} u, & \text{if } u = 1 \text{ or } u \in X; \\ [[v]_c, [w]_c]_c, & \text{if } u \in L, \ell(u) > 1, u = vw \text{ is the Shirshov decomposition}; \\ [u_1]_c \dots [u_t]_c, & \text{if } u \in \mathbb{X} - L \text{ and its Lyndon decomposition is } u = u_1 \dots u_t. \end{cases}$$

Definition 1.7. The *hyperletter* corresponding to $l \in L$ is $[l]_c$. An *hyperword* is a word whose letters are hyperletters, and a *monotone hyperword* is an hyperword $W = [u_1]_c^{k_1} \dots [u_m]_c^{k_m}$ such that $u_1 > \dots > u_m$.

Remark 1.8. For any $u \in L$, $[u]_c$ is a $\mathbb{Z}[q_{ij}]$ -linear combination of words with the same \mathbb{Z}^θ -graduation than u , such that $[u]_c \in u + \mathbf{k}\mathbb{X}_{>u}^{\ell(u)}$.

Theorem 1.9. [R, Thm. 10] *Let $u, v \in L$, $u < v$. Then $[[u]_c, [v]_c]_c$ is a $\mathbb{Z}[q_{ij}]$ -linear combination of monotone hyperwords $[l_1]_c \dots [l_r]_c$, $l_i \in L$, such that the corresponding hyperletters satisfy $v > l_i \geq uv$. Moreover, $[uv]_c$ appears in such combination with non-zero coefficient and each hyperword has the same \mathbb{Z}^θ -graduation than uv . \square*

The comultiplication of hyperwords in $T(V)$ has a nice expression, as we can see in the following result.

Lemma 1.10. [R, Thm.13] *Let $u_1, \dots, u_r, v \in L$, whit $v < u_r \leq \dots \leq u_1$. Then,*

$$\begin{aligned} \Delta([u_1]_c \cdots [u_r]_c [v]_c^m) &= 1 \otimes [u_1]_c \cdots [u_r]_c [v]_c^m + \sum_{i=0}^m \binom{m}{i}_{q_{v,v}} [u_1]_c \cdots [u_r]_c [v]_c^i \otimes [v]_c^{m-i} \\ &+ \sum_{\substack{l_1 \geq \dots \geq l_p > v, \\ 0 \leq j \leq m}} x_{l_1, \dots, l_p}^{(j)} \otimes [l_1]_c \cdots [l_p]_c [v]_c^j; \end{aligned}$$

Where $x_{l_1, \dots, l_p}^{(j)}$ is \mathbb{Z}^θ -homogeneous, and $\deg(x_{l_1, \dots, l_p}^{(j)}) + \deg(l_1 \dots l_p v^j) = \deg(u)$. \square

Another useful result from [R] is the following one.

Lemma 1.11. *For each $l \in L$ let W_l be the subspace of $T(V)$ generated by*

$$(12) \quad [l_1]_c [l_2]_c \cdots [l_k]_c, \quad k \in \mathbb{N}_0, l_i \in L, l_1 \geq \dots \geq l_k \geq l.$$

Then W_l is a left coideal subalgebra of $T(V)$. \square

We consider another order in \mathbb{X} as in [U]; it was implicitly used in [Kh]. Let $u, v \in \mathbb{X}$. We say that $u \succ v$ if $\ell(u) < \ell(v)$, or $\ell(u) = \ell(v)$ and $u > v$ for the lexicographical order. This order \succ is total, and it is called the *deg-lex order*.

The empty word 1 is the maximal element for \succ , and this order is invariant by left and right multiplication.

Let I be an ideal of $T(V)$, and $R = T(V)/I$. Let $\pi : T(V) \rightarrow R$ be the canonical projection. We set:

$$G_I := \{u \in \mathbb{X} : u \notin \mathbf{k}\mathbb{X}_{\succ u} + I\}.$$

Note that if $u \in G_I$ and $u = vw$, then $v, w \in G_I$. Therefore each $u \in G_I$ is a non-increasing product of Lyndon words of G_I .

Proposition 1.12. [Kh, R] *The set $\pi(G_I)$ is a basis of R .* \square

In what follows I will denote a Hopf ideal. Consider the set $S_I := G_I \cap L$. Define $h_I : S_I \rightarrow \{2, 3, \dots\} \cup \{\infty\}$ according to the following condition:

$$(13) \quad h_I(u) := \min \{t \in \mathbb{N} : u^t \in \mathbf{k}\mathbb{X}_{\succ u^t} + I\}.$$

We recall the following result and its corollaries following [Kh].

Theorem 1.13. $B'_I := B(\{1 + I\}, [S_I]_c + I, \langle, h_I)$ is a PBW basis of $H = T(V)/I$. \square

Corollary 1.14. *A word u belongs to G_I if and only if the corresponding hyperletter $[u]_c$ is not a linear combination, modulo I , of greater hyperwords $[w]_c$, $w \succ u$, whose hyperletters are in S_I .* \square

Corollary 1.15. *If $v \in S_I$ is such that $h_I(v) < \infty$, then $q_{v,v}$ is a root of unity. Moreover, if $\text{ord } q_{v,v} = h$, then $h_I(v) = h$, and $[v]_c^h$ is a linear combination of hyperwords $[w]_c$, $w \succ v^h$.* \square

1.2. Weyl groupoids and root systems. We follow the notation in [CH1]. Fix a non-empty set \mathfrak{X} , and a finite set I . For each $i \in I$ we fix a bijective function $r_i : \mathfrak{X} \rightarrow \mathfrak{X}$, and for each $X \in \mathfrak{X}$ a generalized Cartan matrix $A^X = (a_{ij}^X)_{i,j \in I}$.

Definition 1.16. [HY, CH1] The 4-uple $\mathcal{C} := \mathcal{C}(I, \mathfrak{X}, (r_i)_{i \in I}, (A^X)_{X \in \mathfrak{X}})$ is a *Cartan scheme* if it holds:

- for any $i \in I$, $r_i^2 = id$, and
- for any $X \in \mathfrak{X}$ and any pair $i, j \in I$: $a_{ij}^X = a_{ij}^{r_i(X)}$.

For each $i \in I$ and each $X \in \mathfrak{X}$ we denote by s_i^X the automorphism of \mathbb{Z}^I given by

$$s_i^X(\alpha_j) = \alpha_j - a_{ij}^X \alpha_i, \quad j \in I.$$

The *Weyl groupoid* of \mathcal{C} is the groupoid $\mathcal{W}(\mathcal{C})$ whose set of objects is \mathfrak{X} and whose morphisms are generated by s_i^X , considered as elements $s_i^X \in \text{Hom}(X, r_i(X))$, $i \in I$, $X \in \mathfrak{X}$.

In general we denote $\mathcal{W}(\mathcal{C})$ simply by \mathcal{W} , and for each $X \in \mathfrak{X}$:

$$(14) \quad \text{Hom}(\mathcal{W}, X) := \cup_{Y \in \mathfrak{X}} \text{Hom}(Y, X),$$

$$(15) \quad \Delta^{X \text{ re}} := \{w(\alpha_i) : i \in I, w \in \text{Hom}(\mathcal{W}, X)\}.$$

$\Delta^{X \text{ re}}$ is the set of *real roots* of X . Each $w \in \text{Hom}(\mathcal{W}, X_1)$ is written as a product $s_{i_1}^{X_1} s_{i_2}^{X_2} \cdots s_{i_m}^{X_m}$, where $X_j = r_{i_{j-1}} \cdots r_{i_1}(X_1)$, $i \geq 2$. We denote it by $w = \text{id}_{X_1} s_{i_1} \cdots s_{i_m}$: it means that $w \in \text{Hom}(\mathcal{W}, X_1)$, because each $X_j \in \mathfrak{X}$ is univocally determined by this condition. The *length* of w is defined by

$$\ell(w) = \min\{n \in \mathbb{N}_0 : \exists i_1, \dots, i_n \in I \text{ tales que } w = \text{id}_X s_{i_1} \cdots s_{i_n}\}.$$

We assume that \mathcal{W} is *connected*: that is, $\text{Hom}(Y, X) \neq \emptyset$, for any pair $X, Y \in \mathfrak{X}$.

Definition 1.17. [HY, CH1] Given a Cartan scheme \mathcal{C} , consider for each $X \in \mathfrak{X}$ a set $\Delta^X \subset \mathbb{Z}^I$. We say that $\mathcal{R} := \mathcal{R}(\mathcal{C}, (\Delta^X)_{X \in \mathfrak{X}})$ is a *root system of type \mathcal{C}* if

- (1) for any $X \in \mathfrak{X}$, $\Delta^X = (\Delta^X \cap \mathbb{N}_0^I) \cup -(\Delta^X \cap \mathbb{N}_0^I)$,
- (2) for any $i \in I$ and any $X \in \mathfrak{X}$, $\Delta^X \cap \mathbb{Z}\alpha_i = \{\pm\alpha_i\}$,
- (3) for any $i \in I$ and any $X \in \mathfrak{X}$, $s_i^X(\Delta^X) = \Delta^{r_i(X)}$,
- (4) if $m_{ij}^X := |\Delta^X \cap (\mathbb{N}_0\alpha_i + \mathbb{N}_0\alpha_j)|$, then $(r_i r_j)^{m_{ij}^X}(X) = (X)$ for any pair $i \neq j \in I$ and any $X \in \mathfrak{X}$.

$\Delta_+^X := \Delta^X \cap \mathbb{N}_0^I$ is called the set of *positive roots*, and $\Delta_-^X := -\Delta_+^X$ is the set of *negative roots*.

Remark 1.18. From (2) and (3) we deduce that $\Delta^{X \text{ re}} \subset \Delta^X$, for any $X \in \mathfrak{X}$.

For each positive root $\alpha = \sum_i n_i \alpha_i$, the *support* of α is the set

$$\text{supp } \alpha := \{i : 1 \leq i \leq \theta, n_i \neq 0\}.$$

By (3) we have that $w(\Delta^X) = \Delta^Y$ for any $w \in \text{Hom}(Y, X)$. We say that \mathcal{R} is *finite* if Δ^X is finite for some $X \in \mathfrak{X}$. By [CH1, Lemma 2.11], it is equivalent to the fact that all the sets Δ^X are finite, $X \in \mathfrak{X}$, and also that \mathcal{W} is finite. Moreover, for any pair $i \neq j \in I$ and any $X \in \mathfrak{X}$, we have that $k\alpha_i + \alpha_j \in \Delta^X$ if and only if $0 \leq k \leq -a_{ij}^X$. Therefore,

$$(16) \quad a_{ij}^X = -\max\{k \in \mathbb{N}_0 : k\alpha_i + \alpha_j \in \Delta^X\}.$$

A fundamental result involving root systems is the following one:

Theorem 1.19. [CH2, Thm. 2.10] *For every $\alpha \in \Delta_+^X \setminus \{\alpha_i : i = 1, \dots, \theta\}$, there exist $\beta, \gamma \in \Delta_+^X$ such that $\alpha = \beta + \gamma$. \square*

We give now some results about real roots and the length of elements.

Lemma 1.20. [HY, Cor. 3] *Let $m \in \mathbb{N}$, $X, Y \in \mathfrak{X}$, $i_1, \dots, i_m, j \in I$, $w = \text{id}_X s_{i_1} \cdots s_{i_m} \in \text{Hom}(Y, X)$, where $\ell(w) = m$. Then,*

- $\ell(ws_j) = m + 1$ if and only if $w(\alpha_j) \in \Delta_+^X$,
- $\ell(ws_j) = m - 1$ if and only if $w(\alpha_j) \in \Delta_-^X$. □

Proposition 1.21. [CH1, Prop. 2.12] *For any $w = \text{id}_X s_{i_1} \cdots s_{i_m} \in \mathcal{W}$ such that $\ell(w) = m$, the roots $\beta_j = s_{i_1} \cdots s_{i_{j-1}}(\alpha_{i_j}) \in \Delta^X$ are positive and all different. If \mathcal{R} is finite and w is an element of maximal length, then $\{\beta_j\} = \Delta_+^X$. Therefore all the roots are different: that is, for each $\alpha \in \Delta_+^X$ there exist $i_1, \dots, i_k, j \in I$ such that $\alpha = s_{i_k} \cdots s_{i_1}(x_j)$. □*

Call Δ_+^V the set of degrees of a PBW basis of $\mathcal{B}(V)$, counted with their multiplicities, as in [H2]. It does not depend on the PBW basis, see [H2, AA]. We can attach a Cartan scheme \mathcal{C} , a Weyl groupoid \mathcal{W} and a root system \mathcal{R} , see [HS, Thms. 6.2, 6.9]. To do this, define for each $1 \leq i \neq j \leq \theta$,

$$(17) \quad -a_{ij} := \min \{n \in \mathbb{N}_0 : (n+1)_{q_{ii}}(1 - q_{ii}^n \widetilde{q}_{ij}) = 0\},$$

and set $a_{ii} = 2$, $s_i \in \text{Aut}(\mathbb{Z}^\theta)$ such that $s_i(\alpha_j) = \alpha_j - a_{ij}\alpha_i$.

Set $\widetilde{q}_{rs} = \chi(s_i(\alpha_r), s_i(\alpha_s))$. Let V_i be another vector space of dimension θ , and attach to it the matrix $\widetilde{\mathbf{q}} = (\widetilde{q}_{rs})$. By [H2],

$$\Delta_+^{V_i} = s_i(\Delta_+^V \setminus \{\alpha_i\}) \cup \{\alpha_i\}.$$

If we consider $\Delta^V = \Delta_+^V \cup (-\Delta_+^V)$, last equation lets us to define the Weyl groupoid of V , whose root system is defined by the sets $\Delta^{V'}$, V' obtained after to apply some reflections to the matrix of V .

1.3. Defining relations of Nichols algebras of diagonal type.

Proposition 1.22. [A2, Prop. 3.1] *Assume that the braiding matrix is symmetric. Then a PBW basis of Lyndon hyperwords of $\mathcal{B}(V)$ is orthogonal with respect to the bilinear form in Proposition 1.2. □*

Corollary 1.23. [A2, Cor. 3.2] *If $u = x_{\beta_M}^{n_M} \cdots x_{\beta_1}^{n_1}$, where $0 \leq n_j < N_{\beta_j}$, then*

$$(18) \quad c_u := (u|u) = \prod_{j=1}^M n_j!_{q_{\beta_j}} c_{x_{\beta_j}}^{n_j} \neq 0.$$

□

Remark 1.24. Notice that:

$$(x_{\beta_i} x_{\beta_j} | u) = (x_{\beta_i} | u_{(1)})(x_{\beta_j} | u_{(2)}) = d_{i,j} c_{x_{\beta_i}} c_{x_{\beta_j}},$$

where $d_{i,j}$ is the coefficient $x_{\beta_i} \otimes x_{\beta_j}$ for the expression of $\Delta(u)$ in terms of the PBW basis (both factors of the tensor product).

For each pair $1 \leq i \leq j \leq \theta$, we denote

$$B_{ij} := \left\{ x_{\beta_j}^{n_j} \cdots x_{\beta_i}^{n_i} : 0 \leq n_k < N_{\beta_k} \right\};$$

that is, the set of hyperwords whose hyperletters are between x_{β_i} and x_{β_j}

Let (W, d) be a braided vector space of diagonal type that admits a basis $\widehat{x}_1, \dots, \widehat{x}_\theta$ such that, for some $\widehat{q}_{ij} \in \mathbf{k}^\times$, $d(\widehat{x}_i \otimes \widehat{x}_j) = \widehat{q}_{ij} \widehat{x}_j \otimes \widehat{x}_i$, where $\widehat{q}_{ij} = \widehat{q}_{ji}$, and (V, c) , (W, d) are twist equivalent:

$$\widetilde{q}_{ij} = \widehat{q}_{ij} \widehat{q}_{ji}, \quad q_{ii} = \widehat{q}_{ii}, \quad 1 \leq i \neq j \leq \theta.$$

Call $\widehat{x}_\beta = [l_\beta]_d$, the hyperletter corresponding to l_β for the braiding d . If $u = x_{\beta_M}^{n_M} \cdots x_{\beta_1}^{n_1}$, call also

$$\widehat{u} = \widehat{x}_{\beta_M}^{n_M} \cdots \widehat{x}_{\beta_1}^{n_1}.$$

Let $\sigma : \mathbb{Z}^\theta \times \mathbb{Z}^\theta \rightarrow \mathbf{k}^\times$ be the bicharacter determined by the condition

$$(19) \quad \sigma(g_i, g_j) = \begin{cases} \widehat{q}_{ij} q_{ij}^{-1}, & i \leq j \\ 1, & i > j \end{cases}$$

Define $t_{\alpha_i} = 1$ for any $1 \leq i \leq \theta$, and inductively,

$$t_\beta = \sigma(\beta_1, \beta_2) t_{\beta_1} t_{\beta_2}, \quad \text{Sh}(l_\beta) = (l_{\beta_1}, l_{\beta_2}).$$

For each $u = x_{\beta_M}^{n_M} \cdots x_{\beta_1}^{n_1}$ call

$$(20) \quad f(u) := \prod_{1 \leq i < j \leq M} \sigma(\beta_j, \beta_i)^{n_i n_j} \prod_{1 \leq i \leq M} \sigma(\beta_i, \beta_i)^{\binom{n_i}{2}} t_{\beta_i}^{n_i}.$$

Finally, for each pair $1 \leq i < j \leq \theta$ and $u = x_{\beta_M}^{n_M} \cdots x_{\beta_1}^{n_1}$, let

$$(21) \quad c_{i,j}^u := \frac{f(u) (\widehat{x}_{\beta_i} \widehat{x}_{\beta_j} | \widehat{u})}{\sigma(\beta_i, \beta_j) t_{\beta_i} t_{\beta_j} c_{\widehat{u}}},$$

where $(\cdot | \cdot)$ is the bilinear form corresponding to (W, d) , and $c_{\widehat{u}}$ is the scalar in Corollary 1.23. Such scalars let us to give a presentation by generators and relations as follows.

Theorem 1.25. [A2, Thm. 3.9] *Let (V, c) be a finite-dimensional braided vector space of diagonal type such that Δ_+^V is finite. Let x_1, \dots, x_θ be a basis of V such that $c(x_i \otimes x_j) = q_{ij} x_j \otimes x_i$, where $(q_{ij}) \in (\mathbf{k}^\times)^{\theta \times \theta}$ is the braiding matrix, and let $\{x_{\beta_k}\}_{\beta_k \in \Delta_+^V}$ be the set of hyperletters corresponding to the fixed order of the basis of V .*

Then $\mathcal{B}(V)$ is presented by generators x_1, \dots, x_θ , and relations

$$(22) \quad x_\beta^{N_\beta} = 0, \quad \beta \in \Delta_+^V, \quad \text{ord}(q_\beta) = N_\beta < \infty,$$

$$(23) \quad [x_{\beta_i}, x_{\beta_j}]_c = \sum_{u \in B_{ij} - \{x_{\beta_j}, x_{\beta_i}\}: \deg u = \beta_i + \beta_j} c_{i,j}^u u,$$

$$1 \leq i < j \leq M, \quad \text{Sh}(l_{\beta_i} l_{\beta_j}) = (l_{\beta_i}, l_{\beta_j}), \quad l_{\beta_i} l_{\beta_j} \neq l_{\beta_k}, \forall k,$$

where $c_{i,j}^u$ are as in (21). Moreover, $\{x_{\beta_M}^{n_M} \cdots x_{\beta_1}^{n_1} : 0 \leq n_j < N_{\beta_j}\}$ is a basis of $\mathcal{B}(V)$. \square

2. LUSZTIG ISOMORPHISMS OF NICHOLS ALGEBRAS OF DIAGONAL TYPE

In this Section we recall the Lusztig isomorphisms [H4] of Nichols algebras of diagonal type, which are a generalization of the isomorphisms of quantized enveloping algebras in [L]. We shall consider different quotients of the tensor algebra of a braided vector space of diagonal type and the Drinfeld doubles of their bosonizations by a free abelian group.

Notation: Let $\chi : \mathbb{Z}^\theta \times \mathbb{Z}^\theta \rightarrow \mathbf{k}^\times$ be a bicharacter, $q_{ij} = \chi(\alpha_i, \alpha_j)$. Then χ^{op} and χ^{-1} will denote the bicharacters:

$$\chi^{\text{op}}(\alpha, \beta) := \chi(\beta, \alpha), \quad \chi^{-1}(\alpha, \beta) := \chi(\alpha, \beta)^{-1}, \quad \alpha, \beta \in \mathbb{Z}^\theta.$$

Also, for any automorphism $s : \mathbb{Z}^\theta \rightarrow \mathbb{Z}^\theta$, $s^* \chi$ will denote the bicharacter defined by

$$(24) \quad (s^* \chi)(\alpha, \beta) := \chi(s^{-1}(\alpha), s^{-1}(\beta)), \quad \alpha, \beta \in \mathbb{Z}^\theta.$$

Let (V, c) a braided vector space of diagonal type, whose braiding matrix is (q_{ij}) . We consider $T(V)$ as an algebra in the category of Yetter-Drinfeld modules over $\mathbf{k}\mathbb{Z}^\theta$ as above. We follow the results in [H4, Section 4.1].

Definition 2.1. The Drinfeld double $\mathcal{U}(\chi)$ of the Hopf algebra $T(V)\#\mathbf{k}\mathbb{Z}^\theta$ is the algebra generated by elements $E_i, F_i, K_i^\pm, L_i^\pm, 1 \leq i \leq \theta$, and relations

$$\begin{aligned} XY &= YX, & X, Y &\in \{K_i^\pm, L_i^\pm : 1 \leq i \leq \theta\}, \\ K_i K_i^{-1} &= L_i L_i^{-1} = 1, \\ K_i E_j K_i^{-1} &= q_{ij} E_j, & L_i E_j L_i^{-1} &= q_{ji}^{-1} E_j, \\ K_i F_j K_i^{-1} &= q_{ij}^{-1} F_j, & L_i F_j L_i^{-1} &= q_{ji} F_j, \\ E_i F_j - F_j E_i &= \delta_{i,j} (K_i - L_i). \end{aligned}$$

It admits a Hopf algebra structure, where the comultiplication satisfies

$$\begin{aligned} \Delta(K_i) &= K_i \otimes K_i, & \Delta(E_i) &= E_i \otimes 1 + K_i \otimes E_i, \\ \Delta(L_i) &= L_i \otimes L_i, & \Delta(F_i) &= F_i \otimes L_i + 1 \otimes E_i, \end{aligned}$$

and then $\varepsilon(K_i) = \varepsilon(L_i) = 1, \varepsilon(E_i) = \varepsilon(F_i) = 0$.

Notice that $\mathcal{U}(\chi)$ is a \mathbb{Z}^θ -graded Hopf algebra, where the graduation is characterized by the following conditions:

$$\deg(K_i) = \deg(L_i) = 0, \quad \deg(E_i) = \alpha_i, \quad \deg(F_i) = -\alpha_i.$$

$\mathcal{U}^+(\chi)$ (respectively, $\mathcal{U}^-(\chi)$) denotes the subalgebra generated by E_i (respectively, F_i), $1 \leq i \leq \theta$, $\mathcal{U}^0(\chi)$ (respectively, $\mathcal{U}^{-0}(\chi)$) is the subalgebra generated by K_i (respectively, L_i), $1 \leq i \leq \theta$, and finally $\mathcal{U}^0(\chi)$ is the subalgebra generated by K_i and L_i . Note that $\mathcal{U}^0(\chi)$ is isomorphic to $\mathbf{k}\mathbb{Z}^{2\theta}$ as Hopf algebras. Moreover, the subalgebra generated by $\mathcal{U}^+(\chi)$ and $K_i, 1 \leq i \leq \theta$, is isomorphic to $T(V)\#\mathbf{k}\mathbb{Z}^\theta$, so $\mathcal{U}^+(\chi)$ is isomorphic to $T(V)$ as braided graded Hopf algebras in the category of Yetter-Drinfeld modules over $\mathbf{k}\mathbb{Z}^\theta$, where we consider the actions and coactions:

$$K_i \cdot E_i = q_{ij} E_i, \quad \delta(E_i) = K_i \otimes E_i.$$

We will consider a family of useful isomorphisms as in [H4, Section 4.1].

Proposition 2.2. (a) For any $\underline{a} = (a_1, \dots, a_\theta) \in (\mathbf{k}^\times)^\theta$ there exists a unique algebra automorphism $\varphi_{\underline{a}}$ of $\mathcal{U}(\chi)$ such that

$$(25) \quad \varphi_{\underline{a}}(K_i) = K_i, \quad \varphi_{\underline{a}}(L_i) = L_i, \quad \varphi_{\underline{a}}(E_i) = a_i E_i, \quad \varphi_{\underline{a}}(F_i) = a_i^{-1} F_i.$$

(b) There exists a unique algebra automorphism ϕ_1 of $\mathcal{U}(\chi)$ such that

$$(26) \quad \phi_1(K_i) = K_i^{-1}, \quad \phi_1(L_i) = L_i^{-1}, \quad \phi_1(E_i) = F_i L_i^{-1}, \quad \phi_1(F_i) = K_i^{-1} E_i.$$

(c) There exists a unique algebra isomorphism $\phi_2 : \mathcal{U}(\chi) \rightarrow \mathcal{U}(\chi^{-1})$ such that

$$(27) \quad \phi_2(K_i) = K_i, \quad \phi_2(L_i) = L_i, \quad \phi_2(E_i) = F_i, \quad \phi_2(F_i) = -E_i.$$

(d) There exists a unique Hopf algebra isomorphism $\phi_3 : \mathcal{U}(\chi) \rightarrow \mathcal{U}(\chi^{\text{op}})^{\text{cop}}$ such that

$$(28) \quad \phi_3(K_i) = L_i, \quad \phi_3(L_i) = K_i, \quad \phi_3(E_i) = F_i, \quad \phi_3(F_i) = E_i.$$

(e) There exists a unique algebra antiautomorphism ϕ_4 de $\mathcal{U}(\chi)$ such that

$$(29) \quad \phi_4(K_i) = K_i, \quad \phi_4(L_i) = L_i, \quad \phi_4(E_i) = F_i, \quad \phi_4(F_i) = E_i.$$

(f) Let $\underline{a} = (-1, \dots, -1)$. The antipode \mathcal{S} of $\mathcal{U}(\chi)$ is given by the composition $\mathcal{S} = \phi_1 \phi_4 \varphi_{\underline{a}}$. Also, $\phi_4^2 = \text{id}$. \square

As in [H4] we will consider some skew-derivations. $\underline{\Delta}$ will denote the braided comultiplication of $\mathcal{U}^+(\chi)$, which is \mathbb{N} -graded: if $E \in \mathcal{U}^+(\chi)$ is homogeneous of degree n , and $k \in \{0, 1, \dots, n\}$, $\underline{\Delta}_{n-k,k}(E)$ will denote the component of $\underline{\Delta}(E)$ in $\mathcal{U}^+(\chi)_{n-k} \otimes \mathcal{U}^+(\chi)_k$.

Proposition 2.3. *For any $i \in \{1, \dots, \theta\}$ there exist linear endomorphisms $\partial_i^K, \partial_i^L$ of $\mathcal{U}^+(\chi)$ such that*

$$EF_i - F_iE = \partial_i^K(E)K_i - L_i\partial_i^L(E) \quad \text{for all } E \in \mathcal{U}^+(\chi).$$

Such endomorphisms are given by:

$$\underline{\Delta}_{n-1,1}(E) = \sum_{i=1}^{\theta} \partial_i^K(E) \otimes E_i, \quad \underline{\Delta}_{1,n-1}(E) = \sum_{i=1}^{\theta} E_i \otimes \partial_i^L(E), \quad E \in \mathcal{U}^+(\chi)_n,$$

and satisfy the following conditions:

$$\begin{aligned} \partial_i^K(1) &= \partial_i^L(1) = 0, \\ \partial_i^K(E_j) &= \partial_i^L(E_j) = \delta_{i,j}, \\ \partial_i^K(E E') &= \partial_i^K(E)(K_i \cdot E') + E \partial_i^K(E'), \\ \partial_i^L(E E') &= \partial_i^L(E)E' + (L_i^{-1} \cdot E) \partial_i^L(E'), \end{aligned}$$

for all $j \in \{1, \dots, \theta\}$, and all pair of elements $E, E' \in \mathcal{U}^+(\chi)$. \square

We recall now a characterization of quotients of the algebra $\mathcal{U}(\chi)$ with a triangular decomposition [H4, Section 4.1]. According to [H4, Prop. 4.14], the multiplication

$$(30) \quad m : \mathcal{U}^+(\chi) \otimes \mathcal{U}^0(\chi) \otimes \mathcal{U}^-(\chi) \rightarrow \mathcal{U}(\chi)$$

is an isomorphism of \mathbb{Z}^θ -graded vector spaces.

Proposition 2.4. *Let $\mathcal{I}^+ \subset \mathcal{U}^+ \cap \ker \varepsilon$ (respectively, $\mathcal{I}^- \subset \mathcal{U}^- \cap \ker \varepsilon$) be an ideal of $\mathcal{U}^+(\chi)$ (respectively, $\mathcal{U}^-(\chi)$). The following conditions are equivalent:*

- The multiplication (30) induces an isomorphism

$$m : \mathcal{U}^+(\chi)/\mathcal{I}^+ \otimes \mathcal{U}^0(\chi) \otimes \mathcal{U}^-(\chi)/\mathcal{I}^- \rightarrow \mathcal{U}(\chi)/(\mathcal{I}^+ + \mathcal{I}^-).$$

- The vector spaces $\mathcal{I}^+\mathcal{U}^0(\chi)\mathcal{U}^-(\chi)$ y $\mathcal{U}^+(\chi)\mathcal{U}^0(\chi)\mathcal{I}^-$ are ideals of $\mathcal{U}(\chi)$.
- For all $X \in \mathcal{U}^0(\chi)$ and all $i \in \{1, \dots, \theta\}$ we have

$$\begin{aligned} X \cdot \mathcal{I}^+ &\subset \mathcal{I}^+, & \partial_i^K(\mathcal{I}^+) &\subset \mathcal{I}^+, & \partial_i^K(\phi_4(\mathcal{I}^-)) &\subset \phi_4(\mathcal{I}^-), \\ X \cdot \mathcal{I}^- &\subset \mathcal{I}^-, & \partial_i^L(\mathcal{I}^+) &\subset \mathcal{I}^+, & \partial_i^L(\phi_4(\mathcal{I}^-)) &\subset \phi_4(\mathcal{I}^-). \end{aligned}$$

\square

Lemma 2.5. [H4, Cor. 4.20] *Let \mathcal{I}^+ be an braided biideal of $\mathcal{U}^+(\chi)$, which is also a Yetter-Drinfeld $\mathcal{U}^0(\chi)$ -submodule and satisfies $\mathcal{I}^+ \subset \bigoplus_{n \geq 2} \mathcal{U}^+(\chi)_n$. Then $\mathcal{I}^+\mathcal{U}^0(\chi)\mathcal{U}^-(\chi)$ is a Hopf ideal of $\mathcal{U}(\chi)$. \square*

We will assume that all the integers $-a_{ij}$ of (17) associated with the braiding matrices (q_{ij}) are defined. Then we consider the automorphisms $s_{p,\chi} : \mathbb{Z}^\theta \rightarrow \mathbb{Z}^\theta$. We define also the scalars

$$(31) \quad \lambda_i(\chi) := (-a_{pi})_{q_{pp}} \prod_{s=0}^{-a_{pi}-1} (q_{pp}^s q_{pi} q_{ip} - 1), \quad i \neq p.$$

Denote by $\underline{E}_i, \underline{F}_i, \underline{K}_i, \underline{L}_i$ the generators corresponding to $\mathcal{U}(s_p^* \chi)$, and by $\underline{q}_{ij} = s_p^* \chi(\alpha_i, \alpha_j)$ the coefficients of the braiding matrix corresponding to $s_p^* \chi$.

Definition 2.6. We say that $p \in \{1, \dots, \theta\}$ is a *Cartan vertex* if it satisfies

$$q_{pp}^{m_{pj}} q_{pj} q_{jp} = 1, \quad \text{for every } j \neq p.$$

In such case, note that the existence of the integers m_{pj} implies that $\text{ord } q_{pp} \geq m_{pj} + 1$.

We denote by $\mathcal{O}(\chi)$ the union of the orbits of the simple roots α_p by the action of the Weyl groupoid, where p is a Cartan vertex.

Fix $p \in \{1, \dots, \theta\}$. For any $i \neq p$ we define as in [H4],

$$E_{i,0(p)}^+, E_{i,0(p)}^- := E_i, \quad F_{i,0(p)}^+, F_{i,0(p)}^- := F_i,$$

and recursively,

$$(32) \quad E_{i,m+1(p)}^+ := E_p E_{i,m(p)}^+ - (K_p \cdot E_{i,m(p)}^+) E_p = (\text{ad}_c E_p)^{m+1} E_i,$$

$$(33) \quad E_{i,m+1(p)}^- := E_p E_{i,m(p)}^- - (L_p \cdot E_{i,m(p)}^-) E_p,$$

$$(34) \quad F_{i,m+1(p)}^+ := F_p F_{i,m(p)}^+ - (L_p \cdot F_{i,m(p)}^+) F_p,$$

$$(35) \quad F_{i,m+1(p)}^- := F_p F_{i,m(p)}^- - (K_p \cdot F_{i,m(p)}^-) F_p.$$

When p is explicit, we simply denote $E_{i,m(p)}^\pm$ by $E_{i,m}^\pm$. By [H4, Cor. 5.4] the following identity holds for any $m \in \mathbb{N}_0$:

$$(36) \quad E_{i,m}^+ F_i - F_i E_{i,m}^+ = (m)_{q_{pp}} (q_{pp}^{m-1} q_{pi} q_{ip} - 1) L_p E_{i,m-1}^+.$$

Fix a braided graded Hopf algebra $\mathcal{B} \cong T(V)/I$, where I is a graded Hopf ideal generated by homogeneous elements of degree ≥ 2 . For each $1 \leq j \leq \theta$, $p \neq j$, we define

$$(37) \quad M_{p,j}^\pm(\mathcal{B}) := \left\{ E_{j,m}^\pm : m \in \mathbb{N}_0 \right\}.$$

In what follows we consider $\text{ord}(1) = 1$.

Remark 2.7. If $x_i^N = 0$ in \mathcal{B} , with N minimal (it is called the nilpotency order of x_i), then q_{ii} is a root of unity of order N . Moreover, $(\text{ad}_c x_i)^N x_j = 0$.

We recall a result from [A1] extending [H2, Prop. 1, Eqn. (18)].

Lemma 2.8. For each $p \in \{1, \dots, \theta\}$, let $\mathcal{B}_{\pm p}$ be the subalgebra generated by $\cup_{j \neq p} M_{p,j}^\pm(\mathcal{B})$, and denote $n_p = \text{ord}(q_{pp})$. There exist isomorphisms of graded vector spaces:

- $\ker(\partial_p^K) \cong \mathcal{B}_{+p} \otimes \mathbf{k}[E_p^{n_p}]$, $\ker(\partial_p^L) \cong \mathcal{B}_{-p} \otimes \mathbf{k}[E_p^{n_p}]$, if $1 < \text{ord}(q_{pp}) < \infty$ but E_p is not nilpotent, or
- $\ker(\partial_p^K) \cong \mathcal{B}_{+p}$, $\ker(\partial_p^L) \cong \mathcal{B}_{-p}$, if $\text{ord}(q_{pp})$ is the nilpotency order of E_p or $q_{pp} = 1$.

Moreover, the multiplication induces an isomorphism of graded vector spaces $\mathcal{B} \cong \mathcal{B}_p^\pm \otimes \mathbf{k}[x_p]$. \square

Set $N_p = \text{ord } q_{pp}$. We call, following [H4], $\mathcal{I}_p^+(\chi)$ (respectively, $\mathcal{I}_p^-(\chi)$) to the ideal of $\mathcal{U}^+(\chi)$ (respectively, $\mathcal{U}^-(\chi)$) generated by

- (a) $E_p^{N_p}$ (respectively, $F_p^{N_p}$), if p is not a Cartan vertex,
- (b) $E_{i,m_{pi}+1}^+$ (respectively, $F_{i,m_{pi}+1}^+$), for each i such that $N_p > m_{pi} + 1$.

Notice that $E_{i,m_{pi}+1}^+ \in \mathcal{I}_p^+(\chi)$ for any i such that $N_p = m_{pi} + 1$. We denote:

$$\mathcal{U}_p(\chi) := \mathcal{U}(\chi) / (\mathcal{I}_p^+(\chi) + \mathcal{I}_p^-(\chi)), \quad \mathcal{U}_p^+(\chi) := \mathcal{U}^+(\chi) / \mathcal{I}_p^+(\chi), \quad \mathcal{U}_p^-(\chi) := \mathcal{U}^-(\chi) / \mathcal{I}_p^-(\chi).$$

$I^+(\chi)$ will denote the ideal of $\mathcal{U}^+(\chi)$ such that the quotient $\mathcal{U}^+(\chi)/I^+(\chi)$ is isomorphic to the Nichols algebra of V ; that is, the greatest braided Hopf ideal of $\mathcal{U}^+(\chi)$ generated by elements of degree ≥ 2 . Call $I^-(\chi) = \phi_4(I^+(\chi))$, where ϕ_4 is defined by (29), and

$$\mathfrak{u}^\pm(\chi) := \mathcal{U}^\pm(\chi)/I^\pm(\chi), \quad \mathfrak{u}(\chi) := \mathcal{U}(\chi)/(I^-(\chi) + I^+(\chi)).$$

In such case, $\mathfrak{u}(\chi)$ is the Drinfeld double of the algebra $\mathfrak{u}^+(\chi) \# \mathbf{k}\mathbb{Z}^\theta$, where $\mathbf{k}\mathbb{Z}^\theta = \mathcal{U}^{+0}(\chi)$.

The Lusztig isomorphisms can be defined in this general context.

Theorem 2.9. [H4, Lemma 6.5, Theorem 6.12] *There exist algebra morphisms*

$$(38) \quad T_p, T_p^- : \mathcal{U}_p(\chi) \rightarrow \mathcal{U}_p(s_p^* \chi)$$

univocally determined by the following conditions:

$$\begin{aligned} T_p(K_p) &= T_p^-(K_p) = \underline{K}_p^{-1}, & T_p(K_i) &= T_p^-(K_i) = \underline{K}_p^{m_{pi}} \underline{K}_i, \\ T_p(L_p) &= T_p^-(L_p) = \underline{L}_p^{-1}, & T_p(L_i) &= T_p^-(L_i) = \underline{L}_p^{m_{pi}} \underline{L}_i, \\ T_p(E_p) &= \underline{E}_p \underline{L}_p^{-1}, & T_p(E_i) &= \underline{E}_{i, m_{pi}}^+, \\ T_p(F_p) &= \underline{K}_p^{-1} \underline{E}_p, & T_p(F_i) &= \lambda(s_p^* \chi)^{-1} \underline{F}_{i, m_{pi}}^+, \\ T_p^-(E_p) &= \underline{K}_p^{-1} \underline{E}_p, & T_p^-(E_i) &= \lambda(s_p^* \chi^{-1})^{-1} \underline{E}_{i, m_{pi}}^-, \\ T_p^-(F_p) &= \underline{E}_p \underline{L}_p^{-1}, & T_p^-(F_i) &= \underline{F}_{i, m_{pi}}^-. \end{aligned}$$

for every $i \neq p$. Both are isomorphisms satisfying

$$T_p T_p^- = T_p^- T_p = \text{id}, \quad T_p(\mathcal{U}_{+p}^+(\chi)) = \mathcal{U}_{-p}^+(s_p^* \chi).$$

Moreover, there exists $\lambda \in (\mathbf{k}^\times)^\theta$ such that

$$(39) \quad T_p \circ \phi_4 = \phi_4 \circ T_p^- \circ \varphi_\lambda.$$

Such isomorphisms induce algebra isomorphisms (denoted by the same name):

$$T_p, T_p^- : \mathfrak{u}(\chi) \rightarrow \mathfrak{u}(s_p^* \chi).$$

□

Remark 2.10. If the homogeneous elements $X, Y \in \mathcal{U}_p^+(\chi)$ are such that $T_p(X), T_p(Y) \in \mathcal{U}_p^+(s_p^* \chi)$, as $\deg T_p(X) = s_p(\deg X)$, it follows that

$$T_p([X, Y]_c) = [T_p(X), T_p(Y)]_c.$$

3. AN EXPLICIT PRESENTATION BY GENERATORS AND RELATIONS OF NICHOLS ALGEBRAS OF DIAGONAL TYPE

We shall obtain a family of isomorphisms induced by the ones in the previous Section. In this case we shall consider a quotient of $\mathcal{U}(\chi)$ by an ideal which is smaller than $(I^-(\chi) + I^+(\chi))$. Such ideal will be generated by some of the relations in Theorem 1.25, and will be the smallest one such that it is possible to define all the family of isomorphisms over the Weyl groupoid. It will give us a relation between the Hilbert series of these algebras, and new sets of roots. We shall use at the end the uniqueness of the root system, when the Weyl groupoid is finite.

We introduce some notation. We denote $\widetilde{q}_{ij} = q_{ij} q_{ji}$. Also,

$$x_{i_1 i_2 \dots i_k} = (\text{ad}_c x_{i_1}) \cdots (\text{ad}_c x_{i_{k-1}}) x_{i_k}, \quad i_j \in \{1, \dots, \theta\}.$$

For each $m \in \mathbb{N}$, we define the elements $x_{(m+1)\alpha_i + m\alpha_j} \in \mathcal{U}(\chi)$ recursively:

- if $m = 1$, $x_{2\alpha_i + \alpha_j} := (\text{ad}_c x_i)^2 x_j = x_{ii_j}$,
- $x_{(m+2)\alpha_i + (m+1)\alpha_j} := [x_{(m+1)\alpha_i + m\alpha_j}, x_{ij}]_c$.

We give now the main result of this section, which is Theorem 3.1: it gives an explicit presentation by generators and relations of any Nichols algebra of diagonal type with finite root system. We begin by proving several Lemmata to show the existence of Lusztig isomorphisms for some Hopf algebras. These Hopf algebras are intermediate between the tensor algebra and the Nichols algebras of a given braided vector space. Finally we use those Lusztig isomorphisms to prove the Theorem.

Theorem 3.1. *Let (V, c) be a finite-dimensional braided vector space of diagonal type, with braiding matrix $(q_{ij})_{1 \leq i, j \leq \theta}$, $\theta = \dim V$, and fix a basis x_1, \dots, x_θ of V such that $c(x_i \otimes x_j) = q_{ij} x_j \otimes x_i$. Let χ be the bicharacter associated to (q_{ij}) . Assume that the root system Δ^χ is finite. Then $\mathcal{B}(V)$ is presented by generators x_1, \dots, x_θ and relations:*

$$(40) \quad x_\alpha^{N_\alpha}, \quad \alpha \in \mathcal{O}(\chi);$$

$$(41) \quad (\text{ad}_c x_i)^{m_{ij}+1} x_j, \quad q_{ii}^{m_{ij}+1} \neq 1;$$

$$(42) \quad x_i^{N_i}, \quad i \text{ is not a Cartan vertex};$$

\diamond if $i, j \in \{1, \dots, \theta\}$ satisfy $q_{ii} = \widetilde{q}_{ij} = q_{jj} = -1$, and there exists $k \neq i, j$ such that $\widetilde{q}_{ik}^2 \neq 1$ or $\widetilde{q}_{jk}^2 \neq 1$,

$$(43) \quad x_{ij}^2;$$

\diamond if $i, j, k \in \{1, \dots, \theta\}$ satisfy $q_{jj} = -1$, $\widetilde{q}_{ik} = \widetilde{q}_{ij} \widetilde{q}_{jk} = 1$,

$$(44) \quad [x_{ijk}, x_j]_c;$$

\diamond if $i, j \in \{1, \dots, \theta\}$ satisfy $q_{jj} = -1$, $q_{ii} \widetilde{q}_{ij} \in \mathbb{G}_6$, and also $q_{ii} \in \mathbb{G}_3$ or $m_{ij} \geq 3$,

$$(45) \quad [x_{iij}, x_{ij}]_c;$$

\diamond if $i, j, k \in \{1, \dots, \theta\}$ satisfy $q_{ii} = \pm \widetilde{q}_{ij} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, and also $-q_{jj} = \widetilde{q}_{ij} \widetilde{q}_{jk} = 1$ or $q_{jj}^{-1} = \widetilde{q}_{ij} = \widetilde{q}_{jk} \neq -1$,

$$(46) \quad [x_{iijk}, x_{ij}]_c;$$

\diamond if $i, j, k \in \{1, \dots, \theta\}$ satisfy $\widetilde{q}_{ik}, \widetilde{q}_{ij}, \widetilde{q}_{jk} \neq 1$,

$$(47) \quad x_{ijk} - \frac{1 - \widetilde{q}_{jk}}{q_{kj}(1 - \widetilde{q}_{ik})} [x_{ik}, x_j]_c - q_{ij}(1 - \widetilde{q}_{jk}) x_j x_{ik};$$

\diamond if $i, j, k \in \{1, \dots, \theta\}$ satisfy one of the following situations

- \circ $q_{ii} = q_{jj} = -1$, $\widetilde{q}_{ij}^2 = \widetilde{q}_{jk}^{-1}$, $\widetilde{q}_{ik} = 1$, or
- \circ $\widetilde{q}_{ij} = q_{jj} = -1$, $q_{ii} = -\widetilde{q}_{jk}^2 \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, or
- \circ $q_{kk} = \widetilde{q}_{jk} = q_{jj} = -1$, $q_{ii} = -\widetilde{q}_{ij} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, or
- \circ $q_{jj} = -1$, $\widetilde{q}_{ij} = q_{ii}^{-2}$, $\widetilde{q}_{jk} = -q_{ii}^{-3}$, $\widetilde{q}_{ik} = 1$, or
- \circ $q_{ii} = q_{jj} = q_{kk} = -1$, $\pm \widetilde{q}_{ij} = \widetilde{q}_{jk} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$,

$$(48) \quad [[x_{ij}, x_{ijk}]_c, x_j]_c;$$

\diamond if $i, j, k \in \{1, \dots, \theta\}$ satisfy $q_{ii} = q_{jj} = -1$, $\widetilde{q}_{ij}^3 = \widetilde{q}_{jk}^{-1}$, $\widetilde{q}_{ik} = 1$,

$$(49) \quad [[[x_{ij}, [x_{ij}, x_{ijk}]_c]_c, x_j]_c;$$

\diamond if $i, j, k, l \in \{1, \dots, \theta\}$ satisfy $q_{jj} \widetilde{q}_{ij} = q_{jj} \widetilde{q}_{jk} = 1$, $\widetilde{q}_{jk}^2 = \widetilde{q}_{lk}^{-1} = q_{ll}$, $q_{kk} = -1$, $\widetilde{q}_{ik} = \widetilde{q}_{il} = \widetilde{q}_{jl} = 1$,

$$(50) \quad [[[x_{ijkl}, x_k]_c, x_j]_c, x_k]_c;$$

\diamond if $i, j, k \in \{1, \dots, \theta\}$ satisfy $q_{jj} = \widetilde{q}_{ij}^2 = \widetilde{q}_{jk} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$,

$$(51) \quad [[x_{ijk}, x_j]_c x_j]_c;$$

◇ if $i, j, k \in \{1, \dots, \theta\}$ satisfy $q_{jj} = \widetilde{q}_{ij}^3 = \widetilde{q}_{jk} \in \mathbb{G}_4$, $\widetilde{q}_{ik} = 1$,

$$(52) \quad [[x_{ijk}, x_j]_c, x_j]_c, x_j]_c;$$

◇ if $i, j, k \in \{1, \dots, \theta\}$ satisfy $q_{ii} = \widetilde{q}_{ij} = -1$, $q_{jj} = \widetilde{q}_{jk}^{-1} \neq -1$, $\widetilde{q}_{ik} = 1$,

$$(53) \quad [x_{ij}, x_{ijk}]_c;$$

◇ if $i, j, k \in \{1, \dots, \theta\}$ satisfy $q_{ii} = q_{kk} = -1$, $\widetilde{q}_{ij} = 1$, $\widetilde{q}_{ij} \in \mathbb{G}_3$, $q_{jj} = -\widetilde{q}_{jk} = \pm \widetilde{q}_{ij}$,

$$(54) \quad [x_i, x_{jjk}]_c - (1 + q_{jj}^2)q_{kj}^{-1} [x_{ijk}, x_j]_c - (1 + q_{jj}^2)(1 + q_{jj})q_{ij}x_jx_{ijk};$$

◇ if $i, j, k \in \{1, \dots, \theta\}$ satisfy $\widetilde{q}_{jk} = 1$, $q_{ii} = \widetilde{q}_{ij} = -\widetilde{q}_{ik} \in \mathbb{G}_3$,

$$(55) \quad [x_i, [x_{ij}, x_{ik}]_c]_c + q_{jk}q_{ik}q_{ji} [x_{iik}, x_{ij}]_c + q_{ij}x_{ij}x_{iik};$$

◇ if $i, j, k \in \{1, \dots, \theta\}$ satisfy $q_{jj} = q_{kk} = \widetilde{q}_{jk} = -1$, $q_{ii} = -\widetilde{q}_{ij} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$,

$$(56) \quad [x_{iik}, x_{ijk}]_c;$$

◇ if $i, j \in \{1, \dots, \theta\}$ satisfy $-q_{ii}, -q_{jj}, q_{ii}\widetilde{q}_{ij}, q_{jj}\widetilde{q}_{ij} \neq 1$,

$$(57) \quad (1 - \widetilde{q}_{ij})q_{jj}q_{ji} [x_i, [x_{ij}, x_j]_c]_c - (1 + q_{jj})(1 - q_{jj}\widetilde{q}_{ij})x_{ij}^2;$$

◇ if $i, j \in \{1, \dots, \theta\}$ satisfy $q_{jj} = -1$, $q_{ii}\widetilde{q}_{ij} \notin \mathbb{G}_6$, and also $m_{ij} \in \{4, 5\}$, or $m_{ij} = 3$, $q_{ii} \in \mathbb{G}_4$,

$$(58) \quad [x_i, x_{3\alpha_i+2\alpha_j}]_c - \frac{1 - q_{ii}\widetilde{q}_{ij} - q_{ii}^2\widetilde{q}_{ij}^2q_{jj}}{(1 - q_{ii}\widetilde{q}_{ij})q_{ji}}x_{ij}^2;$$

◇ if $i, j \in \{1, \dots, \theta\}$ satisfy $4\alpha_i + 3\alpha_j \notin \Delta_+^\chi$, $q_{jj} = -1$ or $m_{ji} \geq 2$, and also $m_{ij} \geq 3$, or $m_{ij} = 2$, $q_{ii} \in \mathbb{G}_3$,

$$(59) \quad x_{4\alpha_i+3\alpha_j} = [x_{3\alpha_i+2\alpha_j}, x_{ij}]_c;$$

◇ if $i, j \in \{1, \dots, \theta\}$ satisfy $3\alpha_i + 2\alpha_j \in \Delta_+^\chi$, $5\alpha_i + 3\alpha_j \notin \Delta_+^\chi$, and $q_{ii}^3\widetilde{q}_{ij}, q_{ii}^4\widetilde{q}_{ij} \neq 1$,

$$(60) \quad [x_{iij}, x_{3\alpha_i+2\alpha_j}]_c;$$

◇ if $i, j \in \{1, \dots, \theta\}$ satisfy $4\alpha_i + 3\alpha_j \in \Delta_+^\chi$, $5\alpha_i + 4\alpha_j \notin \Delta_+^\chi$,

$$(61) \quad x_{5\alpha_i+4\alpha_j} = [x_{4\alpha_i+3\alpha_j}, x_{ij}]_c;$$

◇ if $i, j \in \{1, \dots, \theta\}$ satisfy $5\alpha_i + 2\alpha_j \in \Delta_+^\chi$, $7\alpha_i + 3\alpha_j \notin \Delta_+^\chi$,

$$(62) \quad [[x_{iij}, x_{iij}], x_{iij}]_c;$$

◇ if $i, j \in \{1, \dots, \theta\}$ satisfy $q_{jj} = -1$, $5\alpha_i + 4\alpha_j \in \Delta_+^\chi$,

$$(63) \quad [x_{iij}, x_{4\alpha_i+3\alpha_j}]_c - \frac{b - (1 + q_{ii})(1 - q_{ii}\zeta)(1 + \zeta + q_{ii}\zeta^2)q_{ii}^6\zeta^4}{a q_{ii}^3q_{ij}^2q_{ji}^3}x_{3\alpha_i+2\alpha_j}^2,$$

where $\zeta = \widetilde{q}_{ij}$, $a = (1 - \zeta)(1 - q_{ii}^4\zeta^3) - (1 - q_{ii}\zeta)(1 + q_{ii})q_{ii}\zeta$, $b = (1 - \zeta)(1 - q_{ii}^6\zeta^5) - a q_{ii}\zeta$.

In what follows we will use implicitly the isomorphism $\mathcal{B}(V) \cong \mathfrak{u}^+(\chi)$ determined by $x_i \mapsto E_i$; in this way, we identify $\mathcal{B}(V)$ as a subalgebra of $\mathfrak{u}(\chi)$.

For any bicharacter χ whose root system is finite, $\mathcal{J}^+(\chi)$ denotes the ideal of $\mathcal{U}^+(\chi)$ generated by all the relations in Theorem 3.1, except (40), plus the quantum Serre relations $(\text{ad}_c x_i)^{1-a_{ij}}x_j$ for those vertices such that $q_{ii}^{a_{ij}} = q_{ij}q_{ji} = q_{ii}^{-1}$. The last ingredient is to obtain a quotient of all the algebras $\mathcal{U}_p(\chi)$, $1 \leq p \leq \theta$.

Call also $\mathcal{J}^-(\chi) := \phi_4(\mathcal{J}^+(\chi))$, $\mathcal{J}(\chi) := (\mathcal{J}^+(\chi) + \mathcal{J}^-(\chi))$,

$$U(\chi) := \mathcal{U}(\chi)/\mathcal{J}(\chi), \quad U^\pm(\chi) := \mathcal{U}^\pm(\chi)/\mathcal{J}^\pm(\chi).$$

We prove first that $\mathcal{J}^+(\chi)$ is contained in the ideal defining the corresponding Nichols algebra. The following Lemma was proved with Agustín García Iglesias and is implicit in other papers.

Lemma 3.2. *Let $I \subset T(V)$ be a braided homogeneous biideal of $T(V)$, so there exists a surjective morphism of braided graded Hopf algebras $\pi : R := T(V)/I \rightarrow \mathcal{B}(V)$. Let $\mathbf{x} \in \ker \pi$, $\mathbf{x} \neq 0$ of minimal degree $k \geq 2$. Then \mathbf{x} is primitive.*

Proof. As π is a morphism of graded braided bialgebras, $\ker \pi$ is a graded biideal:

$$\Delta(\mathbf{x}) = \mathbf{x} \otimes 1 + 1 \otimes \mathbf{x} + \sum_{j=1}^n b_j \otimes c_j \in \ker \pi \otimes R + R \otimes \ker \pi,$$

for some homogeneous elements $b_j, c_j \in \bigoplus_{i=1}^{k-1} R_i$, such that $\deg(b_j) + \deg(c_j) = k$. For each j we may assume either $b_j \in \ker \pi$ or $c_j \in \ker \pi$. If $b_j \in \ker \pi$, then $b_j = 0$ by the minimality condition on k . Similarly, if $c_j \in \ker \pi$, then $c_j = 0$. Hence \mathbf{x} is primitive in R . \square

We will work with \mathbb{N}_0^θ -graded ideals, so the following notation will be useful: given $\beta = \sum_i b_i \alpha_i$, $\gamma = \sum_i c_i \alpha_i$, for some $b_i, c_i \in \mathbb{N}_0$, we say that $\beta \geq \gamma$ (respectively, $\beta > \gamma$) if $b_i \geq c_i$ (respectively, $b_i > c_i$) for all $i \in \{1, \dots, \theta\}$.

Proposition 3.3. *$\mathcal{J}^+(\chi)$ is a braided biideal of $\mathcal{U}^+(\chi)$, and there exist a canonical projection of Hopf algebras $\pi_\chi : U(\chi) \rightarrow \mathbf{u}(\chi)$ such that $\pi(U^\pm(\chi)) = \mathbf{u}^\pm(\chi)$.*

Moreover, the multiplication $m : U^+(\chi) \otimes U^0(\chi) \otimes U^-(\chi) \rightarrow U(\chi)$ is an isomorphism of graded vector spaces.

Proof. We can order the relations according to their \mathbb{N} -graduation. When we quotient by the relations of degree $n - 1$, the relations of degree n are primitive by Lemma 3.2, because for any of them we can see that the relations in Theorem 1.25 of degree at most $n - 1$ are verified. Moreover, for a relation of degree $\alpha \in \mathbb{N}_0^\theta$, it is enough to verify that the relations of \mathbb{N}^θ -degree lower than α hold in this partial quotient. For example, each quantum Serre relation is primitive, and the same holds for $x_i^{\mathbb{N}_i}$; therefore, when we quotient by these relation we have that $\mathbf{x} = [(\text{ad}_c x_i)^2 x_j, (\text{ad}_c x_i) x_j]_c$ is primitive under the conditions for (45), because we have quotiented by x_i^3, x_j^2 , so it also holds that

$$(\text{ad}_c x_i)^3 x_j = (\text{ad}_c x_j)^2 x_i = 0.$$

We work in a similar way with the other relations so each partial quotient is a braided bialgebra (and then a Hopf algebra with the induced antipode); finally, $U^+(\chi)$ is a braided bialgebra, because $\mathcal{J}^+(\chi)$ is a braided biideal.

By the definition of Nichols algebra we conclude that $\mathcal{J}^+(\chi) \subseteq I^+(\chi)$. By Lemma 2.5, $\mathcal{J}^+(\chi) \mathcal{U}^0(\chi) \mathcal{U}^-(\chi)$ is a Hopf ideal of $\mathcal{U}(\chi)$, and then the equivalent conditions in Proposition 2.4 hold. Therefore there exists a projection of Hopf algebras and a triangular decomposition as in the Proposition. \square

Now we prove that the isomorphisms at the beginning of Theorem 2.9 induce isomorphisms between the corresponding algebras $U(\chi)$. The first step is to prove that $T_p(\mathcal{J}(\chi)) \subset \mathcal{J}(s_p^* \chi)$, which will be proved considering each relation generating the ideal. The following two Lemmata help us to reduce the number of explicit computations.

Lemma 3.4. *Let l be a Lyndon word such that $[l]_c \cong \sum_{w \in S_{I^+(\chi)}, w \succ l} a_w w \pmod{I^+(\chi)}$, for some $a_w \in \mathbf{k}$. Let I be a braided Hopf ideal \mathbb{N}^θ -graded of $\mathcal{U}^+(\chi)$ such that the set of good words $S_{I^+(\chi)}, S_I$ coincide for those terms $w \succ l$, and assume that l is written as a linear combination of words greater than l modulo I . Then $[l]_c \cong \sum_{w \in S_{I^+(\chi)}, w \succ l} a_w w \pmod{I}$.*

Proof. It is a direct consequence of Corollary 1.14: by this result, $[l]_c$ is written as a linear combination of good hyperwords greater than $[l]_c$ modulo I . Such hyperwords coincide with the corresponding good hyperwords for $I^+(\chi)$ by hypothesis, and also $I \subseteq I^+(\chi)$. Hence the linear combination should be the same, because the good hyperwords generate a linear complement of the ideal in $\mathcal{U}^+(\chi)$. \square

Lemma 3.5. *Let I be an ideal \mathbb{N}_0^θ -homogeneous of $T(V)$, $\theta = \dim V$. Let S, T be two minimal sets of homogeneous generators of I . Assume that for each $\alpha \in \mathbb{N}^\theta$ there exists at most one generator in S (respectively in T) of degree α , and denote by $I(S, \alpha)$ (respectively, $I(T, \alpha)$) the ideal generated by the elements of S (respectively, T) of degree $\beta < \alpha$.*

For each $s \in S$ of degree $\alpha \in \mathbb{N}_0^\theta$, there exists $t \in T$ of the same degree, and $c \in \mathbf{k}^\times$ such that $s \cong c t \pmod{(I(S, \alpha))}$, and then $I(S, \alpha) = I(T, \alpha)$.

Proof. We prove it by induction on the degree of the generators. Let s be of degree α , minimal for the partial order defined on \mathbb{N}_0^θ . Therefore $\dim I_\alpha = 1$, so there exists an element of T which belongs to this subspace of I of dimension 1.

If the degree of s is not minimal, we apply inductive hypothesis for all the generators in lower degree, so for each s' of lower degree there exists a corresponding $t' \in T$ of the same degree which satisfies the conditions above, and $I(S, \alpha) = I(T, \alpha)$. Therefore

$$\dim I(S, \alpha)_\alpha = \dim I(T, \alpha)_\alpha = \dim I_\alpha - 1,$$

because S is a minimal set of generators, and by hypothesis there exists a unique generator of degree α . As T is also a set of generators of I , there exists $t \in I - I(T, \alpha) = I - I(S, \alpha)$, of degree α , so the statement follows. \square

Remark 3.6. This Lemma lets us to identify relations of the same degree for two sets of minimal generators of an ideal, up to relations of lower degree and scalars. In this way we can consider relations from Theorem 1.25 for a fixed order on the letters, and consider relations for another order. If we have a minimal set and this set contains relations all in different degrees (as we will have for the set of relations of the Nichols algebra or some partial quotients), then we can find a correspondence as above between the relations of the same \mathbb{Z}^θ -degree.

For example, if $q_{ii}^{m_{ij}+1} \neq 1$ for some pair of vertices i, j , then the quantum Serre relation $(\text{ad}_c x_i)^{m_{ij}+1} x_j$ is a generator for the minimal set of generators corresponding to the order $x_i < x_j$, so for the order $x_i > x_j$ we have:

$$[x_j x_i^{m_{ij}+1}]_c = [\cdots [(\text{ad}_c x_j) x_i, x_i]_c, \cdots, x_i]_c = a (\text{ad}_c x_i)^{m_{ij}+1} x_j,$$

for some scalar $a \in \mathbf{k}^\times$.

Also, if $q_{ii} \in \mathbb{G}_3$, $\widetilde{q}_{ij} \in \{\pm q_{ii}, -1\}$, $q_{jj} = -1$, we notice that

$$[(\text{ad}_c x_i)^2 x_j, (\text{ad}_c x_i) x_j]_c \cong b [(\text{ad}_c x_j) x_i, [(\text{ad}_c x_j) x_i, x_i]_c]_c \pmod{I},$$

for some $b \in \mathbf{k}^\times$, where I is the ideal generated by x_i^3 and x_j^2 , because such relations correspond to different minimal sets of generators of the ideal of relations of the Nichols algebra, and these are the generators of degree $3\alpha_i + 2\alpha_j$ for each set.

Lemma 3.7. *Let I be a \mathbb{Z}^θ -graded ideal of $\mathcal{U}_p^+(\chi)$. Let $Y, Z \in \mathcal{U}_p^+(\chi)/I$ be homogeneous elements of degree $\beta, \gamma \in \mathbb{N}_0^\theta$, respectively, such that $(\text{ad}_c E_p)Y = 0$. Then,*

$$(64) \quad [(\text{ad}_c E_p)Z, Y]_c = (\text{ad}_c E_p)[Z, Y]_c.$$

If also $\chi(\alpha_p, \beta)\chi(\beta, \alpha_p) = 1$, then

$$(65) \quad \chi(\alpha_p, \beta)[Y, (\text{ad}_c E_p)Z]_c = (\text{ad}_c E_p)[Y, Z]_c.$$

Proof. Both identities follow from (9). For example, for the second one,

$$\begin{aligned} (\operatorname{ad}_c E_p) [Y, Z]_c &= [E_p, [Y, Z]_c]_c = [[E_p, Y]_c, Z]_c + \chi(\alpha_p, \beta) Y [E_p, Z]_c - \chi(\beta, \gamma) [E_p, Z]_c Y \\ &= \chi(\alpha_p, \beta) (Y (\operatorname{ad}_c E_p) Z - \chi(\beta, \gamma) \chi(\beta, \alpha_p) (\operatorname{ad}_c E_p) Z Y) \\ &= \chi(\alpha_p, \beta) [Y, (\operatorname{ad}_c E_p) Z]_c, \end{aligned}$$

where we use the condition $\chi(\alpha_p, \beta) \chi(\beta, \alpha_p) = 1$. \square

Lemma 3.8. *Let $i, p \in \{1, \dots, \theta\}$ be such that $m_{pi} \geq 2$ and $m_{ip} = 1$. Then, in $U(s_p^* \chi)$,*

$$\left[\underline{E}_{i, m_{pi}}^+, \underline{E}_{i, m_{pi}-1}^+ \right]_c = [(\operatorname{ad}_c \underline{E}_p)^{m_{pi}} \underline{E}_i, (\operatorname{ad}_c \underline{E}_p)^{m_{pi}-1} \underline{E}_i]_c = 0.$$

Remark 3.9. Such relation belongs to the ideal $I^+(s_p^* \chi)$. In fact, as $2\alpha_i + \alpha_p \notin \Delta_+^\chi$, we have $s_p(2\alpha_i + \alpha_p) = 2\alpha_i + (2m_{pi} - 1)\alpha_p \notin \Delta_+^\chi$, so such relation holds in the corresponding Nichols algebra $\mathfrak{u}^+(s_p^* \chi)$.

On the other hand, some of these relations are generators of the ideal $\mathcal{J}(s_p^* \chi)$ by definition, for example (45). We prove here that the other relations not in the definition of this ideal are redundant; that is, they are generated by relations of lower degree.

Proof. We consider the different possible values of m_{pi} ; we begin with $m_{pi} = 2$. Therefore $q_{pp} \in \mathbb{G}_3$ or $q_{pp}^2 q_{ip} q_{pi} = 1$, and also $q_{ii} = -1$ or $q_{ii} q_{ip} q_{pi} = 1$. If $\underline{m}_{ip} = 1$ for $s_p^* \chi$, then p, i determine a subdiagram of standard type. If $q_{pp}^3 \neq 1$ or $q_{ii} \neq -1$ then $E_p^2 E_i E_p E_i$ is written as a linear combination of words greater than $E_p^2 E_i E_p E_i$, modulo $\mathcal{J}(s_p^* \chi)$, using the quantum Serre relations, because in the first case $E_p^2 E_i E_p$ appears with non-zero coefficient in $(\operatorname{ad}_c E_p)^3 E_i$, so $E_p^2 E_i E_p E_i$ is a linear combination of greater words and $E_p^3 E_i^2$, but this last word is in the ideal if $q_{ii} = -1$, or $E_p E_i^2$ appears in $(\operatorname{ad}_c E_i)^2 E_p$ with non-zero coefficient, so in both cases we obtain $E_p^2 E_i E_p E_i$ as a combination of greater words. Using Lemma 3.4, we conclude that $\left[\underline{E}_{i,2}^+, \underline{E}_{i,1}^+ \right]_c = 0$. A similar proof in the case $q_{pp}^3 = 1$, $q_{ii} \neq -1$ gives us the same conclusion. If $q_{pp}^3 = 1$, $q_{ii} = -1$, the relation corresponds to (45), which is by definition a generator of $\mathcal{J}(s_p^* \chi)$.

If $m_{pi} = 2$ and $\underline{m}_{ip} > 1$ for $s_p^* \chi$, then (57) is a generator of $\mathcal{J}(s_p^* \chi)$, and then $E_p E_i E_p E_i$ is a linear combination of $E_p^2 E_i^2$ and greater words. Therefore $\left[\underline{E}_{i,2}^+, \underline{E}_{i,1}^+ \right]_c \in \mathcal{J}(s_p^* \chi)$, by a similar argument.

If $m_{pi} = 3$, then $\underline{m}_{ip} = 1$ for $s_p^* \chi$, or there exists $\zeta \in \mathbb{G}_{24}$ such that $q_{pp} = \zeta^6$, $q_{ii}^{-1} = q_{pi} q_{ip} = \zeta$. For the first case, we notice that (58) holds also if $q_{pp} \notin \mathbb{G}_4$, because $\underline{E}_p^3 \underline{E}_i \underline{E}_p$ can be written as a linear combination of other words from the quantum Serre relation $(\operatorname{ad}_c \underline{E}_p)^4 \underline{E}_i = 0$, and then $\underline{E}_p^3 \underline{E}_i \underline{E}_p \underline{E}_i$ is a linear combination of greater words multiplying by \underline{E}_i , so we apply Lemma 3.4; from this relation we work as above, so we write $E_p^2 E_i E_p^2 E_i$ as a linear combination of other words and deduce that $E_p^3 E_i E_p^2 E_i$ is a linear combination of greater words, and we can apply Lemma 3.4 again. For the second case, we write $\underline{E}_p^3 \underline{E}_i \underline{E}_p^2 \underline{E}_i$ as a linear combination of greater words using the quantum Serre relations or the relation (58), with the same conclusion.

If $m_{pi} = 4, 5$, then $\underline{m}_{ip} = 1$ for $s_p^* \chi$. Therefore we work as before and we obtain the desired relation from (58) or (45), according to $3\alpha_p + 2\alpha_i$ belongs to $\Delta_+^{s_p^* \chi}$ or not. In both situations, we can write $\underline{E}_p^4 \underline{E}_i \underline{E}_p^3 \underline{E}_i$ or $\underline{E}_p^5 \underline{E}_i \underline{E}_p^4 \underline{E}_i$ as a linear combination of greater words, so we apply Lemma 3.4 again. \square

We will prove now that $T_p(x) \in \mathcal{J}(s_p^* \chi)$ for any generator x of the ideal $\mathcal{J}^+(\chi)$ so we will have a family of morphisms between the elements the algebras $U(\chi)$.

Lemma 3.10. *Let i be a non-Cartan vertex. Then $T_p(E_i^{N_i}) \in \mathcal{J}(s_p^*\chi)$.*

If i, j satisfy $q_{ii} = \widetilde{q}_{ij} = q_{jj} = -1$, and there exists k such that $\widetilde{q}_{ik}^2 \neq -1$ or $\widetilde{q}_{jk}^2 \neq -1$, then $T_p(E_{ij}^2) \in \mathcal{J}(s_p^\chi)$.*

Proof. Consider the first relation. If $i = p$, then p is not Cartan for χ , so p is not Cartan for $s_p^*\chi$ too. Therefore, by the definition of the ideal $\mathcal{J}(s_p^*\chi)$,

$$T_p(E_p^{N_p}) = \underline{E}_p^{N_p} = \phi_4(\underline{E}_p^{N_p}) \in \mathcal{J}(s_p^*\chi).$$

We consider then $i \neq p$. In such case, $T_p(E_i^{N_i}) = \left(\underline{E}_{i, m_{pi}}^+\right)^{N_i}$.

If $m_{pi} = 0$, then $\underline{E}_{i,0}^+ = \underline{E}_i$ and $q_{ip}q_{pi} = 1$, so for each $j \neq p$ we have $q_{ij}q_{ji} = \widetilde{q}_{ij}$.

Therefore i is not Cartan for $s_p^*\chi$, and $T_p(E_i^{N_i}) = \underline{E}_i^{N_i} \in \mathcal{J}(s_p^*\chi)$.

Consider $m_{pi} \neq 0$. As i is not Cartan, there exists $j \neq i$ such that $q_{ii}^{m_{ij}} \widetilde{q}_{ij} \neq 1$.

Assume first that $m_{ip} + 1 = N_i$. If $m_{ip} = 1$, that is $q_{ii} = -1$, there are two possibilities. If $q_{ip}q_{pi} \neq -1$, using Lemma 3.8, the identity (9) and the quantum Serre relation $(\text{ad}_{\underline{c}} \underline{E}_p)^{m_{pi}+1} \underline{E}_i = 0$, we compute in $U(s_p^*\chi)$,

$$\begin{aligned} 0 &= \left[\underline{E}_p, \left[\underline{E}_{i, m_{pi}}^+, \underline{E}_{i, m_{pi}-1}^+ \right]_{\underline{c}} \right]_{\underline{c}} \\ &= (s_p^*\chi(\alpha_p, m_{pi}\alpha_p + \alpha_i) - s_p^*\chi(m_{pi}\alpha_p + \alpha_i, (m_{pi}-1)\alpha_p + \alpha_i)) \left(\underline{E}_{i, m_{pi}}^+\right)^2 \\ &= (\chi(-\alpha_p, \alpha_i) - \chi(\alpha_i, \alpha_p + \alpha_i)) \left(\underline{E}_{i, m_{pi}}^+\right)^2 = q_{pi}^{-1}(1 + q_{ip}q_{pi}) \left(\underline{E}_{i, m_{pi}}^+\right)^2, \end{aligned}$$

so $T_p(E_i^2) = \left(\underline{E}_{i, m_{pi}}^+\right)^2 \in \mathcal{J}(s_p^*\chi)$. If $q_{pi}q_{ip} = -1$, there are 3 possible subdiagrams determined by i, p : it is standard with $q = -1$, or it is Cartan of type B_2 with $q \in \mathbb{G}_4$, or it is Cartan of type G_2 with $q \in G_6$. For the first case, if the diagram is of type A_2 associated to $q = -1$, it follows by definition of the ideal $\mathcal{J}(s_p^*\chi)$; for the other cases, we write $\underline{E}_p^{m_{pi}} \underline{E}_i \underline{E}_p^{m_{pi}} \underline{E}_i$ as a linear combination of greater words using (45) or the quantum Serre relations, and also the previous Lemmata.

If $m_{ip} > 1$ and $m_{pi} = 1$, we compute, using (9) and the relation $(\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_i = 0$,

$$\begin{aligned} \text{ad}_{\underline{c}} \underline{E}_p \left[\underline{E}_{i,1}^+, \underline{E}_i \right]_{\underline{c}} &= (s_p^*\chi(\alpha_p, \alpha_i + \alpha_p) - s_p^*\chi(\alpha_i + \alpha_p, \alpha_i)) \left(\underline{E}_{i,1}^+\right)^2 \\ &= q_{pi}^{-1}(1 - q_{ii}q_{ip}q_{pi}) \left(\underline{E}_{i,1}^+\right)^2. \end{aligned}$$

From this relation and (9) again, we calculate

$$\begin{aligned} \text{ad}_{\underline{c}} \underline{E}_p \left[\underline{E}_{i,1}^+, \left[\underline{E}_{i,1}^+, \underline{E}_i \right]_{\underline{c}} \right]_{\underline{c}} &= (s_p^*\chi(\alpha_p, \alpha_i + \alpha_p) - s_p^*\chi(\alpha_i + \alpha_p, 2\alpha_i + \alpha_p)) \\ &\quad (q_{pi}^{-1} - q_{ii}q_{ip}) \left(\underline{E}_{i,1}^+\right)^3 \\ &= q_{pi}^{-2}(1 - q_{ii}q_{ip}q_{pi})(1 - q_{ii}^2q_{ip}q_{pi}) \left(\underline{E}_{i,1}^+\right)^3. \end{aligned}$$

So if $m_{ip} = 2$ and $m_{pi} = 1$, it follows that $\alpha_p + 3\alpha_i \notin \Delta_+^\chi$, and

$$s_p(\alpha_p + 3\alpha_i) = 3\alpha_i + 2\alpha_p \notin \Delta_+^{s_p^*\chi}.$$

Using the previous Lemma, $[(\text{ad}_{\underline{c}} \underline{E}_i)^2 \underline{E}_p, (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_p]_{\underline{c}} \in \mathcal{J}^+(s_p^*\chi)$, so

$$\left[\underline{E}_{i,1}^+, \left[\underline{E}_{i,1}^+, \underline{E}_i \right]_{\underline{c}} \right]_{\underline{c}} \in \mathcal{J}(s_p^*\chi),$$

because we apply Lemma 3.5 if the relation belongs to a minimal set of generators ($q_{ii}^2 q_{ip} q_{pi} \neq 1$, so $q_{ii} \in \mathbb{G}_3$), or we compute it directly for the cases in which it is Cartan of type B_2 or standard with $q_{pp} = -1$. Then, by a similar argument,

$$T_p(E_i^3) = \left(\underline{E}_{i,1}^+ \right)^3 \in \mathcal{J}(s_p^* \chi).$$

If $m_{ip} = 3$, $m_{pi} = 1$, we have that $s_p(\alpha_p + 4\alpha_i) = 4\alpha_i + 3\alpha_p \notin \Delta_+^{s_p^* \chi}$, so

$$\left[\underline{E}_{i,1}^+, \left[\underline{E}_{i,1}^+, \left[\underline{E}_{i,1}^+, \underline{E}_i \right]_{\underline{c}} \right]_{\underline{c}} \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi),$$

by a similar argument, using (59). In this case we deduce that $\left(\underline{E}_{i,1}^+ \right)^4 \in \mathcal{J}(s_p^* \chi)$.

If $m_{ip} = 4$, then $q_{ii}^4 q_{ip} q_{pi} \neq 1$, so $\left(\underline{E}_{i,1}^+ \right)^5 \in \mathcal{J}(s_p^* \chi)$ in a similar way, using (61). We notice that there are no diagrams such that $q_{ii}^{m_{ip}+1} = 1$ and $m_{ip} \geq 5$.

Now we consider $m_{ip}, m_{pi} > 1$, so there are 3 possibilities:

- $m_{ip} = m_{pi} = 2$, so (59) is a generator of $\mathcal{J}(s_p^* \chi)$, and $q_{pp} \in \mathbb{G}_3$. Therefore we write $\underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i$ as a linear combination of other words, which begin with \underline{E}_p or they contain \underline{E}_p^3 as a factor. If we multiply by \underline{E}_p^2 on the left, $\underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i$ is a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, because $\underline{E}_p^3 \in \mathcal{J}(s_p^* \chi)$, so $T_p(E_i^3) = ((\text{ad}_c \underline{E}_p)^2 \underline{E}_i)^3 \in \mathcal{J}(s_p^* \chi)$.
- $m_{ip} = 3$, $m_{pi} = 2$, so $q_{ii} = \zeta^6$, $q_{pp} = \zeta^8$, $q_{ip} q_{pi} = \zeta^{11}$ for some $\zeta \in \mathbb{G}_{24}$, and (63) is a generator of the ideal $\mathcal{J}(s_p^* \chi)$. Using this relation we write $\underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i$ as a linear combination of words beginning with \underline{E}_p or words containing \underline{E}_p^3 as a factor. Multiplying by \underline{E}_p^2 on the left, we write $(\underline{E}_p^2 \underline{E}_i)^4$ as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, because $\underline{E}_p^3 \in \mathcal{J}(s_p^* \chi)$, so $T_p(E_i^4) = ((\text{ad}_c \underline{E}_p)^2 \underline{E}_i)^4 \in \mathcal{J}(s_p^* \chi)$.
- $m_{ip} = 2$, $m_{pi} = 3$, so there are two possible diagrams; in both cases (62) is a generator of $\mathcal{J}(s_p^* \chi)$. From this relation we write $\underline{E}_i \underline{E}_p^3 \underline{E}_i \underline{E}_p^3 \underline{E}_i$ as a combination of words beginning with \underline{E}_p or containing \underline{E}_p^4 . Multiplying on the left by \underline{E}_p^3 , $\underline{E}_p^3 \underline{E}_i \underline{E}_p^3 \underline{E}_i \underline{E}_p^3 \underline{E}_i$ can be written as a linear combination of greater words, modulo $\mathcal{J}(s_p^* \chi)$, because $\underline{E}_p^4 \in \mathcal{J}(s_p^* \chi)$ or $(\text{ad}_c \underline{E}_p)^4 \underline{E}_i \in \mathcal{J}(s_p^* \chi)$, so, in both cases, $T_p(E_i^3) = ((\text{ad}_c \underline{E}_p)^3 \underline{E}_i)^3 \in \mathcal{J}(s_p^* \chi)$.

Finally we consider $q_{ii}^{m_{ip}} q_{ip} q_{pi} = 1$, $m_{ip} < N_i - 1$, so there exists $j \neq p$ such that $1 \leq m_{ip} < m_{ij} = N_i - 1$. In this case, i, j, p determine a connected diagram, where i is not Cartan, connected with j and p , and also q_{ii} is a root of unity of order $N_i > 2$. We have the following possible diagrams under the previous conditions:

- $q_{ii} \in \mathbb{G}_3$, $q_{pp} \in \{q_{ii}, -1\}$, $m_{ip} = m_{pi} = 1$, $m_{ij} = 2$, $m_{pj} = m_{jp} = 0$, or
- $q_{ii} \in \mathbb{G}_4$, $q_{pp} = -1$, $q_{ip} q_{pi} q_{ii} = 1$, $\tilde{q}_{ij} = q_{ii}$, $m_{pj} = m_{jp} = 0$, $m_{ij} = 3$ (a diagram of type super $G(3)$, with $q \in \mathbb{G}_4$).

Both possibilities follow in a similar way to the case $m_{pi} = 1$.

We analyze now the second relation. As $\tilde{q}_{ij} = -1$,

$$((\text{ad}_c E_j) E_i)^2 = q_{ji}^2 ((\text{ad}_c E_i) E_j)^2 + 2q_{ji} (E_i E_j^2 E_i + E_j E_i^2 E_j).$$

By the first part $T_p(E_i^2), T_p(E_j^2) \in \mathcal{J}(s_p^* \chi)$, and as T_p is an algebra morphism, it is enough to prove that $T_p\left(\left((\text{ad}_c E_i) E_j\right)^2\right) \in \mathcal{J}(s_p^* \chi)$, to conclude that also $T_p\left(\left((\text{ad}_c E_j) E_i\right)^2\right) \in$

$\mathcal{J}(s_p^*\chi)$, and vice versa. Moreover we need just one of these two relations in order to have a minimal set of relations.

If $p = j$, we have

$$\begin{aligned} T_p \left(((\text{ad}_c E_i)E_p)^2 \right) &= \left(\underline{E}_{i,1}^+ \underline{E}_p \underline{L}_p^{-1} - q_{ip} \underline{E}_p \underline{L}_p^{-1} \underline{E}_{i,1}^+ \right)^2 \\ &= \left(\left(\underline{E}_p \underline{E}_{i,1}^- 2 \underline{L}_p \underline{E}_i \right) \underline{L}_p^{-1} - q_{ip} q_{pp} q_{ip} \underline{E}_p \underline{E}_{i,1}^+ \underline{L}_p^{-1} \right)^2 \\ &= \left(-2 \underline{q}_{ip}^{-1} \underline{E}_i \right)^2 = 4 \underline{q}_{ip}^2 \underline{E}_i^2 \in \mathcal{J}(s_p^*\chi), \end{aligned}$$

because $\underline{q}_{pp} = \underline{q}_{pi} \underline{q}_{ip} = \underline{q}_{ii} = -1$.

Now we consider $p \neq i, j$. If $m_{pi}, m_{pj} \neq 0$, we have two possibilities:

- $q_{pp} = -1$, $q_{ip} q_{pi} q_{jp} q_{pj} = -1$ so it is a diagram of type super $D(2, 1; \alpha)$, or
- $q_{pp} = q_{ip}^{-1} q_{pi}^{-1} = -q_{jp} q_{pj} \in \mathbb{G}_3 \cup \mathbb{G}_4 \cup \mathbb{G}_6$.

For the first case, or the second when $q_{pp} \in \mathbb{G}_4$,

$$T_p \left(((\text{ad}_c E_i)E_j)^2 \right) = [(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_j]_{\underline{c}}^2.$$

Using (47) and \underline{E}_p^2 if $\underline{q}_{pp} = -1$, or the quantum Serre relations

$$(\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_i = (\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_j = 0$$

if $q_{pp} \in \mathbb{G}_4$, $\underline{E}_i \underline{E}_p \underline{E}_j \underline{E}_p$ is a linear combination of greater words, so $(\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_j)^2$ is also a linear combination of greater words. Then,

$$[(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_j]_{\underline{c}}^2 \in \mathcal{J}^+(s_p^*\chi),$$

by an analogous statement to Lemma 3.4 but for powers of hyperwords, and such relation is in $I^+(s_p^*\chi)$.

For the remaining cases, $q_{pp} \in \mathbb{G}_3 \cup \mathbb{G}_6$ and

$$T_p \left(((\text{ad}_c E_i)E_j)^2 \right) = [(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_j]_{\underline{c}}^2.$$

We write $(\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_j)^2$ as a linear combination of greater words using the quantum Serre relations or (47), so

$$T_p \left(((\text{ad}_c E_i)E_j)^2 \right) \in \mathcal{J}^+(s_p^*\chi)$$

by an analogous argument.

If $m_{pi} = 1$, $m_{pj} = 0$, we have two possibilities. If $q_{pp} = -1$, then $\widetilde{q}_{pi} \neq -1$ and $\underline{q}_{ii} = \widetilde{q}_{pi} = \widetilde{q}_{pi}^{-1}$, so (53) is a generator of $\mathcal{J}(s_p^*\chi)$, as well as \underline{E}_p^2 and \underline{E}_{pj} . By (9), we have that

$$q_{pi} q_{pj} (1 + \widetilde{q}_{pi}) \underline{E}_{pij}^2 = [\underline{E}_{ppi}, \underline{E}_{ij}]_{\underline{c}} - [\underline{E}_p, [\underline{E}_{pij}, \underline{E}_{ij}]_{\underline{c}}]_{\underline{c}} \in \mathcal{J}(s_p^*\chi),$$

so $T_p(\underline{E}_{ij}^2) = \underline{E}_{pij}^2 \in \mathcal{J}(s_p^*\chi)$. If not, then $q_{pp}^{-1} = \widetilde{q}_{pi} \neq -1$, and we write $\underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_p \underline{E}_i \underline{E}_j$ as a linear combination of greater words modulo $\mathcal{J}(s_p^*\chi)$, using the quantum Serre relations (observe that (\underline{q}_{ij}) is twist equivalent to the original braiding). Therefore $T_p(\underline{E}_{ij}^2) = \underline{E}_{pij}^2 \in \mathcal{J}(s_p^*\chi)$.

If $m_{pi} > 1$, $m_{pj} = 0$, then $q_{pp} = -\widetilde{q}_{pi} \in \mathbb{G}_3$, and the proof follows in a similar way to the case $q_{pp} = -1$, but using the relation (56).

If $m_{pi} = m_{pj} = 0$, the proof follows easily, because $\underline{q}_{ii} = \underline{q}_{ij} \underline{q}_{ji} = \underline{q}_{jj} = -1$, and $T_p(\underline{E}_{ij}^2) = \underline{E}_{ij}^2 \in \mathcal{J}(s_p^*\chi)$ by definition of the ideal. \square

Lemma 3.11. *Let $i, j \in \{1, \dots, \theta\}$ be such that $q_{ii}^{m_{ij}} \widetilde{q}_{ij} = 1$. Then*

$$T_p((\text{ad}_c E_i)^{m_{ij}+1} E_j) \in \mathcal{J}(s_p^* \chi).$$

Proof. (i) The case $p = i$ was considered in the first part of Theorem 2.9.

(ii) Let $p = j$: we analyze all the possible values of m_{ip} . If $m_{ip} = 0$, then $q_{ip} q_{pi} = 1$, and

$$T_p((\text{ad}_c E_i) E_p) = \underline{E}_i \underline{F}_p \underline{L}_p^{-1} - q_{ip} \underline{F}_p \underline{L}_p^{-1} \underline{E}_i = (\underline{E}_i \underline{F}_p - \underline{F}_p \underline{E}_i) \underline{L}_p^{-1} \in \mathcal{J}(s_p^* \chi).$$

CONSIDER $m_{ip} = 1$; by (36) we have

$$\begin{aligned} T_p((\text{ad}_c E_i) E_p) &= \underline{E}_{i, m_{pi}}^+ \underline{F}_p \underline{L}_p^{-1} - q_{ip} \underline{F}_p \underline{L}_p^{-1} \underline{E}_{i, m_{pi}}^+ \\ &= \left(\underline{F}_p \underline{E}_{i, m_{pi}}^+ + (m_{pi})_{q_{pp}} (q_{pp}^{m_{pi}-1} q_{pi} q_{ip} - 1) \underline{L}_p \underline{E}_{i, m_{pi}-1}^+ \right) \underline{L}_p^{-1} \\ &\quad - q_{ip} s_p^+ \chi(m_{pi} \alpha_p + \alpha_i, \alpha_p) \underline{F}_p \underline{E}_{i, m_{pi}}^+ \underline{L}_p^{-1} \\ &= (m_{pi})_{q_{pp}} (q_{pp}^{m_{pi}-1} q_{pi} q_{ip} - 1) s_p^* \chi((m_{pi}-1) \alpha_p + \alpha_i, \alpha_p)^{-1} \underline{E}_{i, m_{pi}-1}^+ \\ &\quad + \underline{F}_p \underline{E}_{i, m_{pi}}^+ \underline{L}_p^{-1} - q_{ip} \chi(\alpha_i, -\alpha_p) \underline{F}_p \underline{E}_{i, m_{pi}}^+ \underline{L}_p^{-1} \\ (66) \quad &= (m_{pi})_{q_{pp}} (q_{pp}^{-1-m_{pi}} q_{pi}^{-1} q_{ip}^{-1} - 1) q_{ip} q_{pp} \underline{E}_{i, m_{pi}-1}^+. \end{aligned}$$

If $m_{pi} = 1$, we have by this identity and Remark 2.10:

$$T_p((\text{ad}_c E_i)^2 E_p) = T_p([E_i, (\text{ad}_c E_i) E_p]_c) = [(\text{ad}_c \underline{E}_p) \underline{E}_i, a_1 \underline{E}_i]_c,$$

where $a_{m_{pi}} := (m_{pi})_{q_{pp}} (q_{pp}^{-1-m_{pi}} q_{pi}^{-1} q_{ip}^{-1} - 1) q_{ip} q_{pp} \in \mathbf{k}^\times$. This element is in $\mathcal{J}(s_p^* \chi)$ because $\underline{m}_{ip} = 1$, so $(\text{ad}_c \underline{E}_i)^2 \underline{E}_p \in \mathcal{J}(s_p^* \chi)$. We consider now $m_{pi} \geq 2$; by Lemma 3.8,

$$T_p((\text{ad}_c E_i)^2 E_p) = \left[\underline{E}_{i, m_{pi}}^+, a_{m_{pi}} \underline{E}_{i, m_{pi}-1}^+ \right]_c \in \mathcal{J}(s_p^* \chi).$$

CONSIDER NOW $m_{ip} = 2$. We look at all the possible diagrams with two vertices and note that $m_{pi} = 1$, or there exists $\zeta \in \mathbb{G}_9$ such that $q_{ii} = -\zeta$, $q_{ip} q_{pi} = \zeta^7$, $q_{pp} = \zeta^3$. In the first case, $q_{pp} \in \{-1, q_{ii}^2\}$, so this diagram is standard of type B_2 , and (45) belongs to $\mathcal{J}(s_p^* \chi)$ by Lemma 3.8. Therefore

$$T_p((\text{ad}_c E_i)^3 E_p) = a_1 \left[(\text{ad}_c \underline{E}_p) \underline{E}_i, [(\text{ad}_c \underline{E}_p) \underline{E}_i, \underline{E}_i]_c \right]_c \in \mathcal{J}(s_p^* \chi).$$

For the second case, the braiding matrix of $s_p^* \chi$ is $q_{ii} = -1$, $q_{ip} q_{pi} = \zeta^8$, $q_{pp} = \zeta^3$. Then

$$T_p((\text{ad}_c E_i)^3 E_p) = \left[(\text{ad}_c \underline{E}_p)^2 \underline{E}_i, [(\text{ad}_c \underline{E}_p)^2 \underline{E}_i, (\text{ad}_c \underline{E}_p) \underline{E}_i]_c \right]_c \in \mathcal{J}(s_p^* \chi),$$

because (60) is a generator of this ideal.

IT REMAINS TO CONSIDER $m_{ip} \in \{3, 4, 5\}$. The unique diagram with $m_{pi} > 1$ is

$$\circ - \zeta \xrightarrow{-\zeta^{12}} \circ \zeta^5,$$

where $\zeta \in \mathbb{G}_{15}$, $q_{ii} = -\zeta$ and $m_{ip} = 3$, $m_{pi} = 2$; applying s_p we obtain

$$\circ^{-1} \xrightarrow{-\zeta^{13}} \circ \zeta^5.$$

By (61) we write $\underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p \underline{E}_i$ as a linear combination of words beginning with \underline{E}_p , or containing \underline{E}_p^3 as a factor, or greater than this word for the order $p < i$. Multiplying

on the left by \underline{E}_p^2 and using that $\underline{E}_p^3 \in \mathcal{J}(s_p^*\chi)$, $(\underline{E}_p^2 \underline{E}_i)^3 \underline{E}_p \underline{E}_i$ can be written as a linear combination of greater words, modulo $\mathcal{J}(s_p^*\chi)$. By Lemma 3.4 we conclude that

$$T_p((\text{ad}_c E_i)^4 E_p) = \left[\underline{E}_{i,2}^+, \left[\underline{E}_{i,2}^+, \left[\underline{E}_{i,2}^+, \underline{E}_{i,1}^+ \right]_{\underline{c}} \right]_{\underline{c}} \right]_{\underline{c}} = [(\underline{E}_p^2 \underline{E}_i)^3 \underline{E}_p \underline{E}_i]_{\underline{c}} \in \mathcal{J}(s_p^*\chi).$$

Finally we consider $m_{pi} = 1$, so we have

$$\begin{aligned} s_p(m_{ip}\alpha_i + \alpha_p) &= m_{ip}\alpha_i + (m_{pi} - 1)\alpha_p \in \Delta_+^{s_p^*\chi}, \\ s_p((m_{ip} + 1)\alpha_i + \alpha_p) &= (m_{ip} + 1)\alpha_i + m_{pi}\alpha_p \notin \Delta_+^{s_p^*\chi}. \end{aligned}$$

If $q_{pp} = \underline{q}_{pp} \neq -1$ then $m_{pi} = 3$ and $(\underline{E}_p \underline{E}_i)^2 \underline{E}_p \underline{E}_i^2$ is a linear combination of greater words modulo $\mathcal{J}(s_p^*\chi)$, where we use first the quantum Serre relation $(\text{ad}_c \underline{E}_p)^2 \underline{E}_i = 0$ to write $\underline{E}_p \underline{E}_i \underline{E}_p$ as a combination of the words $\underline{E}_p^2 \underline{E}_i$, $\underline{E}_i \underline{E}_p^2$ and then (58), which also holds in $U(s_p^*\chi)$. By this relation,

$$T_p((\text{ad}_c E_i)^4 E_p) = \left[\underline{E}_{i,1}^+, \left[\underline{E}_{i,1}^+, \left[\underline{E}_{i,1}^+, \underline{E}_i \right]_{\underline{c}} \right]_{\underline{c}} \right]_{\underline{c}} \in \mathcal{J}(s_p^*\chi).$$

In other case, $q_{pp} = -1$ and $m_{ip} \in \{3, 4, 5\}$, so we also have that

$$T_p((\text{ad}_c E_i)^{m_{ip}+1} E_p) = [\underline{E}_{m_{pi}\alpha_i + (m_{pi}-1)\alpha_p}, (\text{ad}_c \underline{E}_p) \underline{E}_i]_{\underline{c}} \in \mathcal{J}(s_p^*\chi),$$

by (59), (61) or (63), depending on the value of m_{ip} .

(iii) Let $p \neq j$: if $m_{pi} = 0$ (i.e. $\widetilde{q}_{ip} = 1$), then $q_{ii} = q_{jj}$, $q_{ij}q_{ji} = \widetilde{q}_{ij}$, so $\underline{m}_{ij} = m_{ij}$, and $(\text{ad}_c \underline{E}_i)^{m_{ij}+1} \underline{E}_j = 0$ holds in $U(s_p^*\chi)$. Moreover, in $U(s_p^*\chi)$ we have $\underline{E}_p \underline{E}_i = \underline{q}_{pi} \underline{E}_p \underline{E}_i$, so

$$(\text{ad}_c \underline{E}_i)(\text{ad}_c \underline{E}_p) X = \underline{q}_{ip} (\text{ad}_c \underline{E}_p)(\text{ad}_c \underline{E}_i) X,$$

for any $X \in U(s_p^*\chi)$, by the second part of Lemma 3.7. By Remark 2.10 and the previous results, in $U(s_p^*\chi)$ we have

$$\begin{aligned} T_p((\text{ad}_c E_i)^{m_{ij}+1} E_j) &= (\text{ad}_c \underline{E}_i)^{m_{ij}+1} (\text{ad}_c \underline{E}_p)^{m_{pj}} \underline{E}_j \\ &= \underline{q}_{ip}^{m_{pj}(m_{ij}+1)} (\text{ad}_c \underline{E}_p)^{m_{pj}} (\text{ad}_c \underline{E}_i)^{m_{ij}+1} \underline{E}_j = 0. \end{aligned}$$

Consider now $m_{pi} \neq 0$. If $m_{ij} = m_{pj} = 0$, we apply Lemma 3.7 to obtain

$$T_p((\text{ad}_c E_i) E_j) = [(\text{ad}_c \underline{E}_p)^{m_{pi}} \underline{E}_i, \underline{E}_j]_{\underline{c}} = (\text{ad}_c \underline{E}_p)^{m_{pi}} \left([\underline{E}_i, \underline{E}_j]_{\underline{c}} \right) = 0.$$

where we use that $q_{ij}q_{ji} = \widetilde{q}_{ij} = 1$, so in $U(s_p^*\chi)$ it holds that $[\underline{E}_i, \underline{E}_j]_{\underline{c}} = 0$. It remains to consider the case in which i, j and p determine a connected diagram, and $m_{pi} \neq 0$.

FIRST WE ANALYZE THE CASE $m_{ij} = 0$, $m_{pj} \neq 0$. If $q_{pp} = -1$ it follows that $m_{pi} = m_{pj} = 1$. Then $q_{ij}q_{ji} = q_{ip}q_{pi}q_{jp}q_{pj}$, and $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_j$ is a linear combination of greater words for the order $p < i < j$, modulo $\mathcal{J}(s_p^*\chi)$:

- if $q_{ij}q_{ji} = 1$, it follows from (44),
- if $q_{ij}q_{ji} \neq 1$, we write $\underline{E}_i \underline{E}_p \underline{E}_j$ as a linear combination of other words by (47), where those words are greater than $\underline{E}_i \underline{E}_p \underline{E}_j$ or begin with \underline{E}_p , so we multiply on the left by \underline{E}_p and use that $\underline{E}_p^2 \in \mathcal{J}(s_p^*\chi)$.

In this way, $T_p((\text{ad}_c E_i) E_j) = [(\text{ad}_c \underline{E}_p) \underline{E}_i, (\text{ad}_c \underline{E}_p) \underline{E}_j]_{\underline{c}} \in \mathcal{J}(s_p^*\chi)$. If $m_{pi} = m_{pj} = 1$ and $q_{pp} \neq -1$, then $(\text{ad}_{cb} \underline{E}_p)^2 \underline{E}_i, (\text{ad}_{cb} \underline{E}_p)^2 \underline{E}_j \in \mathcal{J}(s_p^*\chi)$; by these relations and $(\text{ad}_c \underline{E}_i) \underline{E}_j, \underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_j$ can be written as a linear combination of greater words for the order $p < i < j$, modulo $\mathcal{J}(s_p^*\chi)$, and also $T_p((\text{ad}_c E_i) E_j) \in \mathcal{J}(s_p^*\chi)$ in this case.

If $m_{pj} = 1$ and $m_{pi} > 1$ (or analogously, $m_{pj} > 1$, $m_{pi} = 1$), then $q_{pp}q_{pj}q_{jp} = 1$, and $q_{pp} \neq -1$. Note that

$$\widetilde{q}_{ij} = s_p^* \chi(\alpha_i, \alpha_j) s_p^* \chi(\alpha_j, \alpha_i) = q_{pp}^{m_{pi}} q_{pi} q_{ip}.$$

If $q_{pp}^{m_{pi}} q_{pi} q_{ip} \neq 1$, then (47) holds in $U(s_p^* \chi)$, so we can write $\underline{E}_i \underline{E}_p \underline{E}_j$ as a linear combination of other words, greater than $\underline{E}_i \underline{E}_p \underline{E}_j$ for the order $p < i < j$, or beginning with \underline{E}_p . Multiplying on the left by $\underline{E}_p^{m_{pi}}$, we express $\underline{E}_p^{m_{ij}} \underline{E}_i \underline{E}_p \underline{E}_j$ as a linear combination of greater words, using that $\underline{E}_p^{m_{pi}+1} \in \mathcal{J}(s_p^* \chi)$, or $(\text{ad}_{\underline{c}} \underline{E}_p)^{m_{pi}+1} \underline{E}_i \in \mathcal{J}(s_p^* \chi)$, so

$$T_p((\text{ad}_c E_i) E_j) = [(\text{ad}_{\underline{c}} \underline{E}_p)^{m_{pi}} \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_j]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

If $q_{pp}^{m_{pi}} q_{pi} q_{ip} = 1$ and $q_{pp}^{m_{pi}+1} \neq 1$, $\underline{E}_p^{m_{ij}} \underline{E}_i \underline{E}_p \underline{E}_j$ is written as a linear combination of greater words for the same order using $(\text{ad}_{\underline{c}} \underline{E}_p)^{m_{pi}+1} \underline{E}_i$, $(\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_j$ and $(\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j$, so we obtain the same conclusion. If $q_{pp}^{m_{pi}} q_{pi} q_{ip} = 1$ and $q_{pp}^{m_{pi}+1} = 1$, then $m_{pi} = 2$ or $m_{pi} = 3$, and the conclusion follows from (51) or (52), respectively.

If $m_{pi}, m_{pj} > 1$, there is only one possibility: $m_{pi} = m_{pj} = 2$. The proof is as above, expressing $\underline{E}_p^2 \underline{E}_i \underline{E}_p \underline{E}_j$ as a linear combination of greater words in the two possible cases: if $q_{pp} \notin \mathbb{G}_3$, using the quantum Serre relations; if $q_{pp} \in \mathbb{G}_3$, by (55) and \underline{E}_p^3 .

WE CONSIDER NOW $m_{pj} = 0$, $m_{ij} \neq 0$. Note that $m_{ij} \leq 3$, because we have a connected diagram with three vertices and $q_{ii} \neq -1$: $q_{ii}^{m_{ij}+1} \neq 1$. If $m_{ij} = 3$, it corresponds to a diagram of type super $G(3)$:

$$\chi : \circ^{-1} \xrightarrow{q^{-1}} \circ^q \xrightarrow{q^{-3}} \circ^{q^3} \quad \rightsquigarrow_{s_p} \quad s_p^* \chi : \circ^{-1} \xrightarrow{q} \circ^{-1} \xrightarrow{q^{-3}} \circ^{q^3}.$$

Using (49), $\underline{E}_i (\underline{E}_p \underline{E}_i)^3 \underline{E}_j$ can be written as a linear combination of other words modulo $\mathcal{J}(s_p^* \chi)$, which are greater than this word for the order $p < i < j$, or begin with \underline{E}_p (recall that $\underline{E}_i^2 \in \mathcal{J}(s_p^* \chi)$). Multiplying on the left by \underline{E}_p , $(\underline{E}_p \underline{E}_i)^4 \underline{E}_j$ is expressed as a linear combination of greater words, modulo $\mathcal{J}(s_p^* \chi)$, using that $\underline{E}_p^2 \in \mathcal{J}(s_p^* \chi)$. By Lemma 3.4,

$$\begin{aligned} T_p((\text{ad}_c E_i)^4 E_j) &= \left[(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i, \left[(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i, \left[(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p) (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j \right]_{\underline{c}} \right]_{\underline{c}} \right]_{\underline{c}} \\ &= [(\underline{E}_p \underline{E}_i)^4 \underline{E}_j]_{\underline{c}} \in \mathcal{J}(s_p^* \chi). \end{aligned}$$

If $m_{ij} = 2$, then $m_{pi} = 2$ for the diagram

$$\chi : \circ^{\zeta^6} \xrightarrow{\zeta^4} \circ^{\zeta^5} \xrightarrow{\zeta^8} \circ^{\zeta} \quad \rightsquigarrow_{s_p} \quad s_p^* \chi : \circ^{\zeta^6} \xrightarrow{\zeta^8} \circ^{\zeta} \xrightarrow{\zeta^8} \circ^{\zeta},$$

where $\zeta \in \mathbb{G}_9$, or $m_{pi} = 1$. In the first case, using that $\underline{E}_p^3 \in \mathcal{J}(s_p^* \chi)$ and the quantum Serre relations we write $\underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_j \underline{E}_p \underline{E}_i$ as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, for the order $p < i < j$, so

$$[\underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_j \underline{E}_p \underline{E}_i]_{\underline{c}} = \left[[(\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p)^2 (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}}, (\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

If we call $c = s_p^* \chi(4\alpha_p + 2\alpha_i + \alpha_j, \alpha_p + \alpha_i) = \chi(2\alpha_i + \alpha_j, \alpha_p + \alpha_i)$, we have

$$\begin{aligned} T_p((\text{ad}_c E_i)^3 E_j) &= \left[(\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_i, [(\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p)^2 (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}} \right]_{\underline{c}} \\ &= (\text{ad}_{\underline{c}} \underline{E}_p) \left(\left[(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i, [(\text{ad}_{\underline{c}} \underline{E}_p)^2 \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p)^2 (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}} \right]_{\underline{c}} \right) \\ &= -c (\text{ad}_{\underline{c}} \underline{E}_p) ([\underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_j \underline{E}_p \underline{E}_i]_{\underline{c}}) \in \mathcal{J}(s_p^* \chi). \end{aligned}$$

If $m_{pi} = 1$ and $q_{pp} \neq -1$, using $(\text{ad}_{\underline{c} \underline{E}_p})^2 \underline{E}_i$ we write $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_i$ as a linear combination of greater words and $\underline{E}_p^2 \underline{E}_i^2 \underline{E}_j \underline{E}_i$, for the order induced by $p < i < j$, modulo $\mathcal{J}(s_p^* \chi)$. Using now $(\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_j$, and $(\text{ad}_{\underline{c} \underline{E}_i})^3 \underline{E}_j$ when $\widetilde{q}_{ip}^3 \neq 1$, or (51) in other case, $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_i$ is expressed as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$. If $m_{pi} = 1$ and $q_{pp} = -1$, then (48) is a generator of $\mathcal{J}(s_p^* \chi)$, so

$$\left[[(\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_i, (\text{ad}_{\underline{c} \underline{E}_p})(\text{ad}_{\underline{c} \underline{E}_i}) \underline{E}_j]_{\underline{c}}, \underline{E}_i \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi),$$

or (45) (considered for the pair p, i) is a generator of the ideal. In any case we have that

$$\begin{aligned} T_p((\text{ad}_{\underline{c} \underline{E}_i})^3 \underline{E}_j) &= \left[(\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_i, [(\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_i, (\text{ad}_{\underline{c} \underline{E}_p})(\text{ad}_{\underline{c} \underline{E}_i}) \underline{E}_j]_{\underline{c}} \right]_{\underline{c}} \\ &= (\text{ad}_{\underline{c} \underline{E}_p}) \left[\underline{E}_i, [(\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_i, (\text{ad}_{\underline{c} \underline{E}_p})(\text{ad}_{\underline{c} \underline{E}_i}) \underline{E}_j]_{\underline{c}} \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi). \end{aligned}$$

Now we fix $m_{ij} = 1$. If $m_{pi} = 1$, we analyze each different possible diagram.

- If $q_{pp} \neq -1$, then $s_p^* \chi$ is twist equivalent to χ (restricted to the vertices p, i, j), and $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j$ can be expressed as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, using the quantum Serre relations

$$(\text{ad}_{\underline{c} \underline{E}_p})^2 \underline{E}_i = (\text{ad}_{\underline{c} \underline{E}_i})^2 \underline{E}_j = (\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_j = 0.$$

- If $q_{pp} = -1$ and $q_{ii} q_{ip} q_{pi} = 1$, then $q_{ii} = -1$ and $q_{ip} q_{pi} q_{ij} q_{ji} = 1$. In this way (44) is a generator of the ideal, and by Lemma 3.7,

$$\begin{aligned} T_p((\text{ad}_{\underline{c} \underline{E}_i})^2 \underline{E}_j) &= [(\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_i, (\text{ad}_{\underline{c} \underline{E}_p})(\text{ad}_{\underline{c} \underline{E}_i}) \underline{E}_j]_{\underline{c}} \\ &= (\text{ad}_{\underline{c} \underline{E}_p}) \left([\underline{E}_i, (\text{ad}_{\underline{c} \underline{E}_p})(\text{ad}_{\underline{c} \underline{E}_i}) \underline{E}_j]_{\underline{c}} \right) \in \mathcal{J}(s_p^* \chi). \end{aligned}$$

- If $q_{pp} = -1$ and $q_{ii} q_{ip} q_{pi} \neq 1$, then $q_{pj} q_{jp} = q_{pj} q_{jp} = 1$, $q_{ij} q_{ji} = \widetilde{q}_{ij} \neq -1$ and

$$q_{ii}^{-1} = q_{pp}^{-1} q_{ip}^{-1} q_{pi}^{-1} q_{ii}^{-1} = -q_{ip} q_{pi} q_{ij} q_{ji} \neq -1,$$

so (53) or (54) are generators of $\mathcal{J}(s_p^* \chi)$, and then $T_p((\text{ad}_{\underline{c} \underline{E}_i})^2 \underline{E}_j) \in \mathcal{J}(s_p^* \chi)$.

Now we fix $m_{pi} \neq 1$. The possible connected diagrams of rank three with these conditions must verify $m_{pi} = 2$, $m_{ip} = 1$. Using the quantum Serre relations $(\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_j = (\text{ad}_{\underline{c} \underline{E}_p})^3 \underline{E}_i = 0$ if $q_{pp}^3 \neq -1$, or (46) in other case, $\underline{E}_p^2 \underline{E}_i \underline{E}_j \underline{E}_p \underline{E}_i$ can be expressed as a linear combination of greater words for the order $p < i < j$, and by Lemma 3.4,

$$[(\text{ad}_{\underline{c} \underline{E}_p})^2 (\text{ad}_{\underline{c} \underline{E}_i}) \underline{E}_j, (\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_i]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

By Lemma (3.7) we conclude that

$$\begin{aligned} T_p((\text{ad}_{\underline{c} \underline{E}_i})^2 \underline{E}_j) &= [(\text{ad}_{\underline{c} \underline{E}_p})^2 \underline{E}_i, (\text{ad}_{\underline{c} \underline{E}_p})^2 (\text{ad}_{\underline{c} \underline{E}_i}) \underline{E}_j]_{\underline{c}} \\ &= (\text{ad}_{\underline{c} \underline{E}_p}) \left([(\text{ad}_{\underline{c} \underline{E}_p}) \underline{E}_i, (\text{ad}_{\underline{c} \underline{E}_p})^2 (\text{ad}_{\underline{c} \underline{E}_i}) \underline{E}_j]_{\underline{c}} \right) \in \mathcal{J}(s_p^* \chi). \end{aligned}$$

FINALLY WE CONSIDER $m_{ij}, m_{pj} \neq 0$, so each pair of vertices is connected. If $m_{ij} = 2$, there is just one possibility,

$$\chi : \begin{array}{ccc} & \circ^{-1} & \\ q^{-1} \swarrow & & \searrow q^3 \\ \circ^q & \text{---} q^{-2} \text{---} & \circ^{-1} \end{array} \quad \longleftrightarrow_{s_p} \quad s_p^* \chi : \circ^{-1} \xrightarrow{q} \circ^{-1} \xrightarrow{q^{-3}} \circ^q$$

which is a diagram of type super $G(3)$. By (49) and Lemma 3.5 we have that

$$T_p((\text{ad}_{\underline{c}} E_i)^3 E_j) = \left[(\text{ad}_{\underline{c}} E_p) \underline{E}_i, \left[(\text{ad}_{\underline{c}} E_p) \underline{E}_i, \left[(\text{ad}_{\underline{c}} E_p) \underline{E}_i, (\text{ad}_{\underline{c}} E_p) \underline{E}_j \right]_{\underline{c}} \right]_{\underline{c}} \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

The remaining case is $m_{ij} = 1$. If $m_{pi} = m_{pj} = 1$, there are two possible cases:

- $q_{pp} = -1$; in this case (48) is a generator of the ideal $\mathcal{J}(s_p^* \chi)$ by definition, and by Lemma 3.5 we have that

$$T_p((\text{ad}_{\underline{c}} E_i)^2 E_j) = \left[(\text{ad}_{\underline{c}} E_p) \underline{E}_i, \left[(\text{ad}_{\underline{c}} E_p) \underline{E}_i, (\text{ad}_{\underline{c}} E_p) \underline{E}_j \right]_{\underline{c}} \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

- $q_{pp} \neq -1$, $q_{pp} q_{pi} q_{ip} = q_{pp} q_{pj} q_{jp} = 1$; $s_p^* \chi$ is twist equivalent to χ , so (47) is a generator of $\mathcal{J}(s_p^* \chi)$. Using also the quantum Serre relations $(\text{ad}_{\underline{c}} E_p)^2 \underline{E}_i$, $(\text{ad}_{\underline{c}} E_p)^2 \underline{E}_j$, $(\text{ad}_{\underline{c}} E_i)^2 \underline{E}_j$, $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_j$ is written as a linear combination of greater words, modulo $\mathcal{J}(s_p^* \chi)$, so as before $T_p((\text{ad}_{\underline{c}} E_i)^2 E_j) \in \mathcal{J}(s_p^* \chi)$.

It remains to consider the following braiding:

$$\begin{array}{ccc} & \circ^{\zeta} & \\ -\zeta^{-1} \swarrow & & \searrow \zeta^{-1} \\ \circ^{-\zeta} & \xrightarrow{-\zeta^{-1}} & \circ^{-1} \end{array}, \quad q_{pp} = \zeta \in \mathbb{G}_3, \quad m_{ij} = m_{pj} = 1, \quad m_{pi} = 2.$$

The diagram of $s_p^* \chi$ is $\circ^{-1} \xrightarrow{-1} \circ^{\zeta} \xrightarrow{\zeta^{-1}} \circ^{-1}$. Then (45) holds for p, i , and so $\underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p$ is expressed as a linear combination of other words of the same \mathbb{Z}^θ -degree. Multiplying on the left by \underline{E}_p^2 , on the right by \underline{E}_j , and using that $\underline{E}_p^3 = 0$, $\underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p \underline{E}_j$ can be written as a linear combination of greater words, so

$$T_p((\text{ad}_{\underline{c}} E_i)^2 E_j) = \left[(\text{ad}_{\underline{c}} E_p)^2 \underline{E}_i, \left[(\text{ad}_{\underline{c}} E_p)^2 \underline{E}_i, (\text{ad}_{\underline{c}} E_p) \underline{E}_j \right]_{\underline{c}} \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

Therefore we analyze all the cases and the proof is completed. \square

Lemma 3.12. *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{jj} = -1$, $\widetilde{q}_{ik} = \widetilde{q}_{ij} \widetilde{q}_{jk} = 1$. Then,*

$$T_p([E_{ijk}, E_j]_{\underline{c}}) \in \mathcal{J}(s_p^* \chi).$$

Proof. Note that in $U(\chi)$ we have the following identity, using the condition on the scalars, (9) and $(\text{ad}_{\underline{c}} E_i) E_k = E_j^2 = 0$:

$$[E_{ijk}, E_j]_{\underline{c}} = q_{ij} q_{kj} [E_j, E_{ijk}]_{\underline{c}} = q_{ij} q_{kj} [E_{ji}, E_{jk}]_{\underline{c}},$$

so it is enough to prove that one of these relations is applied in $\mathcal{J}(s_p^* \chi)$ by T_p for each possible diagram. Let $p = j$. Note that $q_{pp} = -1$, $q_{ip} q_{pi} q_{pk} q_{kp} = q_{ik} q_{ki} = 1$, so $(\text{ad}_{\underline{c}} E_i) \underline{E}_k$ and (44) are generators of $\mathcal{J}(s_p^* \chi)$. By (36) we have:

$$\begin{aligned} T_p([E_{ipk}, E_p]_{\underline{c}}) &= [a_1 \underline{E}_i, (\text{ad}_{\underline{c}} E_p) \underline{E}_k]_{\underline{c}} \underline{E}_p \underline{L}_p^{-1} + q_{ip} q_{kp} \underline{E}_p \underline{L}_p^{-1} [a_1 \underline{E}_i, (\text{ad}_{\underline{c}} E_p) \underline{E}_k]_{\underline{c}} \\ &= a_1 \left(q_{ip}^{-1} q_{pp}^{-1} q_{kp}^{-1} + q_{ip} q_{kp} \right) \underline{E}_p \underline{L}_p^{-1} [\underline{E}_i, (\text{ad}_{\underline{c}} E_p) \underline{E}_k]_{\underline{c}} \\ &\quad + a_1 q_{pk}^{-1} (1 - q_{pk} q_{kp}) (\underline{E}_i \underline{E}_k + q_{ip} q_{pk} q_{ik} \underline{E}_k \underline{E}_i) \\ &= a_1 q_{pk}^{-1} (1 - q_{pk} q_{kp}) (\text{ad}_{\underline{c}} E_i) \underline{E}_k \in \mathcal{J}(s_p^* \chi). \end{aligned}$$

Let $p = i$, which is analogous to the case $p = k$. By (64) and (66),

$$T_p([E_{jpk}, E_k]_{\underline{c}}) = [a_{m_{pj}} (\text{ad}_{\underline{c}} E_p)^{m_{pj}-1} \underline{E}_j, (\text{ad}_{\underline{c}} E_p)^{m_{pj}} (\text{ad}_{\underline{c}} E_j) \underline{E}_k]_{\underline{c}}.$$

Here $q \neq -1$. For the three diagrams, $m_{pi} = m_{pj} = 1$. Consider the order $p < j < i < k$:

$$T_p([E_{ji}, E_{jk}]_c) = \left[[E_{pj}, E_{pi}]_c, [E_{pj}, E_k]_c \right]_c = [E_p E_j E_p E_i E_p E_j E_k].$$

We can write $E_p E_j E_p E_i E_p E_j E_k$ as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$: in the first case, using (48); for the second one, we write $E_j E_p E_i E_p E_j$ as a linear combination of other words of the same degree by (53), where those words are greater than $E_j E_p E_i E_p E_j$ or begin with E_p , and then we use the quantum Serre relations; in the last case, $E_j E_p E_i$ is a linear combination of other words of the same degree by (47), where those words are greater than $E_j E_p E_i$ or begin with E_p . In all the cases we conclude that $T_p([E_{ji}, E_{jk}]_c) \in \mathcal{J}(s_p^* \chi)$. \square

Lemma 3.13. *Let $i, j \in \{1, \dots, \theta\}$ be such that $q_{jj} = -1$, and also $q_{ii} = \pm \widetilde{q}_{ij} \in \mathbb{G}_3$, or $q_{ii} \widetilde{q}_{ij} \in \mathbb{G}_6$. Then, $T_p([E_{ii}, E_{ij}]_c) \in \mathcal{J}(s_p^* \chi)$, for any $p \in \{1, \dots, \theta\}$.*

Proof. We denote $\mathbf{x} := [E_{ii}, E_{ij}]_c$. We begin with the case $p = j$. Note that $m_{pi} = 1$ (because $q_{pp} = -1$, $\widetilde{q}_{pi} \neq 1$), $3\alpha_i + 2\alpha_p \notin \Delta_+^\chi$, so

$$s_p(3\alpha_i + 2\alpha_p) = 3\alpha_i + \alpha_p \notin \Delta_+^{s_p^* \chi}.$$

Using (66) and $(\text{ad}_c E_i)^3 E_p \in \mathcal{J}(s_p^* \chi)$, we obtain that

$$T_p(\mathbf{x}) = a_2 a_1 \left[[E_{pi}, E_i]_c, E_i \right]_c \in \mathcal{J}(s_p^* \chi).$$

Now let $p = i$. By Lemma 3.4 it is equivalent to prove that

$$T_p(\mathbf{x}') \in \mathcal{J}(s_p^* \chi), \quad \text{where } \mathbf{x}' := \left[E_{jp}, [E_{jp}, E_p]_c \right]_c,$$

because we have proved that T_p apply the generating relations of degree less than \mathbf{x} in elements of $\mathcal{J}(s_p^* \chi)$. By (36),

$$T_p(\mathbf{x}') = a_1^2 a_2 \left[(\text{ad}_c E_p) E_j, E_j \right]_c \in \mathcal{J}(s_p^* \chi),$$

because it holds that $q_{jj} = -1$, or $q_{jj} q_{jp} q_{pj} = 1$.

Finally, let $p \neq i, j$; the case $m_{pi} = m_{pj} = 0$ follows easily as in the previous Lemmata, so consider the case in which p, i, j determine a connected subdiagram of rank three. We note that $q_{ii} \in \mathbb{G}_3$.

We take first $m_{pi} \neq 0$, $m_{pj} = 0$. The possible braidings verify that $m_{pi} = 1$, so for the order $p < i < j$,

$$T_p(\mathbf{x}) = \left[[E_{pi}, [E_{pi}, E_j]_c]_c, [E_{pi}, E_j]_c \right]_c = [E_p E_i (E_p E_i E_j)^2]_c,$$

where we use that $(\text{ad}_c E_p) E_j = 0$ in $U(s_p^* \chi)$. As $q_{pi} q_{ip} q_{ii} = 1$ or $q_{pi} q_{ip} = \pm q_{ii}$, we have that $q_{ii} = -1$, or $q_{ii} q_{ip} q_{pi} = 1$, or $q_{ii}^2 q_{ip} q_{pi} = 1$, or $q_{ii} = -q_{ip} q_{pi} \in \mathbb{G}_3$, so $E_p E_i E_p E_i E_j E_p E_i E_j$ can be expressed as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, using the quantum Serre relations or (45). We deduce that $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$, using the Lemma 3.4.

Now let $m_{pi} = 0$, $m_{pj} \neq 0$. We note that $m_{pj} = 1$ for any possible diagram, and in $U(s_p^*\chi)$ we have that:

$$\begin{aligned} T_p(\mathbf{x}) &= [(\text{ad}_{\underline{c}} \underline{E}_i)^2 (\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_j, (\text{ad}_{\underline{c}} \underline{E}_i) (\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_j]_{\underline{c}} \\ &= \underline{q}_{ip}^3 [(\text{ad}_{\underline{c}} \underline{E}_p) (\text{ad}_{\underline{c}} \underline{E}_i)^2 \underline{E}_j, (\text{ad}_{\underline{c}} \underline{E}_p) (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}} \\ &= \underline{q}_{ip}^3 (\text{ad}_{\underline{c}} \underline{E}_p) [(\text{ad}_{\underline{c}} \underline{E}_i)^2 \underline{E}_j, (\text{ad}_{\underline{c}} \underline{E}_p) (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}} \\ &= \underline{q}_{ip}^3 (\text{ad}_{\underline{c}} \underline{E}_p) [(\text{ad}_{\underline{c}} \underline{E}_i)^2 (\text{ad}_{\underline{c}} \underline{E}_j) \underline{E}_p, (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}} = 0, \end{aligned}$$

where we apply (64) (because $(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i = 0$), (9), and finally that (46) is a generator of $\mathcal{J}(s_p^*\chi)$, because $\underline{q}_{ii} = q_{ii} \in \mathbb{G}_3$, $\underline{m}_{ji} = \underline{m}_{jp} = 1$, and (45) is also a generator by Lemma 3.8.

Finally, consider $m_{pi}, m_{pj} \neq 0$. There exists just one possible braiding: $q_{pp} = -1 = q_{pj}q_{jp}$, $q_{ii} = -\widetilde{q}_{ij} = q_{pi}^{-1}q_{ip}^{-1}$. The diagram of $s_p^*\chi$ is $\circ^{-1} \xrightarrow{-1} \circ^{-1} \xrightarrow{q_{ii}} \circ^{-1}$, and the solution is analogous to the previous case, but now we use (48). \square

Lemma 3.14. *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = \pm \widetilde{q}_{ij} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, and $q_{jj}\widetilde{q}_{ij} = q_{jj}\widetilde{q}_{jk} = 1$ or $q_{jj} = -1$, $\widetilde{q}_{ij}\widetilde{q}_{jk} = 1$. Then, for any $p \in \{1, \dots, \theta\}$,*

$$T_p([E_{iijk}, E_{ij}]_{\underline{c}}) \in \mathcal{J}(s_p^*\chi).$$

Proof. We denote $\mathbf{x} = [E_{iijk}, E_{ij}]_{\underline{c}}$. We begin with the case $p = k$. In all the cases we have that $m_{kj} = 1$, and $s_p^*\chi$ satisfies the same conditions, so (46) is a generator of $\mathcal{J}(s_p^*\chi)$. Then,

$$\begin{aligned} T_p(\mathbf{x}) &\cong [(\text{ad}_{\underline{c}} \underline{E}_i)^2 (a_1 \underline{E}_j), (\text{ad}_{\underline{c}} \underline{E}_i) (\text{ad}_{\underline{c}} \underline{E}_k) \underline{E}_j]_{\underline{c}} \cong a_1 q_{ik} [E_{iij}, E_{kij}]_{\underline{c}} \\ &\cong a_1 q_{ik} [E_{iijk}, E_{ij}]_{\underline{c}} \cong 0 \pmod{\mathcal{J}(s_p^*\chi)}, \end{aligned}$$

where we apply first (66), then (64), $(\text{ad}_{\underline{c}} \underline{E}_k) \underline{E}_i = 0$ for the second line, and finally (9) plus the fact that (45) is a generator of $\mathcal{J}(s_p^*\chi)$. The cases $p = i$, $p = j$ are proved in a similar way to the case $p = i$ of previous Lemma.

Finally take $p \neq i, j, k$, and assume that p is connected with at least one of the other vertices; in other case the proof is easy as above. We have two possible cases: $m_{pi} = 1$, $m_{pj} = m_{pk} = 0$, or $m_{pk} = 1$, $m_{pj} = m_{pi} = 0$. For the first one,

$$T_p(\mathbf{x}) = \left[[E_{pi}, [E_{pi}, E_{jk}]_{\underline{c}}]_{\underline{c}}, [E_{pi}, E_j]_{\underline{c}} \right]_{\underline{c}},$$

and we have two possibilities:

- if $q_{pp} = -1$, then $q_{ii}\widetilde{q}_{ip} = 1$, and $q_{ii} = -1$, so (48) is a generator of $\mathcal{J}(s_p^*\chi)$ for the subdiagram determined by p, i, j . Therefore $T_p(\mathbf{x}) \in \mathcal{J}(s_p^*\chi)$.
- if $q_{pp} \neq -1$, then $q_{pp} = q_{ip}^{-1}q_{pi}^{-1} = q_{ii}$, so $s_p^*\chi$ is twist equivalent to χ and (46) is a generator of $\mathcal{J}(s_p^*\chi)$. Then $T_p(\mathbf{x}) \in \mathcal{J}(s_p^*\chi)$, because it is obtained after to apply $(\text{ad}_{\underline{c}} \underline{E}_p)^3$ to (46) and multiply by a non-zero scalar, where we use also the quantum Serre relations involving \underline{E}_p .

For the second case, we use $(\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_i, (\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_j \in \mathcal{J}(s_p^*\chi)$ to obtain that

$$\begin{aligned} T_p(\mathbf{x}) &\cong [(\text{ad}_{\underline{c}} \underline{E}_i)^2 (\text{ad}_{\underline{c}} \underline{E}_j) (\text{ad}_{\underline{c}} \underline{E}_p) \underline{E}_k, (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}} \\ &\cong \underline{q}_{ip}^2 \underline{q}_{jp} [(\text{ad}_{\underline{c}} \underline{E}_p) (\text{ad}_{\underline{c}} \underline{E}_i)^2 (\text{ad}_{\underline{c}} \underline{E}_j) \underline{E}_k, (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}} \\ &\cong \underline{q}_{ip}^2 \underline{q}_{jp} (\text{ad}_{\underline{c}} \underline{E}_p) \left([(\text{ad}_{\underline{c}} \underline{E}_i)^2 (\text{ad}_{\underline{c}} \underline{E}_j) \underline{E}_k, (\text{ad}_{\underline{c}} \underline{E}_i) \underline{E}_j]_{\underline{c}} \right) \cong 0 \pmod{\mathcal{J}(s_p^*\chi)}, \end{aligned}$$

by (64) and the fact that (46) is a generator of $\mathcal{J}(s_p^*\chi)$. \square

Lemma 3.15. *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $\widetilde{q}_{ik}, \widetilde{q}_{ij}, \widetilde{q}_{jk} \neq 1$. Then, for any p ,*

$$T_p \left(E_{ijk} - \frac{1 - \widetilde{q}_{jk}}{q_{kj}(1 - \widetilde{q}_{ik})} [E_{ik}, E_j]_c - q_{ij}(1 - \widetilde{q}_{jk}) E_j E_{ik} \right) \in \mathcal{J}(s_p^*\chi).$$

Proof. Let $\mathbf{x} = E_{ijk} - \frac{1 - \widetilde{q}_{jk}}{q_{kj}(1 - \widetilde{q}_{ik})} [E_{ik}, E_j]_c - q_{ij}(1 - \widetilde{q}_{jk}) E_j E_{ik}$. By a direct computation we obtain the same relation, up to an scalar, if we permute the vertices i, j, k , where we use that $\widetilde{q}_{ik}\widetilde{q}_{ij}\widetilde{q}_{jk} = 1$, so it is enough to consider one of these permutations for each p .

Consider then $p = k$, which is analogous to take $p = i$ or $p = j$. Note that $\{m_{pi}, m_{pj}\} = \{1, 1\}$, or $\{m_{pi}, m_{pj}\} = \{1, 2\}$, so we fix $m_{pj} = 1$, $m_{pi} \in \{1, 2\}$. By (66),

$$\begin{aligned} T_p(E_{ijk}) &= (q_{pp}^{-1} q_{pj}^{-1} q_{jp}^{-1} - 1) q_{jp} q_{pp} [(\text{ad}_{\underline{E}_p})^{m_{pi}} \underline{E}_i, \underline{E}_j]_{\underline{c}}, \\ T_p([E_{ik}, E_j]_c) &= (q_{pp}^{-1 - m_{pi}} q_{pi}^{-1} q_{ip}^{-1} - 1) q_{ip} q_{pp} [(\text{ad}_{\underline{E}_p})^{m_{pi} - 1} \underline{E}_i, (\text{ad}_{\underline{E}_p}) \underline{E}_j]_{\underline{c}}, \\ T_p(E_j E_{ik}) &= (q_{pp}^{-1 - m_{pi}} q_{pi}^{-1} q_{ip}^{-1} - 1) q_{ip} q_{pp} (\text{ad}_{\underline{E}_p}) \underline{E}_j (\text{ad}_{\underline{E}_p})^{m_{pi} - 1} \underline{E}_i. \end{aligned}$$

If $m_{pi} = 2$, or $m_{pi} = 1$, $q_{pp} \neq -1$, then $q_{ik} q_{ki}, q_{ij} q_{ji}, q_{jk} q_{kj} \neq 1$ and we deduce that $T_p(\mathbf{x}) \in \mathcal{J}(s_p^*\chi)$ from the fact that (47) is a generator of $\mathcal{J}(s_p^*\chi)$, because we can write then $\underline{E}_p^{m_{pi}} \underline{E}_i \underline{E}_j$ as a linear combination of greater words (for the order on the letters $p < i < j$), modulo $\mathcal{J}(s_p^*\chi)$, y apply then Lemma 3.4. If $q_{pp} = -1$ then $q_{ij} q_{ji} = 1$, so $(\text{ad}_{\underline{E}_i}) \underline{E}_j \in \mathcal{J}(s_p^*\chi)$. By a direct computation, there exists $a \in \mathbf{k}^\times$ such that

$$T_p(\mathbf{x}) = a (\text{ad}_{\underline{E}_p}) (\text{ad}_{\underline{E}_i}) \underline{E}_j \in \mathcal{J}(s_p^*\chi).$$

Let $p \neq i, j, k$. We note that p is not connected with any of the other vertices (so the proof follows easily as in the previous Lemmata), or p is connected just with one of these vertices. For the last case we can assume that $m_{pi} \neq 0$, so the unique possibility is $m_{pi} = m_{ip} = 1$. Then $\widetilde{q}_{ik} = q_{ik} q_{ki}$, $\widetilde{q}_{ij} = q_{ij} q_{ji}$, $\widetilde{q}_{jk} = q_{kj} q_{jk} \neq 1$, so (47) is a generator of $\mathcal{J}(s_p^*\chi)$. By Lemma 3.7 and the relations $(\text{ad}_{\underline{E}_p}) \underline{E}_j = (\text{ad}_{\underline{E}_p}) \underline{E}_k = 0$, we deduce that $T_p(\mathbf{x})$ is obtained, up to a non-zero scalar, after to apply $(\text{ad}_{\underline{E}_p})$ to (47), modulo $\mathcal{J}(s_p^*\chi)$, so $T_p(\mathbf{x}) \in \mathcal{J}(s_p^*\chi)$. \square

Lemma 3.16. *Let $i, j, k \in \{1, \dots, \theta\}$ be such that*

- (i) $q_{ii} = q_{jj} = -1$, $\widetilde{q}_{ij}^2 = \widetilde{q}_{jk}^{-1}$, $\widetilde{q}_{ik} = 1$, or
- (ii) $\widetilde{q}_{ij} = q_{jj} = -1$, $q_{ii} = -\widetilde{q}_{jk}^2 \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, or
- (iii) $q_{kk} = \widetilde{q}_{jk} = q_{jj} = -1$, $q_{ii} = -\widetilde{q}_{ijk} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, or
- (iv) $q_{jj} = -1$, $\widetilde{q}_{ij} = q_{ii}^{-2}$, $\widetilde{q}_{jk} = -q_{ii}^3$, $\widetilde{q}_{ik} = 1$, or
- (v) $q_{ii} = q_{jj} = q_{kk} = -1$, $\pm \widetilde{q}_{ij} = \widetilde{q}_{jk} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$,

Then, for any p , $T_p([E_{ij}, E_{ijk}]_c, E_j]_c) \in \mathcal{J}(s_p^\chi)$.*

Proof. Denote $\mathbf{x} = [[E_{ij}, E_{ijk}]_c, E_j]_c$; we analyze each case.

(i) We begin with the case $p = k$; by (66) and as $\underline{E}_i^2, \underline{E}_j^2, (\text{ad}_{\underline{E}_i}) \underline{E}_p$ are generators of $\mathcal{J}(s_p^*\chi)$ (note that $s_p^*\chi$ is twist equivalent to χ), we have that

$$\begin{aligned} T_p(\mathbf{x}) &\cong a_1 \left[[\underline{E}_{ipj}, \underline{E}_{ij}]_{\underline{c}}, \underline{E}_{pj} \right]_{\underline{c}} \cong a_1 q_{ip} \left[[\underline{E}_{pij}, \underline{E}_{ij}]_{\underline{c}}, \underline{E}_{pj} \right]_{\underline{c}} \\ &\cong a \left[[\underline{E}_{pji}, \underline{E}_{ji}]_{\underline{c}}, \underline{E}_{pj} \right]_{\underline{c}} \cong a \left[\underline{E}_p \underline{E}_j \underline{E}_i \underline{E}_j \underline{E}_i \underline{E}_p \underline{E}_j \right]_{\underline{c}} \pmod{\mathcal{J}(s_p^*\chi)}, \end{aligned}$$

for some $a \in \mathbf{k}^\times$, where we use the order on the letters $p < j < i$. As (48) is also a generator of $\mathcal{J}(s_p^*\chi)$, we can write $\underline{E}_j \underline{E}_i \underline{E}_j \underline{E}_i \underline{E}_p \underline{E}_j$ as a linear combination of other

words, greater than this word or beginning with \underline{E}_p . Multiplying on the left by \underline{E}_p and using the quantum Serre relations $\underline{E}_p \underline{E}_j \underline{E}_i \underline{E}_j \underline{E}_i \underline{E}_p \underline{E}_j$ is expressed as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, so by Lemma 3.4, $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$.

Let $p = j$; note that $m_{pi} = m_{pk} = 1$. Also, $q_{ii}^{-1} = q_{ik} q_{ki}$, so $(\text{ad}_{\underline{c}} \underline{E}_i)^2 \underline{E}_k \in \mathcal{J}(s_p^* \chi)$; use (66) and work as in the case $p = i$ of Lemma 3.13 to obtain that

$$T_p(\mathbf{x}) = a_1^2 [\underline{E}_i, \underline{E}_{ipk}]_{\underline{c}} \underline{E}_p \underline{L}_p^{-1} - a_1^2 q_{ip}^2 q_{kp} \underline{F}_p \underline{L}_p^{-1} [\underline{E}_i, \underline{E}_{ipk}]_{\underline{c}} = b (\text{ad}_{\underline{c}} \underline{E}_i)^2 \underline{E}_k \in \mathcal{J}(s_p^* \chi),$$

for some $b \in \mathbf{k}^\times$.

Let now $p = i$. As in the previous Lemmata, it is enough to prove the statement for

$$\mathbf{x}' := [[E_{kjp}, E_{jp}]_{\underline{c}}, E_j]_{\underline{c}}.$$

We apply (66) to obtain, for the order on the letters $k < i < j$,

$$T_p(\mathbf{x}') = \left[[[E_k, a_1 E_j]_{\underline{c}}, E_j]_{\underline{c}}, (\text{ad}_{\underline{c}} \underline{E}_p) E_j \right]_{\underline{c}} = a_1^2 [E_k E_j^2 E_p E_j]_{\underline{c}}.$$

As $q_{jj} q_{ji} q_{ij} = q_{jj}^2 q_{jk} q_{kj} = 1$, we deduce by (51) if $q_{jj} \in \mathbb{G}_3$, or by $(\text{ad}_{\underline{c}} \underline{E}_j)^2 \underline{E}_p = (\text{ad}_{\underline{c}} \underline{E}_j)^3 \underline{E}_k = 0$, if $q_{jj} \notin \mathbb{G}_3$, that $E_k E_j^2 E_p E_j$ is a linear combination of greater words, so $T_p(\mathbf{x}') \in \mathcal{J}(s_p^* \chi)$, and then $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$.

If $p \neq i, j, k$, then p is not connected with these three vertices, or p is connected just with i , satisfying also $q_{pp} \widetilde{q}_{pi} = 1$, or $q_{pp} = -1$, $\widetilde{q}_{ip} \widetilde{q}_{ij} = 1$. For the last case we have:

$$\begin{aligned} T_p(\mathbf{x}) &\cong \left[[E_{pij}, E_{pijk}]_{\underline{c}}, E_j \right]_{\underline{c}} \cong (\text{ad}_{\underline{c}} \underline{E}_p) \left([E_{pij}, E_{ijk}]_{\underline{c}}, E_j \right)_{\underline{c}} \\ &\cong (\text{ad}_{\underline{c}} \underline{E}_p) \left([E_p E_i E_j E_i E_j E_k E_j]_{\underline{c}} \right) \pmod{\mathcal{J}(s_p^* \chi)} \end{aligned}$$

by using first Lemma 3.7, then $(\text{ad}_{\underline{c}} \underline{E}_p) E_j$, $(\text{ad}_{\underline{c}} \underline{E}_p) E_j$, $(\text{ad}_{\underline{c}} \underline{E}_p) E_j \in \mathcal{J}(s_p^* \chi)$, and fixing the order $p < i < j < k$. We conclude that $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$ by using (50) if $q_{pp} = -1$, or using the quantum Serre relations corresponding to $\text{ad}_{\underline{c}} \underline{E}_p$ to write $\underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_i \underline{E}_j \underline{E}_k \underline{E}_j$ as a linear combination of greater words and apply Lemma 3.4 to deduce that

$$[E_p E_i E_j E_i E_j E_k E_j]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

(ii), (iii), (iv), (v) If $p \in \{i, j, k\}$ the proof is completely analogous to the previous case.

Let $p \neq i, j, k$, so p is not connected with any of these vertices, or it connected only with i , or only with k . The first case is easy. For the second case, $m_{pi} = 1$, because $q_{pp} = 1$ or $q_{pp}^{-1} = \widetilde{q}_{ip} \neq -1$, and the solution follows as in the previous case. For the last case, $m_{pk} = 1$ and the proof is also analogous, considering the previous \mathbf{x}' . \square

Lemma 3.17. (i) Let $i, j, k, l \in \{1, \dots, \theta\}$ be such that $q_{kk} = -1$, $\widetilde{q}_{jk}^2 = \widetilde{q}_{lk}^{-1} = ql$, $q_{jj} \widetilde{q}_{ij} = q_{jj} \widetilde{q}_{jk} = \widetilde{q}_{ik} = \widetilde{q}_{il} = \widetilde{q}_{jl} = 1$. Then, for all p , $T_p \left([[E_{ijkl}, E_k]_{\underline{c}}, E_j]_{\underline{c}}, E_k \right)_{\underline{c}} \in \mathcal{J}(s_p^* \chi)$.

(ii) Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = q_{jj} = -1$, $\widetilde{q}_{ij}^3 = \widetilde{q}_{jk}^{-1} = q_{kk} \neq \pm 1$, $\widetilde{q}_{ik} = 1$. Then, for all p , $T_p \left([[E_{ij}, [E_{ij}, E_{ijk}]_{\underline{c}}]_{\underline{c}}, E_j]_{\underline{c}} \right) \in \mathcal{J}(s_p^* \chi)$.

Proof. (i) The proof is analogous to (i) of the previous Lemma, because if $p \neq i, j, k, l$ is connected with some of them, then p is connected only with i with the same conditions.

(ii) If $p \in \{i, j, k\}$ the proof is completely analogous to the previous Lemma. If $p \neq i, j, k$ is connected with some of them, then p is connected only with i and $q_{pp} = -1$, $\widetilde{q}_{pi} = -\widetilde{q}_{ij} \in \mathbb{G}_4$. Anyway, the proof is analogous to the previous Lemma. \square

Lemma 3.18. (i) Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{jj} = \widetilde{q}_{ij}^2 = \widetilde{q}_{jk} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$. Then, for all p , $T_p \left([[E_{ijk}, E_j]_c, E_j]_c \right) \in \mathcal{J}(s_p^* \chi)$.
(ii) Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{jj} = \widetilde{q}_{ij}^3 = \widetilde{q}_{jk} \in \mathbb{G}_4$, $\widetilde{q}_{ik} = 1$. Then, for all p , $T_p \left([[[[E_{ijk}, E_j]_c, E_j]_c, E_j]_c \right) \in \mathcal{J}(s_p^* \chi)$.

Proof. (i) Let $\mathbf{x} = [[E_{ijk}, E_j]_c, E_j]_c$. For the case $p = k$, note that $m_{pj} = 1$ in all the cases, so for the order $i < p < j$ on the letters we have by (66):

$$T_p(\mathbf{x}) = a_1 \left[[[E_{ij}, E_{pj}]_c, E_{pj}]_c = [E_i E_j E_p E_j E_p E_j]_c \right].$$

As $(\text{ad}_c E_p)^2 E_j \in \mathcal{J}(s_p^* \chi)$, or (48) is a generator of $\mathcal{J}(s_p^* \chi)$, and also $E_j^3, (\text{ad}_c E_k) E_i \in \mathcal{J}(s_p^* \chi)$, $E_i E_j E_k E_j E_k E_j$ can be written as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, so $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$.

Consider now $p = j$. By (66) and the relations defining $\mathcal{J}(s_p^* \chi)$,

$$T_p(\mathbf{x}) = \left[[[E_i, (\text{ad}_c E_p)^2 E_k]_c, E_p E_p^{-1}]_c, E_p E_p^{-1}]_c = a(\text{ad}_c E_i) E_k,$$

for some $a \in \mathbf{k}^\times$, so $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$.

Let $p = i$. It is equivalent to prove that $T_p(\mathbf{x}') \in \mathcal{J}(s_p^* \chi)$, where $\mathbf{x}' = [[E_{kjp}, E_j]_c, E_j]_c$. We note that $m_{ij} = 1$ for all the possible diagrams, so $T_p(\mathbf{x}) = a_1 \left[[E_{kj}, E_{pj}]_c, E_{pj}]_c \right]$, and a proof similar to the case $p = k$ tells us that $T_p(\mathbf{x}') \in \mathcal{J}(s_p^* \chi)$, so $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$.

Finally, if $p \neq i, j, k$, then p is not connected to any of these vertices, or p is connected only with i , or it is connected only with k . The proof of the first case is again trivial, and for the other two cases $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$, using Lemma 3.7 and the fact that $s_p^* \chi$ is twist equivalent to χ .

(ii) The proof is analogous to (i). □

Lemma 3.19. (i) Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = \widetilde{q}_{ij} = -1$, $q_{jj} = \widetilde{q}_{jk}^{-1} \neq -1$, $\widetilde{q}_{ik} = 1$. Then, for all p , $T_p \left([E_{ij}, E_{ijk}]_c \right) \in \mathcal{J}(s_p^* \chi)$.

(ii) Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{jj} = q_{kk} = \widetilde{q}_{jk} = -1$, $q_{ii} = -\widetilde{q}_{ij} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$. Then, for all p , $T_p \left([E_{iij}, E_{ijk}]_c \right) \in \mathcal{J}(s_p^* \chi)$.

Proof. (i) Let $\mathbf{x} = [E_{ij}, E_{ijk}]_c$. If $p = k$, $m_{pj} = 1$ in all the possible diagrams and then:

$$T_p(\mathbf{x}) = a_1 [E_{ikj}, E_{ij}]_c = a_1 q_{ip} [E_{kij}, E_{ij}]_c.$$

We consider the two possible values of q_{pp} . When $q_{pp} \neq -1$, the diagram is of Cartan C_3 type, associated to a root of order 4, and $s_p^* \chi$ is twist equivalent to χ . Therefore $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$, using that (53) is again a generator of $\mathcal{J}(s_p^* \chi)$. If $q_{pp} = -1$, then $q_{jj} \in \mathbb{G}_3 \cup \mathbb{G}_4 \cup \mathbb{G}_6$, and $q_{jj} = \widetilde{q}_{ij} = q_{ii} = -1$, so (43) is a generator of $\mathcal{J}(s_p^* \chi)$ and

$$T_p(\mathbf{x}) = a_1 q_{ip} [E_{kij}, E_{ij}]_c = a_1 q_{ip} \text{ad}_c E_k (E_{ij}^2) \in \mathcal{J}(s_p^* \chi).$$

If $p = j$, we consider the different possible orders of q_{pp} . If $q_{pp} \in \mathbb{G}_4 \cup \mathbb{G}_6$, then p is a Cartan vertex and $s_p^* \chi$ is twist equivalent to $s_p^* \chi$, and $m_{pi} = 2, 3$

$$T_p(\mathbf{x}) = a_{m_{pi}} [(\text{ad}_c E_p)^{m_{pi}-1} E_i, [(\text{ad}_c E_p)^{m_{pi}-1} E_i, E_{pk}]_c]_c = [E_p^{m_{pi}-1} E_i E_p^{m_{pi}-1} E_i E_p E_j]_c.$$

We use the quantum Serre relations and (53) to write $\underline{E}_p^{m_{pi}-1} \underline{E}_i \underline{E}_p^{m_{pi}-1} \underline{E}_i \underline{E}_p \underline{E}_j$ as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$. If $q_{pp} \in \mathbb{G}_3$, then

$$\circ^{q_{ii}=-1} \xrightarrow{-1} \circ^{q_{pp}} \xrightarrow{q_{pp}^2} \circ^{q_{kk}=-1} \quad \longleftrightarrow_{s_p} \quad \begin{array}{ccc} & \circ^{q_{pp}} & \\ -q_{pp}^2 \swarrow & & \searrow q_{pp}^2 \\ \circ^{-q_{pp}} & \xrightarrow{-q_{pp}^2} & \circ^{q_{kk}=-1} \end{array}$$

The result follows in a similar way, but using that (45) and (47) are generators of $\mathcal{J}(s_p^* \chi)$ in this case.

If $p = i$, by Lemma 3.5, it is equivalent to prove that

$$T_p(\mathbf{x}') \in \mathcal{J}(s_p^* \chi), \quad \mathbf{x}' := [E_{kjp}, E_{jp}]_c.$$

Note that $(\text{ad}_c \underline{E}_j)^2 \underline{E}_k \in \mathcal{J}(s_p^* \chi)$, because for all the possible diagrams $q_{jj}^{-1} = q_{jk} q_{kj} \neq -1$. By (66), we have

$$T_p(\mathbf{x}') = a_1^2 [(\text{ad}_c \underline{E}_k) \underline{E}_j, \underline{E}_j]_c \in \mathcal{J}(s_p^* \chi).$$

Finally, if $p \neq i, j, k$, then p is not connected with any of these vertices, or p is connected just with i and $m_{pi} = 1$, or p is connected just with k and $m_{pk} = 1$. The proof is analogous to the corresponding case in previous Lemmata.

(ii) If $p \neq i, j, k$, then p is not connected with them and the result follows easily. In consequence, we consider the case $p \in \{i, j, k\}$. If $p = k$, $s_p^* \chi$ is twist equivalent to χ and (45) is a generator of $\mathcal{J}(s_p^* \chi)$. Therefore (66) implies that

$$T_p([E_{iijk}, E_{ijk}]_c) = a_2^2 [E_{iij}, E_{ij}]_c \in \mathcal{J}(s_p^* \chi).$$

If $p = j$, we have that

$$\circ^{q_{ii}} \xrightarrow{-q_{ii}} \circ^{-1} \xrightarrow{-1} \circ^{-1} \quad \longleftrightarrow_{s_p} \quad \begin{array}{ccc} & \circ^{-1} & \\ -q_{ii}^2 \swarrow & & \searrow -1 \\ \circ^{q_{ii}^2} & \xrightarrow{q_{ii}} & \circ^{-1} \end{array}$$

By (66) and fixing the order $p < i < k$, we have that

$$T_p([E_{iijk}, E_{ijk}]_c) = a_2^2 [[E_{pi}, E_{pik}]_c, E_{pik}]_c = [E_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_k \underline{E}_p \underline{E}_i \underline{E}_k]_c$$

We write $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_k \underline{E}_p \underline{E}_i \underline{E}_k$ as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$ using that (47), (45), \underline{E}_p^2 , \underline{E}_k^2 and \underline{E}_{iik} are generators of $\mathcal{J}(s_p^* \chi)$, so Lemma 3.4 implies that $T_p([E_{iijk}, E_{ijk}]_c) \in \mathcal{J}(s_p^* \chi)$.

Finally let $p = i$. By Lemma 3.5, it is equivalent to prove that

$$T_p(\mathbf{x}') \in \mathcal{J}(s_p^* \chi), \quad \mathbf{x}' := [E_{kjp}, E_{kjpp}]_c.$$

Note that $s_p^* \chi$ is twist equivalent to χ , so x_{jp}^2 is a generator of $\mathcal{J}(s_p^* \chi)$. Applying (66),

$$T_p(\mathbf{x}') = a_2^2 a_1 [E_{kjp}, E_{kj}]_c = a_2^2 a_1 [E_{kj}^2, E_p]_c \in \mathcal{J}(s_p^* \chi).$$

Therefore $T_p([E_{iijk}, E_{ijk}]_c) \in \mathcal{J}(s_p^* \chi)$. □

Lemma 3.20. (i) Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = \widetilde{q}_{ij} = -\widetilde{q}_{ik} \in \mathbb{G}_3$, $\widetilde{q}_{jk} = 1$, $q_{jj} = -1$, $q_{kk} \in \{-1, \widetilde{q}_{ik}^{-1}\}$. Then, for all p ,

$$T_p \left([E_i, [E_{ij}, E_{ik}]_c]_c + q_{jk} q_{ik} q_{ji} [E_{ik}, E_{ij}]_c + q_{ij} E_{ij} E_{ik} \right) \in \mathcal{J}(s_p^* \chi).$$

(ii) Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = q_{kk} = -1$, $\widetilde{q}_{ik} = 1$, $\widetilde{q}_{ij} \in \mathbb{G}_3$, $q_{jj} = -\widetilde{q}_{jk} = \pm \widetilde{q}_{ij}$. Then, for all p ,

$$T_p \left([E_i, E_{jjk}]_c - q_{kj}^{-1} (1 + q_{jj}^2) [E_{ijk}, E_j]_c + (1 + q_{jj})(1 + q_{jj}^2) E_j E_{ijk} \right) \in \mathcal{J}(s_p^* \chi).$$

Proof. If $p \neq i, j, k$, then p is not connected with these three vertices and the proof follows easily for both relations. We consider then $p \in \{i, j, k\}$ for each item.

(i) If $p = k$, $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_i$ is a linear combination of greater words by (45) and the quantum Serre relations, depending on the value of q_{kk} . In this way, there exist $a, b \in \mathbf{k}$ such that

$$\mathbf{x}' := [\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_i]_c + a [\underline{E}_p \underline{E}_i^2 \underline{E}_p \underline{E}_i \underline{E}_j]_c + b [\underline{E}_p \underline{E}_i \underline{E}_j]_c [\underline{E}_p \underline{E}_i^2]_c \in \mathcal{J}(s_p^* \chi).$$

On the other hand, by (66) and for the order on the letters $p < i < k$,

$$\begin{aligned} T_p \left([x_i, [(\text{ad}_c x_i) x_j, (\text{ad}_c x_i) x_k]_c]_c \right) &= a_1 \left[(\text{ad}_c \underline{E}_p) \underline{E}_i, [(\text{ad}_c \underline{E}_p) (\text{ad}_c \underline{E}_i) \underline{E}_j, \underline{E}_i]_c \right]_c \\ &= a_1 [\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_i]_c, \\ T_p \left([(\text{ad}_c x_i)^2 x_k, (\text{ad}_c x_i) x_j]_c \right) &= a_1 \left[[(\text{ad}_c \underline{E}_p) \underline{E}_i, \underline{E}_i]_c, (\text{ad}_c \underline{E}_p) (\text{ad}_c \underline{E}_i) \underline{E}_k \right]_c \\ &= a_1 [\underline{E}_p \underline{E}_i^2 \underline{E}_p \underline{E}_i \underline{E}_j]_c; \\ T_p \left((\text{ad}_c x_i) x_j (\text{ad}_c x_i)^2 x_k \right) &= a_1 (\text{ad}_c \underline{E}_p) (\text{ad}_c \underline{E}_i) \underline{E}_k [(\text{ad}_c \underline{E}_p) \underline{E}_i, \underline{E}_i]_c \\ &= a_1 [\underline{E}_p \underline{E}_i \underline{E}_j]_c [\underline{E}_p \underline{E}_i^2]_c. \end{aligned}$$

Calculating explicitly the scalars a, b , we notice that $T_p(\mathbf{x}) = a_1 \mathbf{x}' \in \mathcal{J}(s_p^* \chi)$. The case $p = j$ is analogous.

Finally the case $p = i$ follows as the corresponding case in Lemma 3.15.

(ii) The proof is analogous to the previous case. \square

Lemma 3.21. (i) Let $i, j \in \{1, \dots, \theta\}$ be such that $m_{ij}, m_{ji} > 1$. Then, for all p ,

$$T_p \left((1 - \widetilde{q}_{ij}) q_{jj} q_{ji} [E_i, [E_{ij}, E_j]_c]_c - (1 + q_{jj})(1 - q_{jj} \widetilde{q}_{ij}) E_{ij}^2 \right) \in \mathcal{J}(s_p^* \chi).$$

(ii) Let $i, j \in \{1, \dots, \theta\}$ be such that $q_{jj} = -1$, $q_{ii} \widetilde{q}_{ij} \notin \mathbb{G}_6$ or $m_{jj} = 2$, and $q_{ii} \in \mathbb{G}_4$, $m_{ij} = 4$, or $m_{ij} \in \{4, 5\}$. Then, for all p ,

$$T_p \left([E_i, [E_{iij}, E_{ij}]_c]_c - \frac{1 - q_{ii} \widetilde{q}_{ij} - q_{ii}^2 \widetilde{q}_{ij}^2 q_{jj}}{(1 - q_{ii} \widetilde{q}_{ij}) q_{ji}} E_{iij}^2 \right) \in \mathcal{J}(s_p^* \chi).$$

(iii) Let $i, j \in \{1, \dots, \theta\}$ be such that $q_{jj} = -1$, $5\alpha_i + 4\alpha_j \in \Delta_+^\chi$. Let

$$\begin{aligned} v &= \widetilde{q}_{ij}, & a &= (1 - v)(1 - q_{ii}^4 v^3) - (1 - q_{ii} v)(1 + q_{ii}) q_{ii} v \\ b &= (1 - v)(1 - q_{ii}^6 v^5) - a q_{ii} v, \\ d &= \frac{b - (1 + q_{ii})(1 - q_{ii} v)(1 + v + q_{ii} v^2) q_{ii}^6 v^4}{a q_{ii}^3 q_{ij}^2 q_{ji}^3}. \end{aligned}$$

Then, for all p , $T_p \left([E_{2\alpha_i + \alpha_j}, E_{4\alpha_i + 3\alpha_j}]_c - d E_{3\alpha_i + 2\alpha_j}^2 \right) \in \mathcal{J}(s_p^* \chi)$.

Proof. (i) Let \mathbf{x} be the relation we are considering here. We note that if $p \neq i, j$ then $m_{pi} = m_{pj} = 0$, so the proof follows easily. Moreover the conditions about i, j are the same but one relation implies the other holds in $\mathcal{U}(\chi)$ too by Lemma 3.4. Therefore it is enough to consider one of cases $p = i$ or $p = j$; consider $p = j$, in order to apply (66). Note that $m_{pi} = 2, 3$.

If $m_{pi} = 3$, then $m_{ip} = 2$ and we have that

$$T_p \left([E_i, [E_{ip}, E_p]_c]_c \right) = [\underline{E}_{pppi}, \underline{E}_{pi}]_{\underline{c}}.$$

By (9) we can write $T_p(\mathbf{x})$ as a linear combination of

$$\left[\underline{E}_p, [\underline{E}_{ppi}, \underline{E}_{pi}]_{\underline{c}} \right]_{\underline{c}}, \quad \underline{E}_{ppi}^2;$$

note that $\left[\underline{E}_p, [\underline{E}_{ppi}, \underline{E}_{pi}]_{\underline{c}} \right]_{\underline{c}} = [\underline{E}_p^3 \underline{E}_i \underline{E}_p \underline{E}_i]_{\underline{c}}$ if we consider $p < i$. Using the quantum Serre relations or (58), depending on the case (there exist two possible diagrams), $\underline{E}_p^3 \underline{E}_i \underline{E}_p \underline{E}_i$ is expressed as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, so $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$ by Lemma 3.4.

If $m_{pi} = 2$, there exist three diagrams such that $m_{ip} = 2$, and two such that $m_{ip} = 3$. In all the cases,

$$T_p(E_{ip}^2) = a_2^2 \underline{E}_{pi}^2,$$

$$T_p([E_i, [E_{ip}, E_p]_c]_c) = a_2 a_1 [\underline{E}_{ppi}, \underline{E}_i]_{\underline{c}} = a_2 a_1 \left[\underline{E}_p, [\underline{E}_{pi}, \underline{E}_i]_{\underline{c}} \right]_{\underline{c}} + a_2 a_1 \underline{q}_{pi} (\underline{q}_{pp} - \underline{q}_{ii}) \underline{E}_{pi}^2,$$

where we use (9) for the last equality. If $q_{pp} = -\zeta$, $\widetilde{q}_{ip} = \zeta^7$, $q_{ii} = \zeta^3$, for some primitive root $\zeta \in \mathbb{G}_9$, then $s_p^* \chi$ is twist equivalent to χ and (57) is a generator of $\mathcal{J}(s_p^* \chi)$, so $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$ by this relation and Lemma 3.4. For the other braidings $q_{ii} = -1$, so $[\underline{E}_{pi}, \underline{E}_i]_{\underline{c}} \in \mathcal{J}(s_p^* \chi)$ and the coefficient of \underline{E}_{pi}^2 in the expression of $T_p(\mathbf{x})$ is zero. Then $T_p(\mathbf{x}) \in \mathcal{J}(s_p^* \chi)$.

(ii) Let \mathbf{x} be the relation we are considering in this item. First we consider $p = j$; if $q_{pp} = -1$, then $m_{pi} = 1$, so

$$s_p(3\alpha_i + \alpha_p) = 3\alpha_i + 2\alpha_p, \quad s_p(3\alpha_i + 2\alpha_p) = 3\alpha_i + \alpha_p \in \Delta_+^{s_p^* \chi},$$

so $\underline{m}_{ip} \geq 3$. Applying (66) we have that:

$$T_p(E_{iip}^2) = a_1^2 [\underline{E}_{pi}, \underline{E}_i]_{\underline{c}}^2,$$

$$T_p([E_i, [E_{iij}, E_{ij}]_c]_c) = a_1^2 \left[\underline{E}_{pi}, \left[[\underline{E}_{pi}, \underline{E}_i]_{\underline{c}}, \underline{E}_i \right]_{\underline{c}} \right]_{\underline{c}}.$$

As $\underline{m}_{ip} \geq 3$, (58) is a generator of $\mathcal{J}(s_p^* \chi)$, or $\underline{m}_{ip} = 3$, $q_{ii} \notin \mathbb{G}_4$, so $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i^3$ can be written as a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, for the order $p < i$, using the corresponding quantum Serre relation and \underline{E}_p^2 . In both cases we apply Lemma 3.4 to deduce that $T_p(\mathbf{y}) \in \mathcal{J}(s_p^* \chi)$.

If $m_{pi} = 2$, then $m_{ip} = 3$; in this case,

$$T_p(E_{iip}^2) = a_2^2 [\underline{E}_{ppi}, \underline{E}_{pi}]_{\underline{c}}^2,$$

$$T_p([E_i, [E_{iip}, E_{ip}]_c]_c) = a_2^2 [\underline{E}_{ppi}, \underline{E}_{4\alpha_p + 3\alpha_i}]_{\underline{c}}.$$

We have two possibilities for $s_p^* \chi$:

- $\circ\zeta^8 \frac{\zeta^5}{\circ^{-1}}, \zeta \in \mathbb{G}_{24}$, so (63) is a generator of $\mathcal{J}(s_p^*\chi)$,
- $\circ\zeta^5 \frac{-\zeta^{13}}{\circ^{-1}}, \zeta \in \mathbb{G}_{15}$, so (61) and \underline{E}_p^3 are generators of $\mathcal{J}(s_p^*\chi)$.

Then $\underline{E}_p^2 \underline{E}_i \underline{E}_p^2 \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i$ is written as a linear combination of greater words modulo $\mathcal{J}(s_p^*\chi)$ in both cases, so by Lemma 3.4 we have that $T_p(\mathbf{y}) \in \mathcal{J}(s_p^*\chi)$.

Let $p = i$; by Lemma 3.5, it is equivalent to prove that $T_p(\mathbf{y}') \in \mathcal{J}(s_p^*\chi)$, where

$$\mathbf{y}' := [[E_{jp}, [E_{jp}, E_p]_c]_c, E_p]_c - a ([E_{jp}, E_p]_c)^2,$$

and $a \in \mathbf{k}^\times$ is fixed. Note that

$$\begin{aligned} & T_p \left([[(\text{ad}_c E_j)E_p, [(\text{ad}_c E_j)E_p, E_p]_c]_c, E_p]_c \right) \\ &= a_{m_{pi}}^2 a_{m_{pi}-1} \left([(\text{ad}_c \underline{E}_p)^{m_{pi}-1} \underline{E}_j, (\text{ad}_c \underline{E}_p)^{m_{pi}-2} \underline{E}_j]_{\underline{c}} \underline{F}_p \underline{L}_p^{-1} \right. \\ &\quad \left. - q_{jp}^2 q_{pp}^3 \underline{F}_p \underline{L}_p^{-1} [(\text{ad}_c \underline{E}_p)^{m_{pi}-1} \underline{E}_j, (\text{ad}_c \underline{E}_p)^{m_{pi}-2} \underline{E}_j]_{\underline{c}} \right) \\ & T_p ([E_{jp}, E_p]_c) = a_{m_{pi}} a_{m_{pi}-1} (\text{ad}_c \underline{E}_p)^{m_{pi}-2} \underline{E}_i \end{aligned}$$

In any case, $T_p(\mathbf{y}') \in \ker \pi_{s_p^*\chi}$ is a linear combination of

$$[\underline{E}_p^{m_{ip}-1} \underline{E}_i \underline{E}_p^{m_{ip}-3} \underline{E}_i]_{\underline{c}}, \quad [\underline{E}_p^{m_{ip}-2} \underline{E}_i]_{\underline{c}},$$

so by Lemma 3.4, $T_p(\mathbf{y}') \in \mathcal{J}(s_p^*\chi)$, because (57), (respectively, (58), (63)) is a generator of $\mathcal{J}(s_p^*\chi)$ if $m_{pi} = 3$, (respectively, $m_{pi} = 4$, $m_{pi} = 5$).

Finally we take $p \neq i, j$, so p is not connected with i and j (and the proof follows easily by Lemma 3.7), or p is connected only with i , and $q_{ii} = \widetilde{q}_{ij} = q_{pi}^{-1} q_{ip}^{-1} \in \mathbb{G}_4$, $q_{pp} = -1$. Consider the order $p < i < j$, so

$$T_p(E_{ij}) = [\underline{E}_{pi}, \underline{E}_{pij}]_{\underline{c}},$$

$$T_p([E_i, [E_{ij}, E_{ij}]_c]_c) = \left[\underline{E}_{pi}, \left[[\underline{E}_{pi}, \underline{E}_{pij}]_{\underline{c}}, \underline{E}_{pij} \right]_{\underline{c}} \right]_{\underline{c}} = [\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_p \underline{E}_i \underline{E}_j]_{\underline{c}}.$$

By (52), $\underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_p \underline{E}_i$ can be written as a linear combination of other words modulo $\mathcal{J}(s_p^*\chi)$, which are greater than it or they begin with \underline{E}_p ; multiplying on the left by \underline{E}_p , on the right by \underline{E}_j , and using that $\underline{E}_p^2 \in \mathcal{J}(s_p^*\chi)$, $\underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_p \underline{E}_i \underline{E}_j \underline{E}_p \underline{E}_i \underline{E}_j$ is a linear combination of greater words modulo $\mathcal{J}(s_p^*\chi)$, so $T_p(\mathbf{y}) \in \mathcal{J}(s_p^*\chi)$ by a similar argument to the previous steps.

(iii) The proof is analogous to the previous items, where we note that in the two possible cases $q_{jj} = -1$, and if $p \neq i, j$, then p is not connected with them. \square

Lemma 3.22. (i) Let $i, j \in \{1, \dots, \theta\}$ be such that $4\alpha_i + 3\alpha_j \notin \Delta_+^\chi$, $q_{jj} = -1$ or $m_{ji} = 2$, and also $m_{ij} \geq 3$, or $m_{ij} = 2$, $q_{ii} \in \mathbb{G}_3$. Then, $T_p([E_{3\alpha_i+2\alpha_j}, E_{ij}]_c) \in \mathcal{J}(s_p^*\chi)$, for all p .

(ii) Let $i, j \in \{1, \dots, \theta\}$ be such that $4\alpha_i + 3\alpha_j \in \Delta_+^\chi$, $5\alpha_i + 4\alpha_j \notin \Delta_+^\chi$. Then, for all p , $T_p([E_{4\alpha_i+3\alpha_j}, E_{ij}]_c) \in \mathcal{J}(s_p^*\chi)$.

(iii) Let $i, j \in \{1, \dots, \theta\}$ be such that $3\alpha_i + 2\alpha_j \in \Delta_+^\chi$, $5\alpha_i + 3\alpha_j \notin \Delta_+^\chi$, and $q_{ii}^3 \widetilde{q}_{ij}, q_{ii}^4 \widetilde{q}_{ij} \neq 1$. Then, $T_p([E_{ij}, E_{3\alpha_i+2\alpha_j}]_c) \in \mathcal{J}(s_p^*\chi)$ for all p .

(iv) Let $i, j \in \{1, \dots, \theta\}$ be such that $5\alpha_i + 2\alpha_j \in \Delta_+^\chi$, $7\alpha_i + 3\alpha_j \notin \Delta_+^\chi$. Then, for all p , $T_p([E_{ij}, E_{ij}], E_{ij}]_c) \in \mathcal{J}(s_p^*\chi)$.

Proof. For these four sets of conditions, if $p \neq i, j$ then p is not connected with i and j , so the proof follows easily using Lemma 3.7, or we have a diagram as in Lemma 3.21, (ii), and the proof is analogous to this one. In consequence we will consider $p = i$ and $p = j$ for each one of these cases.

(i) Let $\mathbf{x} = [E_{3\alpha_i+2\alpha_j}, E_{ij}]_c$, and take $p = j$. If $m_{pi} = 1$ (that is, $q_{pp} = -1$ or $q_{pp}\widetilde{q_{ip}} = 1$), we have that

$$s_p(3\alpha_i + 2\alpha_p) = 3\alpha_i + \alpha_p \in \Delta_+^{s_p^*\chi}, \quad s_p(4\alpha_i + 3\alpha_p) = 4\alpha_i + \alpha_p \notin \Delta_+^{s_p^*\chi}.$$

Therefore $\underline{m}_{ip} = 3$, so \underline{E}_i^4 (respectively, $(\text{ad}_c \underline{E}_i)^4 \underline{E}_p$) is a generator of $\mathcal{J}(s_p^*\chi)$, if \underline{q}_{ii} belongs (respectively, does not belong) to \mathbb{G}_4 . By (66) and the previous relations, depending on the case,

$$T_p(\mathbf{x}) = a_1^4 \left[\left[[\underline{E}_{pi}, \underline{E}_i]_c, \underline{E}_i \right]_c, \underline{E}_i \right]_c \in \mathcal{J}(s_p^*\chi).$$

The remaining case is $m_{pi} = 2$, for which there exist two possible diagrams:

$$\circ - \zeta \xrightarrow{\zeta^7} \circ \zeta^3, \zeta \in \mathbb{G}_9; \quad \circ - \zeta \xrightarrow{-\zeta^{12}} \circ \zeta^5, \zeta \in \mathbb{G}_{15}.$$

In both cases $q_{pp} \in \mathbb{G}_3$, and also

$$s_p(3\alpha_i + 2\alpha_p) = 3\alpha_i + 4\alpha_p \in \Delta_+^{s_p^*\chi}, \quad s_p(4\alpha_i + 3\alpha_p) = 4\alpha_i + 5\alpha_p \notin \Delta_+^{s_p^*\chi}.$$

Then (61) is a generator of $\mathcal{J}(s_p^*\chi)$ if $3\alpha_i + 5\alpha_p \in \Delta_+^{s_p^*\chi}$, or (62) is a generator of $\mathcal{J}(s_p^*\chi)$ in other case, so for both braidings $E_p^2 E_i E_p E_i E_p E_i E_p E_i$ is a linear combination of greater words modulo $\mathcal{J}(s_p^*\chi)$, and (61) belongs to $\mathcal{J}(s_p^*\chi)$. Therefore

$$T_p(\mathbf{x}) = a_2^4 \left[\left[[\underline{E}_{ppi}, \underline{E}_{pi}]_c, \underline{E}_{pi} \right]_c, \underline{E}_{pi} \right]_c \in \mathcal{J}(s_p^*\chi).$$

Consider now $p = i$, so by Lemma 3.5 it is enough to prove that

$$T_p(\mathbf{x}') \in \mathcal{J}(s_p^*\chi), \quad \mathbf{x}' := [E_{jp}, [E_{jp}, [E_{jp}, E_p]_c]_c]_c.$$

If $m_{pj} = 2$, then $s_p(3\alpha_p + 2\alpha_j) = \alpha_p + 2\alpha_j \in \Delta_+^{s_p^*\chi}$, $s_p(4\alpha_p + 3\alpha_j) = 2\alpha_p + 3\alpha_j \notin \Delta_+^{s_p^*\chi}$, so $\underline{m}_{jp} = 2$, and (45) is a generator of $\mathcal{J}(s_p^*\chi)$; then

$$T_p(\mathbf{x}') = a_2^3 a_1 \left[\underline{E}_{pj}, [\underline{E}_{pj}, \underline{E}_j]_c \right]_c \in \mathcal{J}(s_p^*\chi).$$

If $m_{pj} = 3$, then $s_p(3\alpha_p + 2\alpha_j) = 3\alpha_p + 2\alpha_j \in \Delta_+^{s_p^*\chi}$, $s_p(4\alpha_p + 3\alpha_j) = 5\alpha_p + 3\alpha_j \notin \Delta_+^{s_p^*\chi}$, so (62) is a generator of $\mathcal{J}(s_p^*\chi)$. By (66),

$$T_p(\mathbf{x}') = a_3^3 a_2 \left[\underline{E}_{ppj}, [\underline{E}_{ppj}, \underline{E}_{pj}]_c \right]_c \in \mathcal{J}(s_p^*\chi).$$

If $m_{pj} = 4$, then $s_p(3\alpha_p + 2\alpha_j) = 5\alpha_p + 2\alpha_j \in \Delta_+^{s_p^*\chi}$; moreover, we note that $7\alpha_p + 3\alpha_j \notin \Delta_+^{s_p^*\chi}$ in both cases, and (60) is a generator of $\mathcal{J}(s_p^*\chi)$. By this relation and the quantum Serre relations, $\underline{E}_p^3 \underline{E}_j \underline{E}_p^3 \underline{E}_j \underline{E}_p^2 \underline{E}_j$ is written as a linear combination of greater words modulo $\mathcal{J}(s_p^*\chi)$, so

$$T_p(\mathbf{x}') = a_4^3 a_3 \left[\underline{E}_{pppj}, [\underline{E}_{pppj}, \underline{E}_{ppj}]_c \right]_c \in \mathcal{J}(s_p^*\chi).$$

(ii) The proof is similar to (i), but more simple: for $p = j$ we have just one possibility, $m_{pi} = 1$.

(iii) Let $\mathbf{x} = [E_{ii}, E_{3\alpha_i+2\alpha_j}]_{\underline{c}}$. Consider $p = j$; we consider i a non-Cartan vertex, because in other case $\underline{E}_i \underline{E}_p \underline{E}_i^2$ or $\underline{E}_i^2 \underline{E}_p \underline{E}_i^2$ can be written as a linear combination of other words using the corresponding quantum Serre relation, and finally $\underline{E}_i^2 \underline{E}_p \underline{E}_i^2 \underline{E}_p \underline{E}_i$ is a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, so the relation is redundant in this case. In consequence we consider $m_{pi} = 1$, and for this case $\underline{E}_p \underline{E}_i^2 \underline{E}_p \underline{E}_i^3$ is a linear combination of greater words modulo $\mathcal{J}(s_p^* \chi)$, so

$$T_p(\mathbf{x}) = \left[[\underline{E}_{pi}, \underline{E}_i]_{\underline{c}}, \left[[\underline{E}_{pi}, \underline{E}_i]_{\underline{c}}, \underline{E}_i \right]_{\underline{c}} \right] \in \mathcal{J}(s_p^* \chi).$$

Let $p = i$; as above, it is enough to prove that

$$T_p(\mathbf{x}') \in \mathcal{J}(s_p^* \chi), \quad \mathbf{x}' := [[E_{jp}, [E_{jp}, E_p]_{\underline{c}}], [E_{jp}, E_p]_{\underline{c}}]_{\underline{c}}.$$

If $m_{pj} = 2$, then $s_p(3\alpha_p + 2\alpha_j) = \alpha_p + 2\alpha_j \in \Delta_+^{s_p^* \chi}$, $s_p(5\alpha_p + 3\alpha_j) = \alpha_p + 3\alpha_j \notin \Delta_+^{s_p^* \chi}$, so $(\text{ad}_{\underline{c}} \underline{E}_j)^3 \underline{E}_p$ (or \underline{E}_i^3) is a generator of $\mathcal{J}(s_p^* \chi)$; therefore

$$T_p(\mathbf{x}') = a_2^3 a_1^2 \left[[\underline{E}_{pj}, \underline{E}_j]_{\underline{c}}, \underline{E}_j \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

If $m_{pj} = 3$, then $s_p(3\alpha_p + 2\alpha_j) = 3\alpha_p + 2\alpha_j \in \Delta_+^{s_p^* \chi}$, $s_p(5\alpha_p + 3\alpha_j) = 4\alpha_p + 3\alpha_j \notin \Delta_+^{s_p^* \chi}$, so (61) is a generator of $\mathcal{J}(s_p^* \chi)$. Therefore

$$T_p(\mathbf{x}') = a_3^3 a_2^2 \left[[\underline{E}_{ppj}, \underline{E}_{pj}]_{\underline{c}}, \underline{E}_{pj} \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

If $m_{pj} = 4$, then $s_p(3\alpha_p + 2\alpha_j) = 5\alpha_p + 2\alpha_j \in \Delta_+^{s_p^* \chi}$, $s_p(5\alpha_p + 3\alpha_j) = 7\alpha_p + 3\alpha_j \notin \Delta_+^{s_p^* \chi}$, so (60) is a generator of $\mathcal{J}(s_p^* \chi)$. In consequence,

$$T_p(\mathbf{x}') = a_4^3 a_3^2 \left[[\underline{E}_{pppj}, \underline{E}_{ppj}]_{\underline{c}}, \underline{E}_{ppj} \right]_{\underline{c}} \in \mathcal{J}(s_p^* \chi).$$

(iv) The proof is analogous to the previous one. \square

Now we are ready to prove that the Lusztig isomorphisms descend to the family of algebras $U(\chi)$, so we will look at the root system of this family of algebras. As we consider finite root systems, they are univocally determined as sets of real roots, and using this result we will obtain the desired Theorem of presentation by generators and relations of Nichols algebras.

Proposition 3.23. *The morphisms (38) induce algebra isomorphisms*

$$T_p, T_p^- : U(\chi) \rightarrow U(s_p^* \chi),$$

such that $T_p T_p^- = T_p^- T_p = \text{id}_{U(\chi)}$.

Proof. By the definition of the ideals $\mathcal{J}(\chi)$ and the previous Lemmata, $T_p(\mathcal{J}(\chi)) \subset \mathcal{J}(s_p^* \chi)$, so there exists an algebra morphism $T_p : U(\chi) \rightarrow U(s_p^* \chi)$. By $\phi_4^2 = \text{id}$ and the definition of the ideal, $\phi_4(\mathcal{J}(\chi)) = \mathcal{J}(\chi)$, and also $\varphi_\lambda(\mathcal{J}(\chi)) = \mathcal{J}(\chi)$ for any $\lambda \in (\mathbf{k}^\times)^\theta$, because the ideal is \mathbb{Z}^θ -homogeneous. By (39) we have that $T_p^-(\mathcal{J}(\chi)) \subset \mathcal{J}(s_p^* \chi)$, so there exists also an algebra morphism $T_p^- : U(\chi) \rightarrow U(s_p^* \chi)$, induced by the corresponding morphism.

These algebras are generated by E_i, F_i, L_i, K_i , and the identities $T_p T_p^- = T_p^- T_p = \text{id}$ hold for each one of these elements, so these identities hold for all the elements of these algebras, and these morphisms are isomorphisms. \square

This result lets us to prove the main result of this Section. The proof is similar to the one for [A1, Thm.5.25].

Proof of Theorem 3.1.

Set $\underline{\Delta}_+^\chi := \Delta^+(U^+(\chi)) \setminus \{N_\alpha \alpha : \alpha \in \Delta_+^\chi\}$. By the triangular decomposition, Lemma 2.8, Theorem 2.9 and Proposition 3.23, we have that

$$(67) \quad \mathcal{H}_{U^+(\chi)} = \mathcal{H}_{U_{+p}^+(\chi)} \mathfrak{q}_{h(E_p)} = s_p(\mathcal{H}_{U_{+p}^+(s_p^* \chi)}) \mathfrak{q}_{h(E_p)},$$

for all $p \in \{1, \dots, \theta\}$, because $\deg(T_p(X)) = s_p(\deg X)$ for each homogeneous element $X \in \mathcal{U}(\chi)$. Recall that $h(E_p) \in \{\text{ord } q_{pp}, \infty\}$, so

$$(68) \quad \Delta^+(U^+(\chi)) = s_p(\Delta^+(U^+(s_p^* \chi)) \setminus \{\alpha_p, N_p \alpha_p\}) \cup S_p,$$

where $S_p = \{\alpha_p\}$, or $S_p = \{\alpha_p, N_p \alpha_p\}$, so $\underline{\Delta}_+^\chi = s_p(\underline{\Delta}_+^{s_p^* \chi} \setminus \{\alpha_p\}) \cup \{\alpha_p\}$.

In this way, if we consider the sets $\underline{\Delta}_+^\chi$, for each χ in a Weyl equivalence class of a fixed braiding with finite root system, then $R = \{\underline{\Delta}_+^\chi\}$ is a root system for our Weyl groupoid, according with the Definition 1.17. As we have a finite root system, it follows that $\underline{\Delta}_+^\chi = \Delta_+^\chi$, for all χ , because by Proposition 1.21 all the roots are real. In this way, $\Delta^+(U^+(\chi))$ is obtained from Δ_+^χ adding $N_\alpha \alpha$, for some $\alpha \in \Delta_+^\chi$. Fix an order on the letters x_i and consider the corresponding PBW basis. We have a projection $\pi_\chi : U(\chi) \rightarrow \mathfrak{u}(\chi)$ of graded braided Hopf algebras, so the corresponding x_α of the PBW basis of $\mathfrak{u}(\chi)$ are generators of the PBW basis of $U(\chi)$, by the definition of Kharchenko's PBW basis. On the other hand, each simple root of a non-Cartan vertex satisfies $E_i^{N_i} = 0$ in $U(\chi)$, so $N_i \alpha_i \notin \Delta^+(U^+(\chi))$. Therefore (67) implies that

$$N_\alpha \alpha \notin \Delta^+(U^+(\chi)), \quad \text{for all } \alpha \in \Delta_+^\chi \setminus \mathcal{O}(\chi),$$

because α is of the way $\alpha = w(\alpha_i)$ for some $w \in \mathcal{W}$ and $i \in \{1, \dots, \theta\}$, i a non-Cartan vertex in the corresponding χ' . Analogously, for each Cartan vertex i , $N_i \alpha_i \in \Delta^+(U^+(\chi))$, because $E_i^{N_i} \neq 0$ in $U(\chi)$, so

$$N_\alpha \alpha \in \Delta^+(U^+(\chi)), \quad \text{for all } \alpha \in \mathcal{O}(\chi).$$

Therefore $\Delta^+(U^+(\chi)) = \Delta_+^\chi \cup \{N_\alpha \alpha : \alpha \in \mathcal{O}(\chi)\}$.

Suppose that the degree $N_\alpha \alpha$ in $\Delta^+(U^+(\chi))$ corresponds to a Lyndon word of this degree: we can assume that it is of minimal length, and we denote it by u ; set $(v, w) = \text{Sh}(u)$. In this way, $\deg v = \beta$, $\deg w = \gamma$, for some $\beta, \gamma \in \Delta_+^\chi$, and $\beta + \gamma = N_\alpha \alpha$. As all the roots are real, we deduce that if $\beta < \gamma$, then $\beta < \alpha < \gamma$, by a similar argument to the convexity properties in [A2]. We can consider then the case $\beta = \alpha_i$, because if $\beta = s_{i_1} \cdots s_{i_k}(\alpha_{i_{k+1}})$, where $w = s_{i_1} \cdots s_{i_k}$ is the beginning of the expression of the element of maximal length, we apply w^{-1} to obtain that $\alpha_{i_{k+1}} + \gamma' = N_\alpha \alpha'$ for some $\alpha', \gamma' \in \Delta_+^\chi$. Note also that $N_\alpha > 2$, because if we suppose $N_\alpha = 2$, then α is applied in a simple root α_i corresponding a Cartan vertex by some element of the Weyl groupoid, and as N_α is invariant by the action of the Weyl groupoid, it should be $q_{ii} = -1$, but it corresponds to an isolated vertex or a non-Cartan vertex, which is a contradiction. Set

$$\alpha = \sum_{j=1}^{\theta} n_j \alpha_j, \quad \gamma = \sum_{j=1}^{\theta} m_j \alpha_j, \quad \text{for some } n_j, m_j \in \mathbb{N}_0.$$

Note that $m_i = N_\alpha n_i - 1 \geq 2$, and for $j \neq i$, $m_j = N_\alpha n_j \geq 3$, so $\text{supp } \gamma = \text{supp } \alpha$. By simplicity assume that $\text{supp } \alpha = \{1, \dots, \theta\}$; note that the vertices of $\text{supp } \beta$ corresponds to a connected subdiagram, for any positive root β .

By these considerations we reduce the problem to an analysis case by case of the positive root systems of connected diagrams, and we do it with the help of the program

SARNA [GHV]. We look for the possible γ such that all the coordinates, except at most one, are divisible by an integer ≥ 3 , and the remaining coordinate is ≥ 2 , so we just have a few 3-uples in rank two or three. Analyzing each of these 3-uples

$$(\alpha, \gamma, i) \in \Delta_+^\chi \times \Delta_+^\chi \times \{1, \dots, \theta\} \text{ such that there exists } N \in \mathbb{N} : \alpha_i + \gamma = N\alpha,$$

we note that $N \neq N_\alpha$ for all of them. Therefore, there are no Lyndon words of degree $N_\alpha\alpha$, so the generators of degree $N_\alpha\alpha$ are $x_\alpha^{N_\alpha}$, and then the elements

$$x_{\beta_1}^{n_1} \cdots x_{\beta_k}^{n_k}, \quad \beta_i \in \Delta_+^\chi, \quad \begin{cases} 0 \leq n_j < N_j, & \text{si } \beta_j \notin \mathcal{O}(\chi) \\ 0 \leq n_j < \infty, & \text{si } \beta_j \in \mathcal{O}(\chi) \end{cases}$$

are a PBW basis of $U(\chi)$. As $x_\alpha^{N_\alpha} = 0$ in $\mathfrak{u}^+(\chi)$, π_χ induces a surjective morphism

$$\pi'_\chi : U(\chi) / \langle x_\alpha^{N_\alpha} : \alpha \in \mathcal{O}(\chi) \rangle \longrightarrow \mathfrak{u}^+(\chi),$$

which applies the set

$$\{x_{\beta_1}^{n_1} \cdots x_{\beta_k}^{n_k}, \quad \beta_i \in \Delta_+^\chi, 0 \leq n_j < N_j\},$$

generating linearly the quotient, to the corresponding PBW basis of $\mathfrak{u}^+(\chi)$. Therefore π'_χ is an isomorphism. \square

4. GENERATION IN DEGREE ONE

Now we answer positively the Conjecture 1, formulated by Andruskiewitsch and Schneider, but restricting to the case in which $G(H)$ is abelian. The technique of the proof is the same that these authors use in [AS4], extended in some works to other families. In particular, the first Lemmata of this Section correspond to relations generating the ideal for standard braidings as in [AGI], but the proof is made in a general context.

In what follows Γ denotes a finite abelian group, and $S = \bigoplus_{n \geq 0} S(n)$ is a graded braided Hopf algebra in ${}^G_C\mathcal{YD}$ such that $S(0) = \mathbf{k}\mathbf{1}$, generated as an algebra by $V := S(1)$. Fix a basis $\{x_1, \dots, x_\theta\}$ of V , so V has a braiding of diagonal type: we can assume that $x_i \in S(1)_{g_i}^{\chi_i}$ for some $g_i \in \Gamma$ and $\chi_i \in \widehat{\Gamma}$. Set then $q_{ij} := \chi_j(g_i) \in \mathbf{k}^\times$.

We will prove that if S is finite dimensional, then S is the Nichols algebra $\mathcal{B}(V)$ associated to V . We will obtain then the main Theorem of this Section, answering this Conjecture.

We begin by extending [AS4, Lemma 5.4] for a general quantum Serre relation, proving that they hold in S , or S is infinite-dimensional.

Proposition 4.1. *Let S be as above. If there exist $i, j \in \{1, \dots, \theta\}$ such that $q_{ii}^{m_{ij}+1} \neq 1$, and also $\text{ad}_c(x_i)^{1+m_{ij}}(x_j) \neq 0$, then S is infinite-dimensional.*

Proof. By definition of m_{ij} , we have that $q_{ii}^{m_{ij}} \widetilde{q}_{ij} = 1$. We begin the proof as in [AS4, Lemma 5.4]. To simplify the notation, call $m = m_{ij}$, $q = q_{ii}$, $y_1 := x_i$, $y_2 := x_j$, $y_3 := \text{ad}_c(x_i)^{1+m}(x_j)$. Set also

$$\begin{aligned} h_1 &= g_i, & h_2 &= g_j, & h_3 &= g_i^{m+1} g_j, \\ \eta_1 &= \chi_i, & \eta_2 &= \chi_j, & \eta_3 &= \chi_i^{m+1} \chi_j, \end{aligned}$$

so $y_k \in S_{h_k}^{\eta_k}$, $1 \leq k \leq 3$. If $W = \mathbf{k}y_1 + \mathbf{k}y_2 + \mathbf{k}y_3$, then $W \subset \mathcal{P}(S)$ (because y_3 is also primitive), so there exists a monomorphism $\mathcal{B}(W) \hookrightarrow S$. We compute the corresponding

braiding matrix $(Q_{kl} = \eta_l(h_k))_{1 \leq k, l \leq 3}$, and consider the corresponding generalized Dynkin diagram:

$$(69) \quad \begin{array}{ccc} & \circ^{q_{jj}} & \\ q^{-m} \swarrow & & \searrow q^{-m(m+1)} q_{jj}^2 \\ \circ^q & \xrightarrow{q^{m+2}} & \circ^{q^{m+1}} q_{jj} \end{array}$$

We will consider the different possible cases and prove that no one of them are in [H3], so $\mathcal{B}(W)$, and in consequence S , is infinite-dimensional. Suppose that the diagram (69) is in Heckenberger's list:

Case I: $Q_{kl}Q_{lk} \neq 1$ for all $1 \leq k < l \leq 3$. By [H3, Lemma 9], $1 = \prod_{k < l} Q_{kl}Q_{lk} = q^{2-m(m+1)} q_{jj}^2$, and at least one of the vertices is labeled with -1 . Note that $q \neq -1$, because in such case $m = 0$ (and we assume $q^{m+1} \neq 1$). Also $q_{jj} \neq q^{m+1} q_{jj}$ by hypothesis, so there is only one vertex labeled with -1 .

- If $q_{jj} = -1$, then $1 = (q^{m+1} q_{jj})(q^{-m(m+1)} q_{jj}^2) = -q^{1-m^2}$ and $m = 1$ by the same Heckenberger's Lemma, which is a contradiction.
- If $q^{m+1} q_{jj} = -1$, then $1 = qq^{m+2} = q^{m+3}$ by the same result, and also

$$1 = q_{jj}(q^{-m(m+1)} q_{jj}^2) = q_{jj}^3 q^{-m(m+3)+2m} = q_{jj}^3 q^{2m},$$

so we deduce that

$$-1 = (-1)^3 = q_{jj}^3 q^{3m+3} = (q_{jj}^3 q^{2m}) q^{m+3} = q^{m+3},$$

which is also a contradiction. Therefore (69) does not belong to Heckenberger list for this case.

Case II: $Q_{12}Q_{21} = q^{-m} = 1$. In this case $m = 0$, so (69) becomes:

$$(70) \quad \circ^q \xrightarrow{q^2} \circ^{qq_{jj}} \xrightarrow{q_{jj}^2} \circ^{q_{jj}}.$$

If $q_{jj} = -1$ we obtain the connected subdiagram $\circ^q \xrightarrow{q^2} \circ^{-q}$, which has no vertices labeled with -1 , and these labels are different. Such diagram is not of finite Cartan type and moreover it does not correspond to any diagram without -1 in the vertices in rows 5, 9, 11, 12, 15 of [H3, Table 1], so $\mathcal{B}(W)$ is infinite-dimensional.

If $q_{jj} \neq -1$ and $q = -1$, we have an analogous situation, so $q \neq -1$ and (70) is a connected diagram of three vertices. If $qq_{jj} \neq -1$, then [H3, Lemma 9] implies that one of the following situations holds:

- the diagram is of finite Cartan type, so it contains a subdiagram of Cartan type A_2 . Then $1 = qq^2 = (qq_{jj})q^2$, or $1 = q_{jj}q_{jj}^2 = (qq_{jj})q_{jj}^2$, so $q = 1$ or $q_{jj} = 1$;
- $q^3 = 1$, $q_{jj}, q_{jj}q \in \mathbb{G}_6 \cup \mathbb{G}_9$, and $q_{jj}q_{jj}^2 = 1$ or $q_{jj}^3 = 1$, $q, q_{jj}q \in \mathbb{G}_6 \cup \mathbb{G}_9$, $qq^2 = 1$.

No one of these situations hold, so $qq_{jj} = -1$. If this diagram is in [H3, Table 2], it follows that $Q_{ii}Q_{i2}Q_{2i} = 1$ for some $i \in \{1, 3\}$ in all the possible cases. We can assume then $i = 1$, $q^3 = 1$. By [H3, Lemma 9], one of the following situations holds:

- $q_{jj}^3 = 1$, but also $q_{jj}^3 = -q^{-3} = -1$,
- $q_{jj}^4 = 1$,
- $q_{jj} = -q$.

No one of these situations are possible, so we obtain a contradiction in this case too.

Case III: $Q_{13}Q_{31} = q^{m+2} = 1$. We obtain the diagram:

$$\circ^q \xrightarrow{q^2} \circ^{q_{jj}} \xrightarrow{q^{-2}q_{jj}^2} \circ^{q^{-1}q_{jj}}.$$

Such diagram is the corresponding to (70), but changing q_{jj} by $q_{jj}q^{-1}$, so it does not belong to [H3, Table 2]. Then $q^{m+2} \neq 1$.

Case IV: $Q_{23}Q_{32} = 1$. This means $q_{jj}^2 = q^{m(m+1)}$, so we have the diagram:

$$(71) \quad \circ^{q_{jj}} \xrightarrow{q^{-m}} \circ^q \xrightarrow{q^{m+2}} \circ^{q^{m+1}q_{jj}}.$$

This diagram is connected by the previous cases. As $m \neq 0$, $q^{m+1} \neq 1$, it follows that $q \neq -1$. Consider the different possible values of the labels of the vertices:

$\mathbf{q}_{jj} = \mathbf{q}^{m+1}\mathbf{q}_{jj} = -\mathbf{1}$: that is, $q^{m+1} = 1$ and we have the diagram:

$$\circ^{-1} \xrightarrow{q} \circ^q \xrightarrow{q} \circ^{-1},$$

which is not in Heckenberger's list.

$\mathbf{q}_{jj} = -\mathbf{1}, \mathbf{q}^{m+1}\mathbf{q}_{jj} \neq -\mathbf{1}$: By [H3, Table 2], it should be $1 = Q_{22}Q_{23}Q_{32} = q^{m+3}$, and we should have the diagram

$$\circ^{-q^{-2}} \xrightarrow{q^{-1}} \circ^q \xrightarrow{q^3} \circ^{-1}.$$

Moreover, $1 = q_{jj}^2 = q^{m(m+1)} = q^{2m} = q^{-6}$. Note that $q^3 \neq 1$ because $q^m \neq 1$, so $q \in \mathbb{G}_6$. But this diagram is not in Heckenberger's list.

$\mathbf{q}_{jj} \neq -\mathbf{1}, \mathbf{q}^{m+1}\mathbf{q}_{jj} = -\mathbf{1}$: as above, $1 = Q_{22}Q_{21}Q_{12} = q^{1-m}$. By definition it should be $m = 1$, with the same diagram of the previous case and $q \in \mathbb{G}_6$, so we obtain the same contradiction.

$\mathbf{q}_{jj}, \mathbf{q}^{m+1}\mathbf{q}_{jj} \neq -\mathbf{1}$: By [H3, Lemma 9], one of the following situations holds:

- the diagram is of Cartan type. Then, $q = q_{jj}$ and $m = 1$, or $q = q^{m+1}q_{jj} = q^{-m-2}$. In both cases we obtain the same diagram,

$$\circ^q \xrightarrow{q^{-1}} \circ^q \xrightarrow{q^3} \circ^{q^3}.$$

We discard easily the cases A_3, C_3 , because $q, q^2 \neq q^3$. If it is of type B_3 , $q = (q^3)^2 = q^{-3}$, which is a contradiction.

- $q_{jj} \in \mathbb{G}_3$, $q \in \mathbb{G}_6 \cup \mathbb{G}_9$ and $1 = q^{1-m} = q_{jj}q^{2m+3}$. Then $m = 1$ and $q^5 = q_{jj}^{-1}$, so $q^{15} = 1$, but we obtain then a contradiction with the fact that $q \in \mathbb{G}_6 \cup \mathbb{G}_9$ is primitive.
- $q^{m+1}q_{jj} \in \mathbb{G}_3$, $q \in \mathbb{G}_6 \cup \mathbb{G}_9$ y $1 = q_{jj}q^{-m} = q^{m+3}$. Again $q^{15} = 1$, and we obtain the same contradiction.

In consequence, (69) is not in Heckenberger's list, and S is infinite-dimensional. \square

Now we continue with another Lemmata from [AGI], just adapted to this general context.

Lemma 4.2. *Let $j, k, l \in \{1, \dots, \theta\}$ be such that $q_{kk} = -1$, $\widetilde{q}_{kj} = \widetilde{q}_{kl}^{-1} \neq 1$, $\widetilde{q}_{jl} = 1$. If $[x_{jkl}, x_k]_c \neq 0$ is a primitive element of S , then S is infinite-dimensional.*

Proof. Set $u := [x_{jkl}, x_k]_c$, $g_u := g_j g_k^2 g_l \in \Gamma$, $\chi_u := \chi_j \chi_k^2 \chi_l \in \widehat{\Gamma}$, $q := \widetilde{q}_{lk}$; we work then as in the previous Lemma.

We compute the braiding corresponding to the primitive elements $y_1 = x_j$, $y_2 = x_k$, $y_3 = x_l$ and $y_4 = u$, with the corresponding elements $h_i \in \Gamma$, $\eta_i \in \widehat{\Gamma}$; we will prove that

such braiding has an associated Nichols algebra of infinite dimension, and so S has infinite dimension. The corresponding generalized Dynkin diagram to $(Q_{rs} = \eta_s(h_r))_{1 \leq r, s \leq 4}$ is:

$$(72) \quad \begin{array}{ccc} \circ^{q_{jj}} & \xrightarrow{q^{-1}} & \circ^{-1} \\ q_{jj}^2 q^{-2} \Big| & & \Big| q \\ \circ^{q_{jj} q_{ii}} & \xrightarrow{q_{ii}^2 q^2} & \circ^{q_{ii}} \end{array}.$$

Suppose that such diagram is in Heckenberger's list. If $q = -1$, then (72) contains (70) as a subdiagram, so it does not appear in the list. Therefore $q \neq -1$. As each diagram in [H3, Table 3] does not contain a 4-cycle, it follows that $q_{jj}^2 q^{-2} = 1$, or $q_{ii}^2 q^2 = 1$. As the conditions are symmetric, it is enough to consider the case $q_{jj} = \pm q$.

If we also have $q_{ii} = \pm q^{-1}$, and as $Q_{44} = q_{jj} q_{ii} \neq 1$, the diagram contains the following

$$\circ^q \xrightarrow{q^{-1}} \circ^{-1} \xrightarrow{q} \circ^{-q^{-1}},$$

which is a contradiction with [H3, Lemma 9]. In consequence we have:

$$\circ^{\pm q} \xrightarrow{q^{-1}} \circ^{-1} \xrightarrow{q} \circ^{q_{ii}} \xrightarrow{q_{ii}^2 q^2} \circ^{q_{jj} q_{ii}}.$$

Suppose that $q_{jj} = -q$. As $Q_{11} Q_{12} Q_{21} \neq 1$, we deduce from [H3, Table 3] that $m_{12} = 2$; that is,

$$0 = (1 - Q_{11}^3)(Q_{11}^2 Q_{12} Q_{21} - 1) = (1 + q^3)(q - 1),$$

which gives conditions about q , but each diagram in [H3, Lemma 9] does not satisfy this condition.

Therefore $q_{jj} = q$. We look at [H3, Table 3] but a diagram in such list does not satisfy $Q_{22} = -1$, $Q_{11} = Q_{44} Q_{33}^{-1} = q \neq \pm 1$, so (72) is not in the list. In consequence, S has infinite dimension. \square

Lemma 4.3. (i) *Let $i, j \in \{1, \dots, \theta\}$ be such that $q_{jj} = -1$, $q_{ii} \widetilde{q}_{ij} \in \mathbb{G}_6$, and also $q_{ii} \in \mathbb{G}_3$ or $m_{ij} \geq 3$. If $[x_{iij}, x_{ij}]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

(ii) *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = \pm \widetilde{q}_{ij} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, and also $-q_{jj} = \widetilde{q}_{ij} \widetilde{q}_{jk} = 1$ or $q_{jj}^{-1} = \widetilde{q}_{ij} = \widetilde{q}_{jk} \neq -1$. If $[x_{iijk}, x_{ij}]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

Proof. (i) We follow the same scheme of proof. Set

$$y_1 = x_i, \quad y_2 = x_j, \quad y_3 = [x_{iij}, x_{ij}]_c,$$

and $h_i \in \Gamma$, $\eta_i \in \widehat{\Gamma}$, $i = 1, 2, 3$ the corresponding elements. Suppose that the braiding matrix $(Q_{rs} = \eta_s(h_r))_{1 \leq r, s \leq 3}$ appears in Heckenberger's list. The associated generalized Dynkin diagram is

$$\begin{array}{ccc} \circ^{q_{ii}} & \xrightarrow{q} & \circ^{-1} \\ & \searrow^{q^2} & \nearrow^{q^3} \\ & \circ^{q_{ii}^3} & \end{array}, \quad q := \widetilde{q}_{ij}.$$

Then $Q_{33} = q_{ii}^3 \neq 1$, so $m_{ij} \geq 3$. Moreover the diagram is connected, so it is of type super $G(3)$, the unique diagram of rank three such that some m_{rs} is ≥ 3 . Therefore $1 = Q_{23} Q_{32} = q^3$, which is a contradiction, so the diagram associated to (Q_{rs}) does not correspond to a finite-dimensional Nichols algebra. In consequence S is infinite-dimensional.

(ii) Set $w := [x_{iijk}, x_{ij}]_c$, and denote as above $y_1 = x_i$, $y_2 = x_j$, $y_3 = x_k$, $y_4 = w$, W the subspace generated by these elements, and $h_i \in \Gamma$, $\eta_i \in \widehat{\Gamma}$, $i = 1, 2, 3, 4$ the

corresponding elements: suppose again that $\mathcal{B}(W)$ is a finite-dimensional Nichols algebra. Set $\zeta = q_{ii} \in \mathbb{G}_3$. We analyze each possible case.

- $q_{jj} = -1$, $\widetilde{q}_{ij}\widetilde{q}_{jk} = 1$: the diagram of (Q_{rs}) becomes

$$\begin{array}{ccccc} \circ\zeta & \xrightarrow{\pm\zeta} & \circ^{-1} & \xrightarrow{\mp\zeta^2} & \circ^{q_{kk}} \\ & \searrow^{\zeta^2} & & \nearrow_{q_{kk}^2\zeta} & \\ & & \circ^{q_{kk}\zeta} & & \end{array} .$$

As $Q_{12}Q_{21}$, $Q_{14}Q_{41}$, $Q_{42}Q_{24} \neq 1$, and the product of these three scalars is not 1, such diagram is not in Heckenberger's list, by [H3, Lemma 9].

- $q_{jj}^{-1} = \widetilde{q}_{ij} = \widetilde{q}_{jk} \neq -1$: now we have the diagram

$$\begin{array}{ccccc} \circ\zeta & \xrightarrow{\pm\zeta} & \circ\pm\zeta^2 & \xrightarrow{\pm\zeta} & \circ^{q_{kk}} \\ & \searrow^{\zeta^2} & & \nearrow_{q_{kk}^2\zeta^2} & \\ & & \circ^{q_{kk}\zeta} & & \end{array} .$$

The lack of 4-cycles in Heckenberger's list implies that $1 = Q_{34}Q_{43} = q_{kk}^2\zeta^2$, so $q_{kk}\zeta = -1$, because $Q_{44} = q_{kk}\zeta \neq 1$. But this diagram does not appear in [H3, Table 3].

We obtain a contradiction in all the cases, so S is infinite-dimensional. \square

Lemma 4.4. *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $\widetilde{q}_{ik}, \widetilde{q}_{ij}, \widetilde{q}_{jk} \neq 1$. Let*

$$w := x_{ijk} - \frac{1 - \widetilde{q}_{jk}}{q_{kj}(1 - \widetilde{q}_{ik})} [x_{ik}, x_j]_c - q_{ij}(1 - \widetilde{q}_{jk}) x_j x_{ik}.$$

If $w \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.

Proof. Set $y_1 = x_i$, $y_2 = x_j$, $y_3 = x_k$, $y_4 = w$, W the subspace generated by these elements, $h_i \in \Gamma$, $\eta_i \in \widehat{\Gamma}$, $i = 1, 2, 3, 4$ the corresponding elements, and $(Q_{rs} = \eta_s(h_r))_{1 \leq r, s \leq 4}$: suppose as above that $\mathcal{B}(W)$ is finite dimensional. Note that

$$Q_{14}Q_{41} = q_{ii}^2 \widetilde{q}_{ij} \widetilde{q}_{ik} = q_{ii}^2 \widetilde{q}_{jk}^{-1},$$

because $\widetilde{q}_{ij}\widetilde{q}_{ik}\widetilde{q}_{jk} = 1$, by [H3, Lemma 9]. By the same Lemma at least one vertex is labeled with -1 . Then, if $q_{ii} = -1$, we have that $Q_{14}Q_{41} \neq -1$; the same holds for the other vertices, so exactly one vertex is labeled with -1 (we have no 4-cycles). We look for possible braiding with these conditions in [H3, Table 3], but no one coincides with this description. Therefore $\mathcal{B}(W)$ is infinite-dimensional, and S too. \square

Lemma 4.5. (i) *Let $i, j, k \in \{1, \dots, \theta\}$ be such that one of the following conditions holds:*

- $q_{ii} = q_{jj} = -1$, $\widetilde{q}_{ij}^2 = \widetilde{q}_{jk}^{-1}$, $\widetilde{q}_{ik} = 1$, or
- $\widetilde{q}_{ij} = q_{jj} = -1$, $q_{ii} = -\widetilde{q}_{jk}^2 \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, or
- $q_{kk} = \widetilde{q}_{jk} = q_{jj} = -1$, $q_{ii} = -\widetilde{q}_{ij} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$, or
- $q_{jj} = -1$, $\widetilde{q}_{ij} = q_{ii}^{-2}$, $q_{kk} = \widetilde{q}_{jk}^{-1} = -q_{ii}^3$, $\widetilde{q}_{ik} = 1$, or
- $q_{ii} = q_{jj} = q_{kk} = -1$, $\pm\widetilde{q}_{ij} = \widetilde{q}_{jk} \in \mathbb{G}_3$, $\widetilde{q}_{ik} = 1$,

If $[[x_{ij}, x_{ijk}]_c, x_j]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.

(ii) *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = q_{jj} = -1$, $(\widetilde{q}_{ij})^3 = (\widetilde{q}_{jk})^{-1}$, $\widetilde{q}_{ik} = 1$. If $[[x_{ij}, [x_{ij}, x_{ijk}]_c]_c, x_j]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

Proof. (i) Set $y_1 = x_i, y_2 = x_j, y_3 = x_k, y_4 = [[x_{ij}, x_{ijk}]_c, x_j]_c$, W the subspace generated by these elements, $h_i \in \Gamma, \eta_i \in \widehat{\Gamma}, i = 1, 2, 3, 4$ the associated elements and $(Q_{rs} = \eta_s(h_r))_{1 \leq r, s \leq 4}$ the braiding matrix. We will consider the associated generalized Dynkin diagram for each case.

For the first case, we have the following diagram, where $q := \widetilde{q}_{ij}$:

$$\begin{array}{ccccc} \circ^{-1} & \xrightarrow{q} & \circ^{-1} & \xrightarrow{q^{-2}} & \circ^{q_{kk}} \\ & \searrow & & \nearrow & \\ & q^3 & & q_{kk}^2 q^{-6} & \\ & & \circ^{-q_{kk}} & & \end{array} .$$

Suppose that $\mathcal{B}(W)$ is finite-dimensional. Then $Q_{33}Q_{32}Q_{23} = 1$, so $q_{kk} = q^2$ and then $Q_{34}Q_{43} = q^{-2} \neq 1$. In consequence such diagram is of type super $F(4)$. Then $1 = Q_{14}Q_{41} = q^3$, which is a contradiction.

For the second case, $Q_{12}Q_{21} = q_{ii} \in \mathbb{G}_3$, and $Q_{14}Q_{41} = -1$, so the diagram contains a 4-cycle and then $\mathcal{B}(W)$ is infinite-dimensional. An analogous situation holds for the third case, because $Q_{12}Q_{21} = -q_{ii} \in \mathbb{G}_6$, and $Q_{14}Q_{41} = -1$.

For the fourth case, $\widetilde{Q}_{23} = \widetilde{Q}_{24} = \widetilde{Q}_{34} = \widetilde{q}_{kj} \neq 1$ and $\widetilde{Q}_{14} = \widetilde{Q}_{12} = \widetilde{q}_{ij} \neq 1$, so $\mathcal{B}(W)$ is infinite-dimensional.

For the last case, $Q_{44} = 1$, and then $\mathcal{B}(W)$ is infinite-dimensional.

Therefore S is infinite-dimensional in all the cases.

(ii) We use the same notation, but in this case $y_4 = [[x_{ij}, [x_{ij}, x_{ijk}]_c], x_j]_c$. So we have the following diagram for (Q_{rs}) :

$$\begin{array}{ccccc} \circ^{-1} & \xrightarrow{q} & \circ^{-1} & \xrightarrow{q^{-3}} & \circ^{q_{kk}} \\ & \searrow & & \nearrow & \\ & q^4 & & q_{kk}^2 q^{-6} & \\ & & \circ^{-q_{kk}} & & \end{array} ,$$

where $q = \widetilde{q}_{ij}$. Suppose that $\mathcal{B}(W)$ is finite-dimensional. By [H3, Table 3], this diagram cannot be connected. In consequence, $1 = Q_{41}Q_{14} = Q_{34}Q_{43}$, so $q_{kk} = \pm 1$. But then $Q_{33} = 1$, or $Q_{44} = 1$, which is a contradiction to the fact that $\mathcal{B}(W)$ is finite-dimensional. So S is infinite-dimensional. \square

Lemma 4.6. *Let $i, j, k, l \in \{1, \dots, \theta\}$ be such that $q_{jj}\widetilde{q}_{ij} = q_{jj}\widetilde{q}_{jk} = 1, \widetilde{q}_{jk}^2 = \widetilde{q}_{kl}^{-1} = qu, q_{kk} = -1, \widetilde{q}_{ik} = \widetilde{q}_{il} = \widetilde{q}_{jl}$. If $[[[x_{ijkl}, x_k]_c, x_j]_c, x_k]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

Proof. We use a similar notation and consider the corresponding subspace W generated by the corresponding primitive elements. Suppose that $\mathcal{B}(W)$ is finite-dimensional. Its associated Dynkin diagram is

$$\begin{array}{ccccc} \circ^{q_{ii}} & \xrightarrow{q^{-1}} & \circ^q & \xrightarrow{q^{-1}} & \circ^{-1} & \xrightarrow{q^2} & \circ^{q^{-2}} \\ & \searrow & & \nearrow & & & \\ & q_{ii}^2 q^{-2} & & q^2 & & & \\ & & \circ^{-q_{ii}} & & & & \end{array} , \quad q = q_{jj} .$$

Note that $1 = Q_{15}Q_{51}$, because there are no 5-cycles and $q^2 \neq 1$. Therefore $q_{ii} = q$, but this diagram is not in Heckenberger's list, so $\mathcal{B}(W)$ is infinite-dimensional, and S too. \square

Lemma 4.7. *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{jj} = \widetilde{q}_{ij}^{-1} = \widetilde{q}_{jk}$.*

- (i) *If $q_{jj} \in \mathbb{G}_3$ and $[[x_{ijk}, x_j]_c x_j]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*
(ii) *If $q_{jj} \in \mathbb{G}_4$ and $[[[x_{ijk}, x_j]_c, x_j]_c, x_j]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

Proof. (i) Using the same notation as in previous Lemmata, we have the diagram

$$\begin{array}{ccccc} \circ^{q_{ii}} & \xrightarrow{\zeta^2} & \circ^{\zeta} & \xrightarrow{\zeta} & \circ^{q_{kk}} \\ & \searrow^{q_{ii}^2} & & \swarrow_{q_{kk}^2} & \\ & & \circ^{q_{kk}q_{ii}} & & \end{array}, \quad \zeta = q_{jj} \in \mathbb{G}_3,$$

for $y_1 = x_i, y_2 = x_j, y_3 = x_k, y_4 = [[x_{ijk}, x_j]_c x_j]_c$, with corresponding matrix (Q_{rs}) , and W is the subspace generated by these elements. Note that

- if $q_{ii} = q_{kk} = -1$, then $Q_{44} = 1$;
- if $q_{ii}, q_{kk} \neq -1$, then the diagram contains a 4-cycle;
- if $q_{ii} = -1, q_{kk} \neq -1$, or $q_{ii} \neq -1, q_{kk} = -1$, the diagram contains $\circ^q \xrightarrow{q^2} \circ^{-q}$ as a subdiagram (where $q = q_{ii}$ or $q = q_{kk}$), and this connected subdiagram of rank two is not in [H3, Table 1].

In all the cases $\mathcal{B}(W)$ is infinite-dimensional, so S too.

(ii) The proof is analogous. □

Lemma 4.8. (i) *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = q_{kk} = \widetilde{q}_{ij} = -1, q_{jj} = \widetilde{q}_{jk}^{-1}, \widetilde{q}_{ik} = 1$. If $[x_{ij}, x_{ijk}]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

(ii) *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{ii} = q_{kk} = -1, \widetilde{q}_{ij} \in \mathbb{G}_3, q_{jj} = \pm \widetilde{q}_{ij} = -\widetilde{q}_{jk}, \widetilde{q}_{ik} = 1$. If $[x_i, x_{jjk}]_c - (1 + q_{jj}^2)q_{kj}^{-1} [x_{ijk}, x_j]_c - (1 + q_{jj})q_{ij} x_j x_{ijk} \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

(iii) *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $\widetilde{q}_{jk} = 1, q_{ii} = \widetilde{q}_{ij} = -\widetilde{q}_{ik} \in \mathbb{G}_3$. If $[x_i, [x_{ij}, x_{ik}]_c]_c + q_{jk}q_{ik}q_{ji} [x_{iik}, x_{ij}]_c + q_{ij} x_{ij} x_{iik} \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

(iv) *Let $i, j, k \in \{1, \dots, \theta\}$ be such that $q_{jj} = q_{kk} = \widetilde{q}_{jk} = -1, q_{ii} = -\widetilde{q}_{ij} \in \mathbb{G}_3, \widetilde{q}_{ik} = 1$. If $[x_{iijk}, x_{ijk}]_c \in \mathcal{P}(S) \setminus \{0\}$, then S is infinite-dimensional.*

Proof. We consider the same notation as before. We consider the subspace W generated by $y_1 = x_i, y_2 = x_j, y_3 = x_k$ and y_4 (the primitive element corresponding to the relation), where $y_i \in W_{h_i}^{\eta_i}$, for some $h_i \in \Gamma, \eta_i \in \widehat{\Gamma}$, and set $(Q_{rs} = \eta_s(h_r))$. We will prove again that $\mathcal{B}(W)$ is infinite-dimensional.

(i) The corresponding diagram of (Q_{rs}) is

$$\begin{array}{ccccc} \circ^{-1} & \xrightarrow{-1} & \circ^q & \xrightarrow{q^{-1}} & \circ^{-1} \\ & & \downarrow^{q^3} & \swarrow_{q^{-2}} & \\ & & \circ^{-q^2} & & \end{array}, \quad q = q_{jj} \in \mathbb{G}_3 \cup \mathbb{G}_4 \cup \mathbb{G}_6.$$

If $q \in \mathbb{G}_4$ then $Q_{44} = 1$, so $\mathcal{B}(W)$ is infinite-dimensional. If $q \in \mathbb{G}_3 \cup \mathbb{G}_6$, the diagram is not in [H3, Table 3], so $\mathcal{B}(W)$ is also again infinite-dimensional.

(ii) We note that $Q_{14}Q_{41} = Q_{34}Q_{43} = q_{jj}^2 \in \mathbb{G}_3$. Therefore, the diagram corresponding to (Q_{rs}) contains a 4-cycle, and then $\mathcal{B}(W)$ is infinite-dimensional.

(iii) In this case, $Q_{24}Q_{42} = -q_{ii}^2$ and $Q_{34}Q_{43} = -q_{kk}^2 \neq 1$, because $\widetilde{q}_{kk} \in \{-1, \widetilde{q}_{ik}^{-1}\}$. The diagram corresponding to $(Q_{rs})_{r,s=2,3,4}$ is a 3-cycle such that $\widetilde{Q}_{24}Q_{34}\widetilde{Q}_{23} \neq 1$. [H3, Lemma 9] implies that $\mathcal{B}(W)$ is infinite-dimensional.

(iv) $\mathcal{B}(W)$ is infinite-dimensional because $Q_{44} = 1$. \square

Lemma 4.9. *Let $i, j \in \{1, \dots, \theta\}$ be such that they satisfy one of the following conditions:*

(i) $-q_{ii}, -q_{jj}, q_{ii}\widetilde{q_{ij}}, q_{jj}\widetilde{q_{ij}} \neq 1$,

$$[x_i, [x_{ij}, x_j]_c]_c - \frac{(1+q_{jj})(1-q_{jj}\widetilde{q_{ij}})}{(1-\widetilde{q_{ij}})q_{jj}q_{ji}} x_{ij}^2 \in \mathcal{P}(S) \setminus \{0\};$$

(ii) $q_{jj} = -1$, $q_{ii}\widetilde{q_{ij}} \notin \mathbb{G}_6$, and also $m_{ij} \in \{4, 5\}$, or $m_{ij} = 3$, $q_{ii} \in \mathbb{G}_4$,

$$[x_i, x_{3\alpha_i+2\alpha_j}]_c - \frac{1 - q_{ii}\widetilde{q_{ij}} - q_{ii}^2\widetilde{q_{ij}}^2 q_{jj}}{(1 - q_{ii}\widetilde{q_{ij}})q_{ji}} x_{ij}^2 \in \mathcal{P}(S) \setminus \{0\};$$

(iii) $4\alpha_i + 3\alpha_j \notin \Delta_+^X$, $q_{jj} = -1$ or $m_{ji} = 2$, and also $m_{ij} \geq 3$ or $m_{ij} = 2$, $q_{ii} \in \mathbb{G}_3$,

$$[x_{3\alpha_i+2\alpha_j}, x_{ij}]_c \in \mathcal{P}(S) \setminus \{0\};$$

(iv) $3\alpha_i + 2\alpha_j \in \Delta_+^X$, $5\alpha_i + 3\alpha_j \notin \Delta_+^X$, and $q_{ii}^3\widetilde{q_{ij}}, q_{ii}^4\widetilde{q_{ij}} \neq 1$, $[x_{ii}, x_{3\alpha_i+2\alpha_j}]_c \in \mathcal{P}(S) \setminus \{0\}$;

(v) $4\alpha_i + 3\alpha_j \in \Delta_+^X$, $5\alpha_i + 4\alpha_j \notin \Delta_+^X$, $[x_{4\alpha_i+3\alpha_j}, x_{ij}]_c \in \mathcal{P}(S) \setminus \{0\}$;

(vi) $5\alpha_i + 2\alpha_j \in \Delta_+^X$, $7\alpha_i + 3\alpha_j \notin \Delta_+^X$, $[[x_{iii}, x_{ii}], x_{ii}]_c \in \mathcal{P}(S) \setminus \{0\}$;

(vii) $q_{jj} = -1$, $5\alpha_i + 4\alpha_j \in \Delta_+^X$, $[x_{ii}, x_{4\alpha_i+3\alpha_j}]_c - ax_{3\alpha_i+2\alpha_j}^2 \in \mathcal{P}(S) \setminus \{0\}$, for some $a \in \mathbf{k}^\times$.

Then S is infinite-dimensional.

Proof. Firstly we note that there exists just one connected generalized Dynkin diagram of rank three such that $3\alpha_i + 2\alpha_j \in \Delta_+^X$, for some pair i, j , which is exactly the unique one such that $m_{kl} \geq 3$ for some pair k, l . Moreover, $4\alpha_i + 3\alpha_j$, $5\alpha_i + 4\alpha_j$, are not positive roots for any pair i, j and any connected Dynkin diagram of rank 3.

We consider as above the subspace W generated by $y_1 = x_i$, $y_2 = x_j$ and y_3 , the relation which is a primitive element by hypothesis, and analyze its generalized Dynkin diagram.

(i) If $Q_{13}Q_{31} \neq 1$ or $Q_{23}Q_{32} \neq 1$, then $\mathcal{B}(W)$ is infinite-dimensional. In other case,

$$Q_{13}Q_{31} = q_{ii}^4 q_{ij}^2 q_{ji}^2 = 1, \quad Q_{23}Q_{32} = q_{jj}^4 q_{ij}^2 q_{ji}^2 = 1,$$

so $Q_{33} = q_{ii}^4 q_{ij}^4 q_{ji}^4 = 1$, and $\mathcal{B}(W)$ is also infinite-dimensional.

(ii) If $q_{ii} \in \mathbb{G}_4$, $\widetilde{q_{ij}} = q_{ii}^{-1}$ (and then (q_{rs}) is Cartan of type G_2), then

$$Q_{33} = q_{ii}^{16} q_{ij}^8 q_{ji}^8 q_{jj}^4 = 1,$$

so $\mathcal{B}(W)$ is infinite-dimensional. In other case, $Q_{13}Q_{31} \neq 1$, or $Q_{23}Q_{32} \neq 1$, or

$$Q_{13}Q_{31} = q_{ii}^8 q_{ij}^2 q_{ji}^2 = 1, \quad Q_{23}Q_{32} = q_{ij}^4 q_{ji}^4 q_{jj}^4, \quad \text{so } Q_{33} = 1,$$

and therefore $\mathcal{B}(W)$ is infinite-dimensional.

(iii) Now we calculate

$$Q_{33} = q_{ii}^{16} q_{ij}^{12} q_{ji}^{12} q_{jj}^9, \quad Q_{13}Q_{31} = q_{ii}^8 q_{ij}^3 q_{ji}^3, \quad Q_{23}Q_{32} = q_{ij}^4 q_{ji}^4 q_{jj}^6.$$

If (q_{rs}) is Cartan of type G_2 and $q_{ii} \in \mathbb{G}_6$, $\widetilde{q_{ij}} = q_{jj} = -1$, then $\mathcal{B}(W)$ is infinite-dimensional, because we have a connected diagram of rank three such that $M_{12} = 3$, and it is not of type super $G(3)$. In other case, we will prove that $Q_{13}Q_{31} \neq 1$ or $Q_{23}Q_{32} \neq 1$ to conclude that $\mathcal{B}(W)$ is infinite-dimensional. If $m_{ji} \geq 2$, we have the following possible cases:

- $q_{ii} = -\zeta$, $\widetilde{q_{ij}} = \zeta^7$, $q_{jj} = \zeta^3$, $\zeta \in \mathbb{G}_9$; in such case, $Q_{23}Q_{32} = \zeta$.
- $q_{ii} = -\zeta$, $\widetilde{q_{ij}} = -\zeta^{12}$, $q_{jj} = \zeta^5$, $\zeta \in \mathbb{G}_{15}$; therefore, $Q_{23}Q_{32} = \zeta^3$.

Also, if $q_{ii} = \zeta^8$, $\widetilde{q}_{ij} = \zeta^3$, $q_{jj} = -1$, $\zeta \in \mathbb{G}_{12}$, then $Q_{13}Q_{31} = \zeta$. In all the remaining cases, $q_{jj} = -1$ and $\widetilde{q}_{ij} \notin \mathbb{G}_4$, so $Q_{23}Q_{32} \neq 1$.

(iv) This relation is not redundant just in the following two cases:

$$\circ\zeta^3 \xrightarrow{\zeta^8} \circ^{-1}, \quad \zeta \in \mathbb{G}_9, \quad \circ\eta^3 \xrightarrow{-\eta^4} \circ^{-\eta^4}, \quad \eta \in \mathbb{G}_{15}.$$

Note that they are not contained in any connected diagram of rank three in [H3, Table 2], so it is enough to verify that $Q_{13}Q_{31} \neq 1$ or $Q_{23}Q_{32} \neq 1$ to conclude that $\mathcal{B}(W)$ is infinite-dimensional. For the first diagram, $Q_{23}Q_{32} = \zeta^4 \neq 1$; and for the second one, $Q_{23}Q_{32} = -\eta^{-4} \neq 1$.

(v) The proof is analogous to (iii). Note that

$$Q_{33} = q_{ii}^{25} q_{ij}^{12} q_{ji}^{20} q_{jj}^{16}, \quad Q_{13}Q_{31} = q_{ii}^{10} q_{ij}^4 q_{ji}^4, \quad Q_{23}Q_{32} = q_{ij}^5 q_{ji}^5 q_{jj}^8.$$

We have that $q_{jj} = -1$ for every diagram satisfying the conditions for this item. Also, if $q_{ii} = \zeta \in \mathbb{G}_5$, $\widetilde{q}_{ij} = \zeta^2$, it follows that $Q_{33} = 1$. In the remaining cases, $\widetilde{q}_{ij} \notin \mathbb{G}_5$, so $Q_{32}Q_{23} \neq -1$, and then $\mathcal{B}(W)$ is infinite-dimensional.

(vi) In this case,

$$Q_{33} = q_{ii}^{49} q_{ij}^{21} q_{ji}^{21} q_{jj}^9, \quad Q_{13}Q_{31} = q_{ii}^{14} q_{ij}^3 q_{ji}^3, \quad Q_{23}Q_{32} = q_{ij}^7 q_{ji}^7 q_{jj}^6.$$

We have that $q_{jj} = -1$ for every diagram satisfying the conditions for this item, and also $\widetilde{q}_{ij} \notin \mathbb{G}_7$, so $Q_{23}Q_{32} \neq 1$. Therefore $\mathcal{B}(W)$ is infinite-dimensional.

(vii) The proof is analogous to the one for (i).

We conclude that S is infinite-dimensional in all the cases. \square

Now we can prove the main Theorem of this Section.

Theorem 4.10. *Let $S = \bigoplus_{n \geq 0} S(n)$ be a finite-dimensional graded Hopf algebra in ${}^G_G\mathcal{YD}$, where Γ is a finite abelian group, such that $S(0) = \mathbf{k}1$. Fix a basis x_1, \dots, x_θ of $V := S(1)$, such that $x_i \in S(1)_{g_i}^{\chi_i}$ for some $g_i \in \Gamma$ and $\chi_i \in \widehat{\Gamma}$, and set $q_{ij} := \chi_j(g_i)$. If S is generated as an algebra by $S(0) \oplus S(1)$, then $S \cong \mathcal{B}(V)$.*

Proof. As S is generated as an algebra by $S(0) \oplus S(1)$, the canonical projection $T(V) \rightarrow \mathcal{B}(V) = T(V)/I(V)$ induces a surjective morphism $\pi : S \rightarrow \mathcal{B}(V)$ of graded braided Hopf algebras; we can consider $S = T(V)/I$, for some graded braided Hopf ideal I of $T(V)$, generated by homogeneous elements of degree ≥ 2 , $I \subseteq I(V)$.

Suppose that $I(V) \not\supseteq I$. Then at least one of the generators of $I(V)$ from Theorem 3.1 does not belong to I . We can assume that $\mathbf{x} \in I(V) \setminus I$ is one of these generators, of minimal degree k . Then \mathbf{x} is primitive in S by Lemma 3.2.

By Proposition 4.1 and Lemmata 4.2-4.9, we deduce that $\mathbf{x} = x_\alpha^{N_\alpha}$ for some $\alpha \in \mathcal{O}$, or a simple root $\alpha = \alpha_i$ such that i is not a Cartan vertex, or $\alpha = \alpha_i + \alpha_j$, such that $N_\alpha = 2$, $q_{ii} = q_{jj} = \widetilde{q}_{ij} = -1$. If $g_\alpha \in \Gamma$, $\chi_\alpha \in \widehat{\Gamma}$ are the associated elements, we have that $q_\alpha = \chi_\alpha(g_\alpha)$, which is a root of unity of order N_α . Therefore $g_\alpha^{N_\alpha} \in \Gamma$ and $\chi_\alpha^{N_\alpha} \in \widehat{\Gamma}$ are the associated elements to \mathbf{x} , and

$$c(\mathbf{x} \otimes \mathbf{x}) = g_\alpha^{N_\alpha} \cdot \mathbf{x} \otimes \mathbf{x} = \chi_\alpha^{N_\alpha} (g_\alpha^{N_\alpha}) \mathbf{x} \otimes \mathbf{x} = \mathbf{x} \otimes \mathbf{x},$$

so \mathbf{x} generates in S an infinite-dimensional braided Hopf subalgebra, and we obtain a contradiction. In consequence, $S \cong \mathcal{B}(V)$. \square

Remark 4.11. Note that we just use the fact that the braiding is diagonal, so we can generalize this Theorem to a general braided Hopf algebra R in ${}^H_H\mathcal{YD}$, where H is a finite-dimensional Hopf algebra which acts diagonally over $R(1)$.

The following Theorem answers positively Conjecture 1 in the case that the group of group-like elements is abelian. It extends [AS4, Thm. 5.5].

Theorem 4.12. *Let H be a finite-dimensional pointed Hopf algebra over an abelian group Γ . Then H is generated by its group-like and skew-primitive elements.*

Proof. Let $\text{gr } H = R\#\mathbf{k}\Gamma$, $V = R(1)$. Then H is generated by its group-like and skew-primitive elements if and only if $\text{gr } H$ satisfies this condition, which is equivalent to the fact that R is the Nichols algebra $\mathcal{B}(V)$. Let S be the graded dual R^* in the category ${}^G_C\mathcal{YD}$, which is generated as an algebra by $S(1) = V^*$. By [AS3, Lemma 2.3] it is enough to prove that S is the Nichols algebra $\mathcal{B}(V^*)$, which follows by Theorem 4.10. \square

REFERENCES

- [An] N. Andruskiewitsch, *About finite dimensional Hopf algebras*, Notes of a course given at the CIMPA School Quantum symmetries in theoretical physics and mathematics Bariloche 2000. *Contemp. Math.* **294**, 1–57 (2002).
- [AA] N. Andruskiewitsch and I. Angiono, *On Nichols algebras with generic braiding*, *Modules and Comodules*. Trends in Mathematics. Brzezinski, T.; Gómez Pardo, J.L.; Shestakov, I.; Smith, P.F. (Eds.), pp. 47-64 (2008). ISBN: 978-3-7643-8741-9.
- [AA Y] N. Andruskiewitsch, I. Angiono and H. Yamane, *On pointed Hopf superalgebras*. Accepted in *Contemp. Math.* arxiv:1009.5148, 18 pp.
- [AS1] N. Andruskiewitsch and H.-J. Schneider, *Lifting of quantum linear spaces and pointed Hopf algebras of order p^3* , *J. Algebra* **209**, 658-691 (1998).
- [AS2] N. Andruskiewitsch and H.-J. Schneider, *Finite quantum groups and Cartan matrices*, *Adv. Math.* **154**, 1–45 (2000).
- [AS3] N. Andruskiewitsch and H.-J. Schneider, *Pointed Hopf algebras*, “New directions in Hopf algebras”, MSRI series Cambridge Univ. Press; 1–68 (2002).
- [AS4] N. Andruskiewitsch and H.-J. Schneider, *On the classification of finite-dimensional pointed Hopf algebras*, *Annals of Mathematics* Vol. **171** (2010), No. 1, 375417.
- [A1] I. Angiono, *On Nichols algebras with standard braiding*, *Algebra and Number Theory* Vol. 3, No. 1, 35-106, (2009).
- [A2] I. Angiono, *A presentation by generators and relations of Nichols algebras of diagonal type and convex orders on root systems*, submitted. arXiv:1008.4144.
- [AGI] I. Angiono and A. García Iglesias. *Pointed Hopf algebras with standard braiding are generated in degree one*, *Contemp. Math.* **537** (2011), 57–70.
- [CH1] M. Cuntz and I. Heckenberger, *Weyl groupoids with at most three objects*. *J. Pure Appl. Algebra* **213**, No. 6 (2009), 1112–1128.
- [CH2] M. Cuntz and I. Heckenberger, *Finite Weyl groupoids of rank three*, math. QA/0912.0212.
- [GHV] M. Graña, I. Heckenberger and L. Vendramín, *SARNA: Arithmetic Root Systems and Nichols Algebras Software*, a program to calculate the Arithmetic Root System associated to a braid of diagonal type. Available at <http://mate.dm.uba.ar/~lvendram/>
- [H1] I. Heckenberger, *Weyl equivalence for rank 2 Nichols algebras of diagonal type*. *Ann. Univ. Ferrara - Sez. VII - Sc. Mat.*, 281–289 (2005).
- [H2] I. Heckenberger, *The Weyl groupoid of a Nichols algebra of diagonal type*, *Invent. Math.* **164**, 175–188 (2006).
- [H3] I. Heckenberger, *Classification of arithmetic root systems*, *Adv. Math.* **220** (2009) 59-124.
- [H4] I. Heckenberger, *Lusztig isomorphisms for Drinfel’d doubles of bosonizations of Nichols algebras of diagonal type*, *J. Alg.* **323** (2010), 2130–2180.
- [HS] I. Heckenberger and H.-J. Schneider, *Right coideal subalgebras of Nichols algebras and the Duflo order on the Weyl groupoid*. math.QA/0909.0293.
- [HY] I. Heckenberger and H. Yamane, *A generalization of Coxeter groups, root systems, and Matsumoto’s theorem*, *Math. Z.* **259** (2008), 255–276.
- [He] M. Helbig, *On the lifting of Nichols algebras*, math. QA/1003.5882.
- [Kh] V. Kharchenko, *A quantum analog of the Poincare-Birkhoff-Witt theorem*, *Algebra and Logic* **38**, (1999), 259-276.
- [L] G. Lusztig, *Introduction to quantum groups*, Birkhäuser (1993).
- [Mo] S. Montgomery, *Hopf algebras and their action on rings*, CBMS Regional Conference Series **82** (1993).

- [R] M. Rosso, *Lyndon words and Universal R-matrices*, talk at MSRI, October 26, 1999, available at <http://www.msri.org>; *Lyndon basis and the multiplicative formula for R-matrices*, preprint (2003).
- [Se] V. Serganova, *On generalizations of root systems*. Commun. Algebra **24** (1996), 4281–4299.
- [U] S. Ufer, *PBW bases for a class of braided Hopf algebras*, J. Alg. **280** (2004) 84–119.
- [Y] H. Yamane, *Quantized enveloping algebras associated to simple Lie superalgebras and their universal R-matrices* Publ. Res. Inst. Math. Sci. **30** (1994), 15–87.

FAMAF-CIEM (CONICET), UNIVERSIDAD NACIONAL DE CÓRDOBA, MEDINA ALLENDE S/N, CIUDAD UNIVERSITARIA, 5000 CÓRDOBA, REPÚBLICA ARGENTINA.

E-mail address: `angiono@famaf.unc.edu.ar`