

UNIVERSIDAD NACIONAL DE CÓRDOBA  
FACULTAD DE MATEMÁTICA, ASTRONOMÍA Y FÍSICA

TECHNICAL REPORT

ASYMPTOTIC BEHAVIOR OF RA-ESTIMATES IN AUTOREGRESSIVE 2D  
PROCESSES

BY

BUSTOS, O., RUIZ, M., OJEDA, S.,  
VALLEJOS, R. AND FRERY, A.

FEBRUARY 2007

# ASYMPTOTIC BEHAVIOR OF RA-ESTIMATES IN AUTOREGRESSIVE 2D PROCESSES

BUSTOS, O.<sup>1</sup>, RUIZ, M.<sup>2</sup>, OJEDA, S.<sup>3</sup>, VALLEJOS, R.<sup>4</sup>, AND FRERY, A.<sup>5</sup>

## Abstract

In this work we study the asymptotic behavior of a robust class of estimator of coefficients of AR-2D process. We established the consistency and asymptotic normality of the RA estimator under precise conditions. This class of models has diverse applications in image modeling and statistical image processing.

KEYWORDS AND PHRASES: Image Processing, Bidimensional Processes, AR-2D Models, Polynomial Coefficients, Robust Estimators, Residual Autocovariance, Asymptotic Normality and Consistency.

---

<sup>1,3</sup> FaMAF-CIEM, UNC, Argentina.

<sup>2</sup> Ciencias Exactas, UNRC, Argentina.

<sup>4</sup> IMAFI, Universidad de Valparaíso, Chile.

<sup>5</sup> Instituto de Computação, UFAL, Brasil.

# 1 INTRODUCTION

Robust inference techniques appear in a diversity of contexts and applications, though the terms “robust” and “robustness” are quite freely used in the image processing and computer vision literature, not necessarily with the usual statistical meaning.

The median and similar order-based filters are basic tools in image processing (Aysal and Barner (2006); Huang and Lee (2006); Palenichka et al. (2000), (1998)), and in some cases particular attention has been devoted to obtain the distribution of those estimators (Steland (2005)). Frery et al. (1997) derived a family of robust estimators for a class of low signal-to-noise ratio images, while Vallejos and Mardesic (2004) proposed the robust estimation of structural parameters for their restoration.

Other resistant approaches have proved being successful in image restoration (see, for instance, Ben Hamza and Krim (2001); Chu et al. (1998); Koivunen (1995); Marroquin et al. (1998); Rabie (2005); Tarel et al. (2002); Voloshynovskiy et al. (2000); Zervakis and Kwon (1992)). A common challenge in these applications is that the number of observations is reduced to a few, typically less than a hundred points.

When it comes to image analysis, many robust techniques have been proposed. In this case, the sample size is usually larger than the one available in filters and, frequently, structure and topology do not impose heavy requirements or constraints. In some cases, strong hypothesis are made on the laws governing the observed process (Allende and Pizarro (2003); Brunelli and Messelodi (1995); Bustos et al. (2002); Butler (1998); Dryden et al. (2002); Van de Weijer and Van den Boomgaard (2005)); other approaches can be seen in the works by Bouzouba and Radouane (2000); Brandle et al. (2003); Nirel et al. (1998); Sim et al. (2004); Tohka et al. (2004); Xu (2005) and Zervakis et al. (1995).

High-level image analysis, or vision, also benefits from the use of robust estimation techniques, as can be seen in Black and Rangarajan (1996), Black et al. (1997), Chen et al. (2003), Comport et al. (2006), Glendinning (1999), Gottardo et al. (2006), Hasler et al. (2003), Kim and Han (2006), Li et al. (1998), Meer et al. (1991), Mirza and Boyer (1993), Prastawa et al. (2004), Roth (2006), Singh et al. (2004), Stewart (1999), Torr and Zisserman (2000) and Wang and Suter (2004a,b).

In a wide variety of situations such as image analysis, remote sensing and agricultural field trials, observations are obtained on two-dimensional lattices or grids. A class of two-dimensional autoregressive processes has been suggested (Whittle (1954)) as a source of models for the spatial correlation in such data (Tjøstheim (1978)). These models are natural extensions of the autoregressive processes used in time series analysis (Basu and Reinsel (1993)).

Most robust techniques developed for parametric models in time series have been implemented for spatial parametric models when the process has been contaminated with innovation or additive outliers (Kashyap and Eom (1988)). Since a single outlier can produce bias and a large variance in the estimators, most of those proposals aim at providing estimators that are more resistant to the presence of contamination.

There are at least three classes of robust estimators that have been studied in this context, namely, the M, GM and RA estimators. Kashyap and Eom (1988) introduced M estimators for two-dimensional autoregressive models, and a recursive image restoration algorithm was implemented using a robust M estimators. Allende et al. (1998) studied the implementation of Generalized M (GM) estimators for the same class of models. The image restoration algorithm previously developed by Kashyap and Eom (1988) was generalized by Allende et al. (2001).

Robust Residual Autocovariance (RA) estimators were introduced by Bustos and Yohai (1986) in the context of time series. They are based on “cleaning” the residuals by the application of a robustifying  $\psi$  function in the recursive estimation procedure. Ojeda (1999) studied the extension of the RA estimators for spatial unilateral autoregressive models. The computational implications of that extension have been studied by Ojeda et al. (2002) and Vallejos et al. (2006). Monte Carlo simulation studies show that the performance of the RA estimators is better than the M estimators and slightly better than the GM estimators when the model has been contaminated with additive outliers.

Although the performance of the M and GM estimators is reasonable under innovation outliers, the asymptotic properties are still open problems.

In this paper we study the asymptotic behavior of the RA estimators for unilateral autoregressive spatial processes, generalizing the results for one-dimensional time series asymptotic behavior established by Bustos et al. (1984). We give precise conditions for the consistency and asymptotic normality of the RA estimators.

The paper is organized as follows. In Section 2, the model and some preliminary notation are established. In Section 3 the RA estimators of a two-dimensional autoregressive process is introduced. Section 4 establishes the Strong Consistency of the RA estimators, while Section 5 proves its Asymptotic Normality. The proofs of the results are organized in two subsections of the Appendix (Section 6). The paper concludes in Section 7 with some final remarks and directions for future work.

## 2 AR-2D MODELS

Throughout this paper we assume that the random variables are defined on the same probability space  $(\Omega, \mathbb{F}, P)$ .

If  $\underline{m} = (m_1, m_2)'$  and  $\underline{k} = (k_1, k_2)'$   $\in \mathbb{Z}^2$ , we write  $\underline{m} \leq \underline{k}$  if  $m_i \leq k_i$  for  $i = 1, 2$ . Let

$$I_0 = \{\underline{m} \in \mathbb{Z}^2 : \underline{0} \leq \underline{m} \text{ and } \underline{m} \neq \underline{0} = (0, 0)'\},$$

and let

$$T = \{\underline{t}_1, \dots, \underline{t}_L\} = \{(t_{1,1}, t_{1,2})', \dots, (t_{L,1}, t_{L,2})'\} \subset I_0,$$

be a finite and not empty set.

Consider  $\phi_0 = (\phi_{01}, \dots, \phi_{0L})' \in \mathbb{R}^L$  such that

$$\sup_{1 \leq p \leq L} \{|\phi_{0p}|\} < \frac{1}{L}, \quad (1)$$

and let  $P_{\phi_0}$  be a polynomial defined over  $\mathbb{C}^2$  as follows

$$P_{\phi_0}(z, w) = 1 - \sum_{p=1}^L \phi_{0p} z^{t_{p,1}} w^{t_{p,2}}. \quad (2)$$

For each  $\underline{m} \in \mathbb{Z}^2$  let  $X_{\underline{m}} : \Omega \rightarrow \mathbb{R}$  be a random variable such that  $\tilde{\mathbf{X}} = (X_{\underline{m}})_{\underline{m} \in \mathbb{Z}^2}$  is a  $AR(P_{\phi_0}, \tilde{\varepsilon})$  process; that is, for each  $\omega \in \Omega$  and  $\underline{m} \in \mathbb{Z}^2$ ,  $X_{\underline{m}}$  satisfies

$$X_{\underline{m}}(\omega) = \mu + \sum_{p=1}^L \phi_{0p} X_{\underline{m}-\underline{t}_p}(\omega) + \varepsilon_{\underline{m}}(\omega), \quad (3)$$

where  $\tilde{\varepsilon} = (\varepsilon_{\underline{m}})_{\underline{m} \in \mathbb{Z}^2}$  is a white noise, i.e., the components of  $\tilde{\varepsilon}$  are independent and identically distributed random variables with common distribution function  $F_{\varepsilon}$  (not necessarily Gaussian), zero mean and finite variance  $\sigma_{\varepsilon}^2 > 0$ . Whenever possible we will drop the argument  $\omega$ .

The following condition is required:

**Assumption 1.** *The distribution function of the errors,  $F_{\varepsilon}$ , is absolutely continuous with density  $f_{\varepsilon}$ .*

Our interest is the estimation of the coefficients  $\phi_0$  from the observed values of  $\tilde{\mathbf{X}}$ , so we assume in this paper that  $\mu$  and  $\sigma_{\varepsilon}$  are known and without loss of generality we take  $\mu = 0$ . If the position  $\mu$  and the scale  $\sigma_{\varepsilon}$  are unknown they can be estimated using (possibly robust) estimators.

**Example 1.** *In practice, typical values for  $L$  are 1, 2 or at most 3. The following case has been frequently found in recent applications (see Ojeda et al. (2002)): consider  $T = \{(1, 0)', (1, 1)', (0, 1)'\}$  and set  $\underline{t}_1 = (1, 0)'$ ,  $\underline{t}_2 = (1, 1)'$  and  $\underline{t}_3 = (0, 1)'$ . So, the polynomial in (2) is given by*

$$P_{\phi_0}(z, w) = 1 - \phi_{01}z - \phi_{02}zw - \phi_{03}w.$$

Hence, the model is described by

$$X_{(m_1, m_2)'} = \phi_{01}X_{(m_1-1, m_2)'} + \phi_{02}X_{(m_1-1, m_2-1)'} + \phi_{03}X_{(m_1, m_2-1)'} + \varepsilon_{(m_1, m_2)'}$$

We will now show an important characterization of  $\tilde{\mathbf{X}}$ . Assuming that  $X_{\underline{m}} \in L^2(\Omega, \mathbb{F}, P)$  with

$$\begin{aligned} \|X_{\underline{m}}\|_{L^2}^2 &= \text{Variance of } X_{\underline{m}} = \sigma_X^2, \\ E(X_{\underline{m}}) &= 0, \end{aligned}$$

(see Guyon (1993), Chapter 1), we have that, for each  $\underline{m}$ ,  $X_{\underline{m}}$  can be expressed as a series of terms of  $(\varepsilon_{\underline{n}})_{\underline{n} \in \mathbb{Z}^2}$  in the  $L^2(\Omega, \mathbb{F}, P)$  space.

Indeed, for each  $\omega \in \Omega$  and  $\underline{m} \in \mathbb{Z}^2$ , after one iteration in (3),  $X_{\underline{m}}(\omega)$  can be written as

$$\begin{aligned} X_{\underline{m}}(\omega) &= \sum_{k_1=1}^L \phi_{0k_1} X_{\underline{m}-\underline{t}_{k_1}}(\omega) + \varepsilon_{\underline{m}}(\omega) \\ &= \sum_{k_1=1}^L \phi_{0k_1} \left[ \sum_{k_2=1}^L \phi_{0k_2} X_{\underline{m}-\underline{t}_{k_1}-\underline{t}_{k_2}}(\omega) + \varepsilon_{\underline{m}-\underline{t}_{k_1}}(\omega) \right] + \varepsilon_{\underline{m}}(\omega) \\ &= \sum_{k_1, k_2=1}^L \phi_{0k_1} \phi_{0k_2} X_{\underline{m}-\underline{t}_{k_1}-\underline{t}_{k_2}}(\omega) + \sum_{k_1=1}^L \phi_{0k_1} \varepsilon_{\underline{m}-\underline{t}_{k_1}} + \varepsilon_{\underline{m}}(\omega); \end{aligned}$$

now, if we iterate twice we have

$$\begin{aligned} X_{\underline{m}}(\omega) &= \sum_{k_1, k_2, k_3=1}^L \phi_{0k_1} \phi_{0k_2} \phi_{0k_3} X_{\underline{m}-\underline{t}_{k_1}-\underline{t}_{k_2}-\underline{t}_{k_3}}(\omega) + \\ &\quad \sum_{k_1, k_2=1}^L \phi_{0k_1} \phi_{0k_2} \varepsilon_{\underline{m}-\underline{t}_{k_1}-\underline{t}_{k_2}} + \sum_{k_1=1}^L \phi_{0k_1} \varepsilon_{\underline{m}-\underline{t}_{k_1}} + \varepsilon_{\underline{m}}(\omega). \end{aligned}$$

So, after  $h-1 \geq 2$  iterations, we arrive to the following expression:

$$X_{\underline{m}}(\omega) = \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^h} I_{h, \tilde{\mathbf{k}}}(\phi_0) X_{\underline{m}-s(h, \tilde{\mathbf{k}})}(\omega) + \sum_{j=1}^{h-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j, \tilde{\mathbf{k}}}(\phi_0) \varepsilon_{\underline{m}-s(j, \tilde{\mathbf{k}})}(\omega) + \varepsilon_{\underline{m}}(\omega),$$

where  $\mathbb{L} = \{1, 2, \dots, L\}$ , for each  $j \in \mathbb{N}$  and  $\tilde{\mathbf{k}} = (k_1, \dots, k_j)' \in \mathbb{L}^j$

$$s(j, \tilde{\mathbf{k}}) = \sum_{i=1}^j \underline{t}_{k_i},$$

and  $I_{h, \tilde{\mathbf{k}}}: \Theta \rightarrow \mathbb{R}$  is the function defined on

$$\Theta = \left\{ \phi = (\phi_1, \dots, \phi_L)' \in \mathbb{R}^L : \sup_{1 \leq p \leq L} |\phi_p| < \frac{1}{L} \right\}, \quad (4)$$

such that

$$I_{h, \tilde{\mathbf{k}}}(\phi) = \phi_{k_1} \dots \phi_{k_h}, \forall \phi \in \Theta.$$

Considering that

$$\left\| \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^h} I_{h, \tilde{\mathbf{k}}}(\phi_0) X_{\underline{m}-s(h, \tilde{\mathbf{k}})} \right\|_{L^2} \leq \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^h} |I_{h, \tilde{\mathbf{k}}}(\phi_0)| \sigma_X = \sigma_X \left( \sum_{p=1}^L |\phi_{0p}| \right)^h,$$

and that, by (1),  $\lim_{h \rightarrow \infty} \left( \sum_{p=1}^L |\phi_{0p}| \right)^h = 0$ , the following proposition is valid.

**Proposition 1.** *For all  $\underline{m} \in \mathbb{Z}^2$  holds that*

$$X_{\underline{m}} = \varepsilon_{\underline{m}} + \sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j, \tilde{\mathbf{k}}}(\phi_0) \varepsilon_{\underline{m}-s(j, \tilde{\mathbf{k}})}, \quad (5)$$

where the series converges in the  $L^2(\Omega, \mathbb{F}, P)$  space.

### 3 ESTIMATORS OF POLYNOMIAL COEFFICIENTS $\phi_0$

In order to define the estimators of  $\phi_0$  we need to introduce some general notation. Let  $\mathbf{H}: \Theta \rightarrow \mathbb{R}^L$  be a differentiable function and, for each  $1 \leq p \leq L$ , let  $H_p: \Theta \rightarrow \mathbb{R}$  be its  $p$ -th component; that is,  $H_p(\phi) = (\mathbf{H}(\phi))_p$ . Denote the

derivative of  $\mathbf{H}$  by  $\mathbb{D}\mathbf{H}$  and, for each  $1 \leq p, q \leq L$ , let  $D_q(H_p)(\phi)$  be the partial derivative with respect to the  $q$ -th component of the function  $H_p$  evaluated at  $\phi$ . Similarly, if  $h: \Theta \rightarrow \mathbb{R}$  is a differentiable function,  $D_q(h)(\phi)$  and  $\nabla h(\phi)$  denote the partial derivative with respect to the  $q$ -th component and the gradient of  $h$  evaluated at  $\phi$ , respectively.

Let  $\tilde{\mathbf{X}}$  be the set of all possible realizations of the random variables  $X_{\underline{m}}$ ,  $\underline{m} \in I_0$ . Since for all  $\underline{m} \in \mathbb{Z}^2$ ,  $E(X_{\underline{m}}) = 0$ , and for convenience (it will be clear in the notation introduced below) we assume in this section that

$$X_{\underline{m}}(\omega) = 0 \text{ if } \underline{m} \notin I_0, \text{ for all } \omega \in \Omega. \quad (6)$$

For each  $\underline{m} \in \mathbb{Z}^2$ , let  $r_{\underline{m}}: \Omega \times \Theta \rightarrow \mathbb{R}$  be a function defined as follows:

$$r_{\underline{m}}(\omega, \phi) = X_{\underline{m}}(\omega) - \sum_{p=1}^L \phi_p X_{\underline{m}-\underline{t}_p}(\omega).$$

Then, by (3) and (6), for every  $\phi \in \Theta$ , the residual with respect to  $\phi$ ,  $r_{\underline{m}}(\phi)$ , is

$$\begin{aligned} r_{\underline{m}}(\phi_0) &\equiv \varepsilon_{\underline{m}}, \text{ if } (\underline{m} - \underline{t}_p) \in I_0, \forall 1 \leq p \leq L, \\ r_{\underline{m}}(\phi) &\equiv 0, \text{ if } \underline{m} \notin I_0. \end{aligned} \quad (7)$$

For each positive integer  $M$  let us define the square window of  $M$  order as

$$W_M = \{\underline{m} \in I_0 : \underline{m} \leq (M, M)\},$$

and

$$\tilde{\mathbf{X}}_M = (X_{\underline{m}})_{\underline{m} \in W_M},$$

denotes the observed process in  $W_M$ .

**Remark 1.** We assume that  $M$  is large enough to have a well defined estimation problem; for example, consider  $M \geq M_0 = \inf\{M': T \subset W_{M'}\}$ .

Now we introduce the classical definition of least squares estimator as follows:

**Definition 1.** The least squares estimator of  $\phi_0$  based on  $\tilde{\mathbf{X}}_M$ , with domain  $\hat{\Omega}_M \subset \Omega$  is defined as the function  $\hat{\phi}_M: \hat{\Omega}_M \rightarrow \Theta$  such that

$$\sum_{\underline{m} \in W_M} \left( r_{\underline{m}}(\omega, \hat{\phi}_M(\omega)) \right)^2 \leq \sum_{\underline{m} \in W_M} \left( r_{\underline{m}}(\omega, \phi) \right)^2,$$

for all  $\omega \in \hat{\Omega}_M$  and  $\phi \in \Theta$ .

Considering that, for each  $\omega \in \hat{\Omega}_M$  and  $\underline{m} \in \mathbb{Z}^2$  the function  $\phi \mapsto r_{\underline{m}}(\omega, \phi)$  is continuously differentiable, then  $\hat{\phi}_M$  satisfies

$$\sum_{\underline{m} \in W_M} \left( r_{\underline{m}}(\omega, \hat{\phi}_M(\omega)) \right) \cdot \nabla \left( r_{\underline{m}}(\omega, \hat{\phi}_M(\omega)) \right) = \mathbf{0},$$

for all  $\omega \in \hat{\Omega}_M$ . Equivalently, by (7),

$$\sum_{\underline{m} \in W_M} \left( r_{\underline{m}}(\omega, \hat{\phi}_M(\omega)) \right) \cdot X_{\underline{m}-\underline{t}_p}(\omega) = 0, \quad (8)$$

for all  $1 \leq p \leq L$  and  $\omega \in \hat{\Omega}_M$ . But, by Proposition 1, (6) and (7),

$$X_{\underline{m}-\underline{t}_p}(\omega) = r_{\underline{m}-\underline{t}_p}(\omega, \phi_0) + \sum_{j=1}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j, \tilde{\mathbf{k}}}(\phi_0) r_{\underline{m}-\underline{t}_p-s(j, \tilde{\mathbf{k}})}(\omega, \phi_0),$$

for all  $\omega \in \hat{\Omega}_M$ . Replacing this expression in (8), the following equality holds:

$$\sum_{\underline{m} \in W_M} \left\{ r_{\underline{m}}(\omega, \hat{\phi}_M(\omega)) r_{\underline{m}-\underline{t}_p}(\omega, \phi_0) + \sum_{j=1}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j, \tilde{\mathbf{k}}}(\phi_0) r_{\underline{m}}(\omega, \hat{\phi}_M(\omega)) r_{\underline{m}-\underline{t}_p-s(j, \tilde{\mathbf{k}})}(\omega, \phi_0) \right\} = 0, \quad (9)$$

for all  $1 \leq p \leq L$  and  $\omega \in \widehat{\Omega}_M$ . Now, replacing the parameter by its estimator in (9) we have the following definition:

**Definition 2.** The least squares estimator of  $\phi_0$  based on the covariance of the residuals corresponding to the observations  $\widetilde{\mathbf{X}}_M$ , with domain  $\Omega_{MQ,M} \subset \Omega$  is defined as the function  $\widehat{\phi}_{Q,M}: \Omega_{MQ,M} \rightarrow \Theta$  such that

$$\sum_{\underline{m} \in W_M} \left( \chi_{\underline{m},M} \left( \omega, \widehat{\phi}_{Q,M}(\omega) \right) \right)_p = 0,$$

for all  $1 \leq p \leq L$  and  $\omega \in \Omega_{MQ,M}$ , where  $\chi_{\underline{n},N}: \Omega \times \Theta \rightarrow \mathbb{R}^L$  is given by

$$\begin{aligned} \left( \chi_{\underline{n},N}(\omega, \phi) \right)_p &= r_{\underline{n}}(\omega, \phi) \cdot r_{\underline{n}-\underline{t}_p}(\omega, \phi) + \\ &\sum_{j=1}^{N-1} \sum_{\mathbf{k} \in \mathbb{L}^j} I_{j,\mathbf{k}}(\phi) r_{\underline{n}}(\omega, \phi) \cdot r_{\underline{n}-\underline{t}_p-s(j,\mathbf{k})}(\omega, \phi), \end{aligned}$$

$1 \leq N$ ,  $\omega \in \Omega$ ,  $\underline{n} \in \mathbb{Z}^2$ ,  $\phi \in \Theta$  and  $1 \leq p \leq L$ .

**Remark 2.** If we assume that  $\widetilde{\mathbf{X}} = (X_{\underline{m}})_{\underline{m} \in \mathbb{Z}^2}$  is a  $AR(P_{\phi_0}, \widetilde{\epsilon})$  Gaussian process, then the asymptotic properties of this estimator can be derived from more general results as in Guyon (1993), Chapter 3.

It is a well known fact that least squares estimator based on the covariances are not robust. Hence, the idea is to make them robust using adequate continuous and bounded score functions.

**Definition 3.** The RA estimator of  $\phi_0$  based on the observations  $\widetilde{\mathbf{X}}_M$  with domain  $\Omega_{RA,M} \subset \Omega$  is defined as the function  $\widehat{\phi}_M^{RA}: \Omega_{RA,M} \rightarrow \Theta$ , such that

$$0 = \sum_{\underline{m} \in W_M} \left( \chi_{\underline{m},M}^\eta(\omega, \widehat{\phi}_M^{RA}(\omega)) \right)_p,$$

for all  $1 \leq p \leq L$  and  $\omega \in \Omega_{RA,M}$ , where  $\chi_{\underline{n},N}^\eta: \Omega \times \Theta \rightarrow \mathbb{R}^L$  is given by

$$\begin{aligned} \left( \chi_{\underline{n},N}^\eta(\omega, \phi) \right)_p &= \eta \left( r_{\underline{n}}^*(\omega, \phi), r_{\underline{n}-\underline{t}_p}^*(\omega, \phi) \right) + \\ &\sum_{j=1}^{N-1} \sum_{\mathbf{k} \in \mathbb{L}^j} I_{j,\mathbf{k}}(\phi) \eta \left( r_{\underline{n}}^*(\omega, \phi), r_{\underline{n}-\underline{t}_p-s(j,\mathbf{k})}^*(\omega, \phi) \right), \end{aligned}$$

and

$$r_{\underline{n}}^*(\omega, \phi) = r_{\underline{n}}(\omega, \phi) / \sigma_\varepsilon,$$

$1 \leq N$ ,  $\omega \in \Omega$ ,  $\underline{n} \in \mathbb{Z}^2$ ,  $\phi \in \Theta$ ,  $1 \leq p \leq L$  and  $\eta: \mathbb{R}^2 \rightarrow \mathbb{R}$  is a score function.

Score functions can be chosen from several families or types (see Bustos and Yohai (1986)). The Mallows type score functions are defined as  $\eta_M(u, v) = \psi_1(u)\psi_2(v)$ , while the Hampel type is  $\eta_H(u, v) = \psi(uv)$ , where the  $\psi$  functions are continuous and odd, and they may be chosen, for example, from the Huber family:

$$\psi_{H,k}(u) = \text{sgn}(u) \cdot \min(|u|, k),$$

where  $\text{sgn}(u)$  is the sign function and  $k$  is a constant; or in a redescending family, for example the bisquare family defined by

$$\psi_{B,k}(u) = k \cdot \psi_B \left( \frac{u}{k} \right),$$

where  $k$  is a constant and

$$\psi_B(u) = \begin{cases} u(1-u^2)^2, & 0 \leq |u| \leq 1 \\ 0, & |u| > 1. \end{cases}$$

Note that if  $\eta(u, v) = uv$  then the RA estimator defined above coincides with the previously defined least squares estimator of  $\phi_0$  based on the covariance of the residuals.

**Continuation of Example 1.** Let  $\Theta = \{(\phi_1, \phi_2, \phi_3)' \in \mathbb{R}^3 : \sup_{1 \leq p \leq 3} \{|\phi_p|\} < 1/3\}$ . The RA estimator of  $\phi_0$  based on the observations  $\tilde{\mathbf{X}}_M$  is the random vector  $\hat{\phi}_M^{RA} = (\hat{\phi}_{M,1}^{RA}, \hat{\phi}_{M,2}^{RA}, \hat{\phi}_{M,3}^{RA})' \in \Theta$ , such that

$$0 = \sum_{\underline{m} \in W_M} \eta \left( r_{\underline{m}}^* (\hat{\phi}_M^{RA}), r_{\underline{m}-\underline{t}_p}^* (\hat{\phi}_M^{RA}) \right) + \sum_{\underline{m} \in W_M} \sum_{j=1}^{2M} \sum_{u=0}^j \sum_{v=0}^u \left( \hat{\phi}_{M,1}^{RA} \right)^v \left( \hat{\phi}_{M,2}^{RA} \right)^{j-u} \left( \hat{\phi}_{M,3}^{RA} \right)^{u-v} \eta \left( r_{\underline{m}}^* (\hat{\phi}_M^{RA}), r_{\underline{m}-\underline{t}_p-(v+j-u, u-v)}^* (\hat{\phi}_M^{RA}) \right),$$

for all  $p = 1, 2, 3$ .

## 4 CONSISTENCY OF RA ESTIMATORS

In order to study the asymptotic behavior of RA estimator we will introduce some changes in the previous definitions. First, we eliminate the condition (6) and we assume that the process  $\tilde{\mathbf{X}}$  is observed in  $\mathbb{Z}^2$ . Now we redefine the residuals as follows

**Definition 4.** For all  $\underline{m} \in \mathbb{Z}^2$ , the residual with respect to  $\phi$  is the function  $R_{\underline{m}}: \Omega \times \Theta \rightarrow \mathbb{R}$  defined as

$$R_{\underline{m}}(\omega, \phi) = \frac{1}{\sigma_\varepsilon} \left( X_{\underline{m}}(\omega) - \sum_{p=1}^L \phi_p X_{\underline{m}-\underline{t}_p}(\omega) \right),$$

with  $\omega \in \Omega$  and  $\phi \in \Theta$ .

Note that  $R_{\underline{m}}(\phi)$  is  $r_{\underline{m}}^*(\phi)$  as given in Definition 3 when  $(\underline{m} - \underline{t}_p) \in I_0$  for all  $1 \leq p \leq L$ . Also

$$R_{\underline{m}}(\phi_0) = \varepsilon_{\underline{m}} / \sigma_\varepsilon, \text{ for all } \underline{m}. \quad (10)$$

**Definition 5.** The RA estimator of  $\phi_0$  based on the observations  $(X_{\underline{m}})_{\underline{m} \leq (M, M)}$  with domain  $\Omega'_{RA, M} \subset \Omega$  is defined as the function  $\hat{\phi}_M^{RA}: \Omega'_{RA, M} \rightarrow \Theta$ , such that

$$\frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \mathbf{\Gamma}_{\underline{m}, M}^\eta(\omega, \hat{\phi}_M^{RA}(\omega)) = \mathbf{0}, \quad (11)$$

for all  $\omega \in \Omega'_{RA, M}$ , and  $\mathbf{\Gamma}_{\underline{n}, N}^\eta: \Omega \times \Theta \rightarrow \mathbb{R}^L$  is given by

$$\left( \mathbf{\Gamma}_{\underline{n}, N}^\eta(\omega, \phi) \right)_p = \eta \left( R_{\underline{n}}(\omega, \phi), R_{\underline{n}-\underline{t}_p}(\omega, \phi) \right) + \sum_{j=1}^{N-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j, \tilde{\mathbf{k}}}(\phi) \eta \left( R_{\underline{n}}(\omega, \phi), R_{\underline{n}-\underline{t}_p-s(j, \tilde{\mathbf{k}})}(\omega, \phi) \right),$$

where  $\eta$  is a score function,  $1 \leq N \leq \infty$ ,  $\underline{n} \in \mathbb{Z}^2$ ,  $\omega \in \Omega$ ,  $\phi \in \Theta$  and  $1 \leq p \leq L$ .

**Remark 3.**

- (a) For  $N = \infty$  the convergence of the series involved in the definition of  $\mathbf{\Gamma}_{\underline{n}, \infty}^\eta$  is granted because of the definition of  $\Theta$  and the assumptions about the  $\eta$  function that we describe below.
- (b) As we mention before, when  $\sigma_\varepsilon$  is unknown, the computation of  $R_{\underline{m}}(\phi_0)$  can be done by plugging in (10) a robust estimator of the scale of  $F_\varepsilon$ .

Let us consider the following assumptions about  $\eta$ :

**Assumption 2.**



(i)  $\eta: \mathbb{R}^2 \rightarrow \mathbb{R}$  is a continuously differentiable function satisfying

$$\begin{aligned}\eta(0, v) &= 0, \\ \eta(u, 0) &= 0, \\ |\eta(u, v)| &\leq K_\eta,\end{aligned}$$

for some constant  $K_\eta < \infty$  and for all  $(u, v) \in \mathbb{R}^2$ .

(ii) Let  $\eta_1(u, v) = D_1\eta(u, v)$  and  $\eta_2(u, v) = D_2\eta(u, v)$  the partial derivatives of  $\eta$  with respect to the first and second components respectively. One of the two following conditions is satisfied:

(a) There are constants  $K_1, K_2$  such that

$$\begin{aligned}|\eta_1(u, v)| &\leq K_1, \\ |\eta_2(u, v)| &\leq K_2,\end{aligned}$$

for all  $(u, v)$  in  $\mathbb{R}^2$ .

(b) There exists a constant  $0 < K_3 < \infty$ , such that

$$\begin{aligned}|\eta_1(u, v)| &\leq K_3 |v|, \\ |\eta_2(u, v)| &\leq K_3 |u|.\end{aligned}$$

(iii)

$$E\left(\eta\left(\frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon}, \frac{\varepsilon_{\underline{m}'}}{\sigma_\varepsilon}\right)\right) = 0 \text{ if } \underline{m} \neq \underline{m}',$$

where  $\varepsilon_{\underline{m}}$  and  $\varepsilon_{\underline{m}'}$  are independent random variables with distribution  $F_\varepsilon$ .

(iv)

$$E\left(\eta_1\left(\frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon}, \frac{\varepsilon_{\underline{m}'}}{\sigma_\varepsilon}\right)\varepsilon_{\underline{m}'}\right) \neq 0 \text{ if } \underline{m} \neq \underline{m}',$$

where  $\varepsilon_{\underline{m}}$  and  $\varepsilon_{\underline{m}'}$  are independent random variables with distribution  $F_\varepsilon$ .

Now, we state the main result of this section:

**Theorem 1** (Existence and Consistency of the RA Estimator). *Given  $M_0 < \infty$ , there exists  $\Omega'' \subset \Omega$  with  $P(\Omega'') = 1$  such that*

$$\begin{aligned}\Omega'' \subset \Omega''' &= \{\omega \in \Omega: \text{there exists } N > M_0 \text{ such that} \\ M \geq N &\Rightarrow \exists \hat{\phi}_M^{RA}(\omega) \in \Theta \text{ with } \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \mathbf{I}_{\underline{m}, M}^\eta(\omega, \hat{\phi}_M^{RA}(\omega)) = \mathbf{0}, \text{ and } \lim_{M \rightarrow \infty} \hat{\phi}_M^{RA}(\omega) = \phi_0\}.\end{aligned}$$

The proof of this Theorem is given in the Subsection 6.1 of the Appendix.

## 5 ASYMPTOTIC NORMALITY OF RA ESTIMATORS

We will prove in the appendix that the covariance matrix of  $\mathbf{X}$ , say  $\Sigma(\mathbf{X}_T)$ , is a positive definite matrix. More precisely,  $\Sigma(\mathbf{X}_T)$  is given by  $\Sigma(\mathbf{X}_T) = \sigma_\varepsilon^2 \mathbf{I}_\infty(\phi_0)$  (see Lemma 1 in Subsection 6.1) where  $\mathbf{I}_\infty(\phi_0)$  is defined as follows.

Let

$$T - T = \{\underline{t} - \underline{s} : \underline{t} \text{ and } \underline{s} \text{ in } T\}, \quad (12)$$

and, for each  $\underline{v} \in T - T$ , let

$$\begin{aligned}(T - T)_1(\underline{v}) &= \{(j, \tilde{\mathbf{k}}) : j \geq 1, \tilde{\mathbf{k}} \in \mathbb{L}^j \text{ and } s(j, \tilde{\mathbf{k}}) = \underline{v}\}, \\ (T - T)_2(\underline{v}) &= \{(j, \tilde{\mathbf{k}}, l, \tilde{\mathbf{h}}) : j, l \geq 1, \tilde{\mathbf{k}} \in \mathbb{L}^j, \tilde{\mathbf{h}} \in \mathbb{L}^l \text{ and } s(j, \tilde{\mathbf{k}}) - s(l, \tilde{\mathbf{h}}) = \underline{v}\}.\end{aligned}$$

Note that

$$\underline{v} \notin I_0 \Rightarrow (T - T)_1(\underline{v}) = \emptyset. \quad (13)$$

So,  $\mathbf{I}_\infty(\phi_0)$  is the  $L \times L$  matrix given by

$$\mathbf{I}_\infty(\phi_0)_{p,p} = 1 + \sum_{(j,\tilde{\mathbf{k}},l,\tilde{\mathbf{h}}) \in (T-T)_2(\underline{0})} I_{j,\tilde{\mathbf{k}}}(\phi_0) I_{l,\tilde{\mathbf{h}}}(\phi_0), \quad (14)$$

for all  $1 \leq p \leq L$ , and

$$\begin{aligned} \mathbf{I}_\infty(\phi_0)_{p,q} = & \sum_{(j,\tilde{\mathbf{k}}) \in (T-T)_1(\underline{t}_p - \underline{t}_q)} I_{j,\tilde{\mathbf{k}}}(\phi_0) + \sum_{(j,\tilde{\mathbf{k}}) \in (T-T)_1(\underline{t}_q - \underline{t}_p)} I_{j,\tilde{\mathbf{k}}}(\phi_0) + \\ & \sum_{(j,\tilde{\mathbf{k}},l,\tilde{\mathbf{h}}) \in (T-T)_2(\underline{t}_p - \underline{t}_q)} I_{j,\tilde{\mathbf{k}}}(\phi_0) I_{l,\tilde{\mathbf{h}}}(\phi_0), \end{aligned} \quad (15)$$

for all  $p \neq q$ ,  $1 \leq p, q \leq L$ .

Now, in addition to the conditions given in Assumption 2 we need to introduce the following restriction on the score function.

**Assumption 3.**

$$E \left( \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \cdot \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon''}{\sigma_\varepsilon} \right) \right) = E \left( \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \cdot \eta \left( \frac{\varepsilon''}{\sigma_\varepsilon}, \frac{\varepsilon}{\sigma_\varepsilon} \right) \right) = 0,$$

where  $\varepsilon$ ,  $\varepsilon'$  and  $\varepsilon''$  are independent random variables with distribution function  $F_\varepsilon$ .

We now state the asymptotic distribution of the RA estimator.

**Theorem 2** (Asymptotic Normality of the RA Estimator). *Let  $(\hat{\phi}_M^{RA})_{M \geq M_0}$  be a sequence of random variables in  $\Theta$  such that*

$$\sqrt{\#(W_M)} \left[ \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \mathbf{\Gamma}_{\underline{m},M}^\eta(\hat{\phi}_M^{RA}) \right] \xrightarrow{P} \mathbf{0}, \text{ as } M \rightarrow \infty, \quad (16)$$

and

$$\hat{\phi}_M^{RA} \xrightarrow{P} \phi_0, \text{ as } M \rightarrow \infty.$$

Then

$$\sqrt{\#(W_M)} (\hat{\phi}_M^{RA} - \phi_0) \xrightarrow{D} N(\mathbf{0}, \sigma_A^2), \text{ as } M \rightarrow \infty,$$

where the asymptotic variance is given by

$$\sigma_A^2 = \sigma_\varepsilon^2 \frac{E \left( \left( \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \right)^2 \right)}{\left( E \left( \eta_1 \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right), \varepsilon' \right) \right)^2} (\mathbf{I}_\infty(\phi_0))^{-1},$$

with  $\varepsilon$ ,  $\varepsilon'$  are independent random variables with distribution function  $F_\varepsilon$ .

The proof of Theorem 2 is given in the Subsection 6.2 of the Appendix.

**Remark 4.** *Theorem 2 gives the asymptotic distribution of the least squares estimator of  $\phi_0$  based on the covariance of the residuals (LS estimator) when  $\eta(u, v) = uv$ . Hence, the efficiency of the RA estimator with respect to the LS estimator is given by*

$$\frac{1}{\sigma_\varepsilon^2 \frac{E \left( \left( \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \right)^2 \right)}{\left( E \left( \eta_1 \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right), \varepsilon' \right) \right)^2}}.$$

So, as discussed by in Bustos and Yohai (1986), the constants involved in the definition of the  $\eta$  function can be tuned.

## 6 APPENDIX: PROOFS

In order to simplify the presentation we introduce some additional notation and considerations. Let

$$\begin{aligned}\mathbf{G}_{M,N}(\omega, \phi) &= \frac{1}{\#(W_M)} \sum_{m \in W_M} \mathbf{\Gamma}_{m,N}^\eta(\omega, \phi), \\ \mathbf{g}_N(\phi) &= E \left( \mathbf{\Gamma}_{0,N}^\eta(\phi) \right), \\ \mathbf{F}_M(\omega, \phi) &= \mathbf{G}_{M,M}(\omega, \phi), \\ \mathbf{f}(\phi) &= \mathbf{g}_\infty(\phi),\end{aligned}\tag{17}$$

where  $1 \leq N \leq \infty$ ,  $M_0 \leq M < \infty$ ,  $\omega \in \Omega$  and  $\phi \in \Theta$ . Note that from (10), and (i) and (iii) of Assumption 2, it follows that

$$\mathbf{f}(\phi_0) = \mathbf{0}.\tag{18}$$

Since  $\eta$  is continuously differentiable, for each  $\omega \in \Omega$  the function  $\phi \mapsto \mathbf{F}_M(\omega, \phi)$  is continuously differentiable on  $\Theta$ , for every  $M$ .

Let  $\delta_0 > 0$  such that  $\Lambda = \{\phi \in \Theta : |\phi - \phi_0| < \delta_0\} \subset \Theta$ ; hence by (4), we have that

$$b = \sup \left\{ L \sup_{1 \leq p \leq L} |\phi_p| : \phi \in \Lambda \right\} < 1.\tag{19}$$

**Remark 5.** Let  $\phi \in \Lambda$ . Then

$$\left| \sum_{j=0}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j,\tilde{\mathbf{k}}}(\phi) \right| \leq \sum_{j=0}^{\infty} b^j.$$

In fact,  $\left| \sum_{j=0}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j,\tilde{\mathbf{k}}}(\phi) \right| \leq \sum_{j=0}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |I_{j,\tilde{\mathbf{k}}}(\phi)| = \sum_{j=0}^{\infty} \left( \sum_{p=1}^L |\phi_p| \right)^j \leq \sum_{j=0}^{\infty} b^j$ . Consider also  $0 < \delta_1 < \delta_0$ , and let  $C = B'(\delta_1) = \{\phi \in \mathbb{R}^L : |\phi - \phi_0| \leq \delta_1\}$ .

### 6.1 Proof of the Consistency of the RA estimator

The following lemmas and propositions are necessary to prove Theorem 1. This proof is given at the end of this subsection.

**Lemma 1.** Let  $\mathbf{I}_\infty(\phi_0)$  be as in (14) and (15) and let  $\Sigma(\mathbf{X}_T)$  be the covariance matrix of  $\mathbf{X}_T = (X_{t_1}, \dots, X_{t_L})'$ . Then  $\Sigma(\mathbf{X}_T)$  is positive definite and

$$\Sigma(\mathbf{X}_T) = \sigma_\varepsilon^2 \mathbf{I}_\infty(\phi_0).$$

Let us prove that the covariance matrix is positive definite. Note first that, since  $E(X_{t_p}) = 0$ , for all  $1 \leq p \leq L$ , then

$$\Sigma(\mathbf{X}_T) = [E(X_{t_p} X_{t_q})]_{1 \leq p, q \leq L}.$$

As  $\Sigma(\mathbf{X}_T)$  is non negative definite, by contradiction we suppose that  $\Sigma(\mathbf{X}_T)$  is not positive definite. Then there exists a vector  $\mathbf{a} = (a_1, \dots, a_L)' \neq (0, \dots, 0)$  such that

$$0 = \mathbf{a} \Sigma(\mathbf{X}_T) \mathbf{a}' = \sum_{p=1}^L \sum_{q=1}^L a_p a_q E(X_{t_p} X_{t_q}) = \sum_{p=1}^L \sum_{q=1}^L \langle a_p X_{t_p}, a_q X_{t_q} \rangle_{L^2} = \left\| \sum_{p=1}^L a_p X_{t_p} \right\|_{L^2}^2.$$

Hence

$$\sum_{p=1}^L a_p X_{t_p} = 0,\tag{20}$$

in  $L^2$ . Without loss of generality we may assume that the set  $T = \{t_1, \dots, t_L\}$  satisfies

$$\begin{cases} t_{p,1} \geq t_{q,1} & 1 \leq p < q \leq L \\ t_{p,2} > t_{q,2} & \text{if } t_{p,1} = t_{q,1}, 1 \leq p < q \leq L. \end{cases}\tag{21}$$

Considering that  $\mathbf{a}$  is different from the null vector and (20) then there exists

$$1 \leq p_1 < \cdots < p_r \leq L, \quad 2 \leq r \leq L, \quad (22)$$

such that

$$a_{p_i} \neq 0, \text{ for all } i = 1, \dots, r. \quad (23)$$

For every  $U \subset L^2(\Omega, F, P)$  we denote with  $\mathbb{H}(U)$  the closed vector subspace of  $L^2(\Omega, F, P)$  generated by  $U$ . So, using Proposition 1, we have that

$$X_{\underline{m}} \in \mathbb{H} \left( \left\{ \varepsilon_{\underline{m}} \right\} \cup \left\{ \varepsilon_{\underline{m}-s(j, \tilde{\mathbf{k}})} : j \geq 1, \tilde{\mathbf{k}} \in \mathbb{L}^j \right\} \right), \text{ for all } \underline{m} \in \mathbb{Z}^2.$$

Then, by (20), (22) and (23) we conclude that

$$\varepsilon_{\underline{t}_{p_1}} \in \mathbb{H} \left( \left\{ \varepsilon_{\underline{t}_{p_2}}, \dots, \varepsilon_{\underline{t}_{p_r}} \right\} \cup \left\{ \varepsilon_{\underline{t}_{p_i}-s(j, \tilde{\mathbf{k}})} : 1 \leq i \leq r, j \geq 1, \tilde{\mathbf{k}} \in \mathbb{L}^j \right\} \right). \quad (24)$$

Now, using (21), (22) and the definition of  $s(j, \tilde{\mathbf{k}})$ ,  $j \geq 1, \tilde{\mathbf{k}} \in \mathbb{L}^j$ , we establish that

$$\varepsilon_{\underline{t}_{p_1}} \notin \left( \left\{ \varepsilon_{\underline{t}_{p_2}}, \dots, \varepsilon_{\underline{t}_{p_r}} \right\} \cup \left\{ \varepsilon_{\underline{t}_{p_i}-s(j, \tilde{\mathbf{k}})} : 1 \leq i \leq r, j \geq 1, \tilde{\mathbf{k}} \in \mathbb{L}^j \right\} \right).$$

Since  $\tilde{\varepsilon} = (\varepsilon_{\underline{m}})_{\underline{m} \in \mathbb{Z}^2}$  is a white noise, (24) is impossible. By this contradiction we conclude that the covariance matrix is positive definite.

The equality  $\Sigma(\mathbf{X}_T) = \sigma_\varepsilon^2 \mathbf{I}_\infty(\phi_0)$  follows from a straightforward calculation.

**Lemma 2.** For each  $\underline{m} \in \mathbb{Z}^2$ , let  $\mathbf{Z}_{\underline{m}}$  be the random vector with values in  $\mathbb{R}^{L+1}$  given by

$$\mathbf{Z}_{\underline{m}} = (X_{\underline{m}}, X_{\underline{m}-\underline{t}_1}, \dots, X_{\underline{m}-\underline{t}_L})'.$$

Let  $c_0: \Lambda \rightarrow \mathbb{R}$  be a continuous function. For each  $j \geq 1$  and  $\tilde{\mathbf{k}} \in \mathbb{L}^j$  let  $c_{j, \tilde{\mathbf{k}}}: \Lambda \rightarrow \mathbb{R}$  be a continuous function. Assume that

**c0)** There exists a sequence of positive real numbers,  $(b_j)_{j \geq 0}$ , such that

$$\begin{aligned} \sum_{j=0}^{\infty} b_j &< \infty, \\ \sup_{\phi \in \Lambda} |c_0(\phi)| &\leq b_0, \text{ and} \\ \sup_{\phi \in \Lambda} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |c_{j, \tilde{\mathbf{k}}}(\phi)| &\leq b_j, \forall j \geq 1. \end{aligned}$$

Let  $W: \mathbb{R}^{L+1} \times \mathbb{R}^{L+1} \times \Lambda \rightarrow \mathbb{R}$  be a function satisfying the following assumptions:

**w1)** There exists  $K_W < \infty$  such that for each  $(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^{L+1} \times \mathbb{R}^{L+1}$

$$\sup_{\phi \in \Lambda} |W(\mathbf{x}, \mathbf{y}, \phi)| \leq K_W |\rho(\mathbf{x}, \mathbf{y})|,$$

where  $\rho$  is the function  $\rho: \mathbb{R}^{L+1} \times \mathbb{R}^{L+1} \rightarrow \mathbb{R}$  such that there exists a constant  $\kappa$  satisfying

$$E(|\rho(\mathbf{Z}_{\underline{m}}, \mathbf{Z}_{\underline{n}})|) \leq \kappa \|\mathbf{X}_0\|_{L^2}, \quad (25)$$

for all  $\underline{m}, \underline{n} \in \mathbb{Z}^2$ .

**w2)** Given  $\mathbb{K} \subset \mathbb{R}^{L+1}$  a compact set and  $d > 0$ , there exists  $d' > 0$  such that

$$|\phi - \phi^*| < d', \phi \text{ and } \phi^* \in \Lambda \implies |W(\mathbf{x}, \mathbf{y}, \phi) - W(\mathbf{x}, \mathbf{y}, \phi^*)| < d, \text{ for all } \mathbf{x}, \mathbf{y} \in \mathbb{K}.$$

Let  $\underline{n}_1 \in \mathbb{Z}^2$ . For each  $1 \leq N$  integer and  $\underline{n} \in \mathbb{Z}^2$  let  $\Psi_{\underline{n}, N}: \Omega \times \Lambda \rightarrow \mathbb{R}$  be given by

$$\Psi_{\underline{n}, N}(\omega, \phi) = c_0(\phi) W(\mathbf{Z}_{\underline{n}}(\omega), \mathbf{Z}_{\underline{n}-\underline{n}_1}(\omega), \phi) + \sum_{j=1}^{N-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} c_{j, \tilde{\mathbf{k}}}(\phi) W(\mathbf{Z}_{\underline{n}}(\omega), \mathbf{Z}_{\underline{n}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}(\omega), \phi).$$

Then

1)  $(\Psi_{\underline{0}, N}(\phi))_{N \geq 1}$  converges uniformly on  $\phi \in \Lambda$  in  $L^2(\Omega, \mathbb{F}, P)$ .

Let us denote the limit as

$$\Psi_{\underline{0}, \infty}(\omega, \phi) = c_0(\phi) W(\mathbf{Z}_{\underline{0}}(\omega), \mathbf{Z}_{\underline{0}-\underline{n}_1}(\omega), \phi) + \sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} c_{j, \tilde{\mathbf{k}}}(\phi) W(\mathbf{Z}_{\underline{0}}(\omega), \mathbf{Z}_{\underline{0}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}(\omega), \phi).$$

2) Writing

$$T_{M, N}(\omega, \phi) = \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \Psi_{\underline{m}, N}(\omega, \phi)$$

there exists a strictly increasing sequence  $(J(N))_N$  such that for all  $N$  there exists  $\Omega_N \subseteq \Omega$  with  $P(\Omega_N) = 1$  satisfying

$$\limsup_{M \rightarrow \infty} \sup_{\phi \in C} |T_{M, M}(\omega, \phi) - E(\Psi_{\underline{0}, \infty}(\phi))| \leq \limsup_{M \rightarrow \infty} \sup_{\phi \in C} |T_{M, J(N)}(\omega, \phi) - E(\Psi_{\underline{0}, J(N)}(\phi))| + \frac{1}{N},$$

if  $\omega \in \Omega_N$ .

3) For each  $N$  there exists  $\Omega'_N$  with  $P(\Omega'_N) = 1$ , satisfying  $\lim_{M \rightarrow \infty} \sup_{\phi \in C} |T_{M, N}(\omega, \phi) - E(\Psi_{\underline{0}, N}(\phi))| = 0$  if  $\omega \in \Omega'_N$ .

4) There exists a subset  $\Omega_0$  of  $\Omega$  with  $P(\Omega_0) = 1$ , such that

$$\Omega_0 \subset \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} \left| \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \Psi_{\underline{m}, M}(\omega, \phi) - E(\Psi_{\underline{0}, \infty}(\phi)) \right| = 0 \right\}.$$

**PROOF:** The proof of 1) is an immediate consequence of c0), w1) and (25). To prove 2), let  $J(N)$  be the minimum positive integer such that for each  $N$ ,

$$K_W \cdot \kappa \cdot \|X_{\underline{0}}\|_{L^2} \sum_{j=J(N)}^{\infty} b_j < \frac{1}{2N}.$$

Then

$$\begin{aligned} & E \left( \sup_{\phi \in C} \sum_{j=J(N)}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |c_{j, \tilde{\mathbf{k}}}(\phi) W(\mathbf{Z}_{\underline{0}}, \mathbf{Z}_{\underline{0}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}(\omega), \phi)| \right) \\ & \leq K_W \cdot \sup_{\phi \in C} \sum_{j=J(N)}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |c_{j, \tilde{\mathbf{k}}}(\phi)| E \left( \left| \rho(\mathbf{Z}_{\underline{0}}(\omega), \mathbf{Z}_{\underline{0}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}(\omega)) \right| \right) \\ & \leq K_W \cdot \kappa \cdot \|X_{\underline{0}}\|_{L^2} \sup_{\phi \in C} \sum_{j=J(N)}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |c_{j, \tilde{\mathbf{k}}}(\phi)| \\ & \leq K_W \cdot \kappa \cdot \|X_{\underline{0}}\|_{L^2} \sum_{j=J(N)}^{\infty} b_j \\ & < \frac{1}{2N}. \end{aligned}$$

Hence, for each  $\omega \in \Omega$  and  $M > J(N)$ ,

$$\begin{aligned} & \sup_{\phi \in C} |T_{M,M}(\omega, \phi) - E(\Psi_{\underline{0}, \infty}(\phi))| \\ & \leq \sup_{\phi \in C} |T_{M, J(N)}(\omega, \phi) - E(\Psi_{\underline{0}, J(N)}(\phi))| \\ & \quad + \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \left( \sup_{\phi \in C} \sum_{j=J(N)}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |c_{j, \tilde{\mathbf{k}}}(\phi) W(\mathbf{Z}_{\underline{m}}(\omega), \mathbf{Z}_{\underline{m}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}(\omega), \phi)| \right) \\ & \quad + E \left( \sup_{\phi \in C} \sum_{j=J(N)}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |c_{j, \tilde{\mathbf{k}}}(\phi) W(\mathbf{Z}_{\underline{0}}, \mathbf{Z}_{\underline{0}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}, \phi)| \right). \end{aligned}$$

By the ergodicity, there exists  $\Omega_N \subseteq \Omega$  with  $P(\Omega_N) = 1$ , such that if  $\omega \in \Omega_N$ , then

$$\begin{aligned} & \limsup_{M \rightarrow \infty} \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \left( \sup_{\phi \in C} \sum_{j=J(N)}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |c_{j, \tilde{\mathbf{k}}}(\phi) W(\mathbf{Z}_{\underline{m}}(\omega), \mathbf{Z}_{\underline{m}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}(\omega), \phi)| \right) \\ & = E \left( \sup_{\phi \in C} \sum_{j=J(N)}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} |c_{j, \tilde{\mathbf{k}}}(\phi) W(\mathbf{Z}_{\underline{0}}, \mathbf{Z}_{\underline{0}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}, \phi)| \right). \end{aligned}$$

Thus, for each  $\omega \in \Omega_N$  we have that

$$\limsup_{M \rightarrow \infty} \sup_{\phi \in C} |T_{M,M}(\omega, \phi) - E(\Psi_{\underline{0}, \infty}(\phi))| \leq \limsup_{M \rightarrow \infty} \sup_{\phi \in C} |T_{M, J(N)}(\omega, \phi) - E(\Psi_{\underline{0}, J(N)}(\phi))| + 2 \left( \frac{1}{2N} \right).$$

Let us prove 3). Consider  $N \geq 1$  and, for each  $\phi \in C$ , let  $\Omega'_N(\phi)$  with  $P(\Omega'_N(\phi)) = 1$ , such that

$$\Omega'_N(\phi) \subseteq \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} |T_{M, N}(\omega, \phi) - E(\Psi_{\underline{0}, N}(\phi))| = 0 \right\}.$$

Let  $n \geq 1$ . Since  $\left| \rho(\mathbf{Z}_{\underline{0}}, \mathbf{Z}_{\underline{0}-\underline{n}_1-s(j, \tilde{\mathbf{k}})}) \right|$  is integrable for all  $\tilde{\mathbf{k}} \in \mathbb{L}^j, 1 \leq j \leq N-1$ , there exists  $0 < K_{N,n} < \infty$ , such that

$$\sup_{\tilde{\mathbf{k}} \in \mathbb{L}^j, 1 \leq j \leq N-1} \left( E(1_{A(K_{N,n})^c} |\rho(\mathbf{Z}_{\underline{0}}, \mathbf{Z}_{\underline{0}-\underline{n}_1})|), E(1_{A(K_{N,n})^c} |\rho(\mathbf{Z}_{\underline{0}}, \mathbf{Z}_{\underline{0}-\underline{n}_1-s(j, \tilde{\mathbf{k}})})|) \right) < \frac{1}{6nK_W \left( \sum_{j=0}^{\infty} b_j \right)}, \quad (26)$$

where

$$\begin{aligned} A(K_{N,n}) & = \left\{ \omega \in \Omega : \sup_{\underline{s} \in S_N} |\mathbf{Z}_{\underline{s}}(\omega)| \leq K_{N,n} \right\}, \\ S_N & = \{ \underline{0}, \underline{0} - \underline{n}_1 \} \cup \left\{ \underline{0} - \underline{n}_1 - s(j, \tilde{\mathbf{k}}) : \tilde{\mathbf{k}} \in \mathbb{L}^j, 1 \leq j \leq N-1 \right\}. \end{aligned}$$

Let  $\phi \in C$ . Considering that the functions  $c_0$  and  $c_{j, \tilde{\mathbf{k}}}, \tilde{\mathbf{k}} \in \mathbb{L}^j, 1 \leq j \leq N-1$ , are continuous and also  $c_0, w_1, w_2$ ,

the definition of  $A(K_{N,n})$  and (26) we conclude that there exists a neighborhood,  $V_{\phi,n}$ , of  $\phi$  contained in  $\Lambda$  such that:

$$\begin{aligned}
& E \left( \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{0},N}(\phi^*) - \Psi_{\underline{0},N}(\phi)| \right) \\
& \leq E \left( \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{0},N}(\phi^*) 1_{A(K_{N,n})} - \Psi_{\underline{0},N}(\phi) 1_{A(K_{N,n})}| \right) \\
& \quad + E \left( \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{0},N}(\phi^*) 1_{A(K_{N,n})^c} - \Psi_{\underline{0},N}(\phi) 1_{A(K_{N,n})^c}| \right) \\
& \leq E \left( \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{0},N}(\phi^*) 1_{A(K_{N,n})} - \Psi_{\underline{0},N}(\phi) 1_{A(K_{N,n})}| \right) \\
& \quad + E \left( \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{0},N}(\phi^*) 1_{A(K_{N,n})^c}| \right) + E (|\Psi_{\underline{0},N}(\phi) 1_{A(K_{N,n})^c}|) \\
& < \frac{1}{6n} + \frac{1}{6n} + \frac{1}{6n} = \frac{1}{2n}.
\end{aligned}$$

Now, for each  $\omega \in \Omega$ ,

$$\begin{aligned}
& \sup_{\phi^* \in C \cap V_{\phi,n}} |T_{M,N}(\omega, \phi^*) - E(\Psi_{\underline{0},N}(\phi^*))| \\
& \leq |T_{M,N}(\omega, \phi) - E(\Psi_{\underline{0},N}(\phi))| \\
& \quad + \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{m},N}(\omega, \phi^*) - \Psi_{\underline{m},N}(\omega, \phi)| \\
& \quad + \sup_{\phi^* \in C \cap V_{\phi,n}} |E(\Psi_{\underline{0},N}(\phi^*)) - E(\Psi_{\underline{0},N}(\phi))|. \tag{27}
\end{aligned}$$

Also,

$$\sup_{\phi^* \in C \cap V_{\phi,n}} |E(\Psi_{\underline{0},N}(\phi^*)) - E(\Psi_{\underline{0},N}(\phi))| \leq E \left( \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{0},N}(\phi^*) - \Psi_{\underline{0},N}(\phi)| \right) < \frac{1}{2n}. \tag{28}$$

Because of the definition of  $\Omega_N''(\phi)$ , it follows that for all  $\omega \in \Omega_N''(\phi)$ :

$$\lim_{M \rightarrow \infty} |T_{M,N}(\omega, \phi) - E(\Psi_{\underline{0},N}(\phi))| = 0. \tag{29}$$

By ergodicity, there exists  $\Omega_N(\phi, n) \subset \Omega_N''(\phi)$  with  $P(\Omega_N(\phi, n)) = 1$  such that, if  $\omega \in \Omega_N(\phi, n)$ , then

$$\lim_{M \rightarrow \infty} \left| \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{m},N}(\omega, \phi^*) - \Psi_{\underline{m},N}(\omega, \phi)| \right| = E \left( \sup_{\phi^* \in C \cap V_{\phi,n}} |\Psi_{\underline{0},N}(\phi^*) - \Psi_{\underline{0},N}(\phi)| \right) < \frac{1}{2n}.$$

Hence, by (27), (28), (29) and (6.1),

$$\limsup_{M \rightarrow \infty} \sup_{\phi^* \in C \cap V_{\phi,n}} |T_{M,N}(\omega, \phi^*) - E(\Psi_{\underline{0},N}(\phi^*))| < \frac{1}{n}, \tag{30}$$

for all  $\omega \in \Omega_N(\phi, n)$ .

Since  $C$  is a compact set, there exists a finite set  $\{\phi_1, \dots, \phi_r\}$  contained in  $C$  such that

$$C = \bigcup_{i=1}^r (C \cap V_{\phi_i,n}).$$

Thus, if we define  $\Omega'_N(n) = \bigcap_{i=1}^r \Omega_N(\phi_i, n)$ , then  $P(\Omega'_N(n)) = 1$ . By (30),

$$\limsup_{M \rightarrow \infty} \sup_{\phi^* \in C} |T_{M,N}(\omega, \phi^*) - E(\Psi_{\underline{0},N}(\phi^*))| \leq \frac{1}{n}, \quad (31)$$

for all  $\omega \in \Omega'_N(n)$ . Setting  $\Omega'_N = \bigcap_{n=1}^{\infty} \Omega'_N(n)$ , then  $P(\Omega'_N) = 1$ , and because of (31) we have that

$$\lim_M \sup_{\phi^* \in C} |T_{M,N}(\omega, \phi^*) - E(\Psi_{\underline{0},N}(\phi^*))| = 0, \quad (32)$$

for all  $\omega \in \Omega'_N$ . The proof of 3) is completed. To prove 4), it is enough to set  $\Omega_0 = \bigcap_{N \geq 1} (\Omega_N \cap \Omega'_N)$ . ■

Next lemma recalls a well known result and useful result in robustness.

**Lemma 3** (The Zeros Lemma (Ruskin (1978))). *Let  $U \subset \mathbb{R}^k$  be an open set,  $\lambda_0 \in U$ , for each  $n = 1, 2, \dots$  let  $\mathbf{q}_n: U \rightarrow \mathbb{R}^k$  and  $\mathbf{q}: U \rightarrow \mathbb{R}^k$  be continuously differentiable functions. Assume that*

1)  $\mathbf{q}(\lambda_0) = \mathbf{0}$ .

2)  $\mathbb{D}\mathbf{q}(\lambda_0)$  is not zero.

3) *There exists  $\gamma > 0$  such that  $(\mathbf{q}_n)_n$  and  $(\mathbb{D}\mathbf{q}_n)_n$  converge uniformly to  $\mathbf{q}$  and  $\mathbb{D}\mathbf{q}$  respectively on  $B(\lambda_0, \gamma) = \{\lambda \in \mathbb{R}^k: \sup_{1 \leq i \leq k} |\lambda_i - \lambda_{0i}| < \gamma\}$ .*

*Then, there exist  $n_0 \geq 1$  and a sequence  $(\lambda_n)_{n \geq n_0}$  in  $B(\lambda_0, \gamma)$  such that  $(\lambda_n)_n$  converges to  $\lambda_0$  and  $\mathbf{q}_n(\lambda_n) = 0$ , for all  $n \geq n_0$ .*

**Proposition 2.** *Let  $\mathbf{f}$  be as in (17). Then  $\mathbf{f}$  is continuously differentiable on  $\Lambda$  and satisfies*

1) *For each  $1 \leq p, q \leq L$  and for each  $\phi \in \Lambda$ :*

$$D_q \left( (\mathbf{f}(\phi))_p \right) = A_{1p,q}(\phi) + A_{2p,q}(\phi) + \sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left( A_{3p,q}(j, \tilde{\mathbf{k}}, \phi) + A_{4p,q}(j, \tilde{\mathbf{k}}, \phi) + A_{5p,q}(j, \tilde{\mathbf{k}}, \phi) \right),$$

where

$$\begin{aligned} A_{1p,q}(\phi) &= -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( R_{\underline{0}}(\phi), R_{-\underline{t}_p}(\phi) \right) X_{-\underline{t}_q} \right), \\ A_{2p,q}(\phi) &= -\frac{1}{\sigma_\varepsilon} E \left( \eta_2 \left( R_{\underline{0}}(\phi), R_{-\underline{t}_p}(\phi) \right) X_{-\underline{t}_p - \underline{t}_q} \right), \\ A_{3p,q}(j, \tilde{\mathbf{k}}, \phi) &= D_q \left( I_{j, \tilde{\mathbf{k}}}(\phi) \right) E \left( \eta \left( R_{\underline{0}}(\phi), R_{-\underline{t}_p - s(j, \tilde{\mathbf{k}})}(\phi) \right) \right), \\ A_{4p,q}(j, \tilde{\mathbf{k}}, \phi) &= -\frac{1}{\sigma_\varepsilon} I_{j, \tilde{\mathbf{k}}}(\phi) E \left( \eta_1 \left( R_{\underline{0}}(\phi), R_{-\underline{t}_p - s(j, \tilde{\mathbf{k}})}(\phi) \right) X_{-\underline{t}_q} \right), \\ A_{5p,q}(j, \tilde{\mathbf{k}}, \phi) &= -\frac{1}{\sigma_\varepsilon} I_{j, \tilde{\mathbf{k}}}(\phi) E \left( \eta_2 \left( R_{\underline{0}}(\phi), R_{-\underline{t}_p - s(j, \tilde{\mathbf{k}})}(\phi) \right) X_{-\underline{t}_p - \underline{t}_q - s(j, \tilde{\mathbf{k}})} \right). \end{aligned}$$

2)  $\mathbb{D}\mathbf{f}(\phi_0) = -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \varepsilon' \right) \mathbf{I}_\infty(\phi_0)$ , where  $\varepsilon$  and  $\varepsilon'$  are independent random variables with distribution  $F_\varepsilon$ , and  $\mathbb{D}\mathbf{f}(\phi_0)$  is not zero.

**PROOF:** First we will prove that

$$(\mathbf{g}_N)_N \text{ converges uniformly to } \mathbf{f} \text{ on } \Lambda. \quad (33)$$

In fact, for any  $\phi \in \Lambda$ ,  $1 \leq p \leq L$  and  $N$  we have

$$\left| (\mathbf{f}(\phi))_p - (\mathbf{g}_N(\phi))_p \right| = \sum_{j=N}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j, \tilde{\mathbf{k}}}(\phi) E \left( \left| \eta \left( R_{\underline{0}}(\phi), R_{-\underline{t}_p - s(j, \tilde{\mathbf{k}})}(\phi) \right) \right| \right) \leq K_\eta \sum_{j=N}^{\infty} b^j,$$

being  $b$  as in (19), this implies (33). Then, by a classical result in analysis, to prove 1) in Proposition 2, it is enough to show that

$$\left( D_q \left( (\mathbf{g}_N(\phi))_p \right) \right)_N \text{ converges uniformly on } \Lambda. \quad (34)$$



Notice that for each  $\phi \in \Lambda$

$$D_q \left( (\mathbf{g}_N(\phi))_p \right) = A1_{p,q}(\phi) + A2_{p,q}(\phi) + \sum_{j=1}^{N-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left( A3_{p,q}(j, \tilde{\mathbf{k}}, \phi) + A4_{p,q}(j, \tilde{\mathbf{k}}, \phi) + A5_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right).$$

To prove (34), it is enough to show that for each  $d > 0$ , there exists a positive integer  $N$  such that for all positive integer  $n$

$$\sup_{\phi \in \Lambda} \left| D_q \left( (\mathbf{g}_{N+n}(\phi))_p \right) - D_q \left( (\mathbf{g}_N(\phi))_p \right) \right| < d. \quad (35)$$

Now,

$$\left| A3_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| = K_\eta \left| D_q \left( I_{j, \tilde{\mathbf{k}}}(\phi) \right) \right|.$$

Then

$$\sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left| A3_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| \leq K_\eta \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left| D_q \left( I_{j, \tilde{\mathbf{k}}}(\phi) \right) \right|. \quad (36)$$

Since

$$\sum_{\tilde{\mathbf{k}} \in \mathbb{L}^h} I_{h, \tilde{\mathbf{k}}}(\phi) = \left( \sum_{p=1}^L \phi_p \right)^h, \quad (37)$$

we have that

$$\sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} D_q \left( I_{j, \tilde{\mathbf{k}}}(\phi) \right) = D_q \left( \left( \sum_{p'=1}^L \phi_{p'} \right)^j \right) = j \left( \sum_{p'=1}^L \phi_{p'} \right)^{j-1}.$$

Then, by (36), it follows that

$$\sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left| A3_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| \leq K_\eta j \left( \sum_{p'=1}^L |\phi_{p'}| \right)^{j-1} \leq K_\eta j (b)^{j-1}. \quad (38)$$

Assume that (a) of Assumption 2-(ii) holds. Then

$$\begin{aligned} \left| A4_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| &\leq \frac{1}{\sigma_\varepsilon} \left| I_{j, \tilde{\mathbf{k}}}(\phi) \right| K_1 E(|X_0|), \\ \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left| A5_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| &\leq \frac{1}{\sigma_\varepsilon} K_2 E(|X_0|) \left( \sum_{p'=1}^L |\phi_{p'}| \right)^j, \end{aligned}$$

for all  $j$ .

Thus, for each  $N, n$  positive integers and  $\phi \in \Lambda$  we have

$$\begin{aligned} &\left| D_q \left( (\mathbf{g}_{N+n}(\phi))_p \right) - D_q \left( (\mathbf{g}_N(\phi))_p \right) \right| \\ &\leq \sum_{j=N}^{N+n} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left( \left| A3_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| + \left| A4_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| + \left| A5_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| \right) \\ &\leq \sum_{j=N}^{N+n} \left( K_\eta j (b)^{j-1} + \frac{1}{\sigma_\varepsilon} K_1 E(|X_0|) b^j + \frac{1}{\sigma_\varepsilon} K_2 E(|X_0|) b^j \right). \end{aligned}$$

Hence (35) is satisfied. Now, let us consider (b) of Assumption 2-(ii) instead of (a). By the Cauchy-Schwartz inequality

$$\left| A4_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| \leq \frac{K_3}{\sigma_\varepsilon} \left| I_{j, \tilde{\mathbf{k}}}(\phi) \right| \left( \sum_{p'=1}^L |\phi_{p'}| \right) E(|X_0|^2).$$

Hence

$$\sum_{j=N}^{N+n} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left| A4_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| \leq \frac{K_3}{\sigma_\varepsilon} E \left( |X_{\underline{0}}|^2 \right) \sum_{j=N}^{N+n} \left( \sum_{p'=1}^L |\phi_{p'}| \right)^{j+1}, \quad (39)$$

for all  $N$ .

Similarly, considering (b) of Assumption 2-(ii) instead of (a) of that assumption:

$$\sum_{j=N}^{N+n} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left| A5_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| \leq \frac{K_3}{\sigma_\varepsilon} E \left( |X_{\underline{0}}|^2 \right) \sum_{j=N}^{N+n} \left( \sum_{p'=1}^L |\phi_{p'}| \right)^{j+1}, \quad (40)$$

for all  $N$ . Thus, using (38), (39) and (40), we have that

$$\left| D_q \left( (\mathbf{g}_{N+n}(\phi))_p \right) - D_q \left( (\mathbf{g}_N(\phi))_p \right) \right| \leq \sum_{j=N}^{N+n} \left( K_\eta j (b)^{j-1} + 2 \frac{K_3}{\sigma_\varepsilon} E \left( |X_{\underline{0}}|^2 \right) b^j \right).$$

Hence the proof of (35) is completed. This finishes the proof of 1) in Proposition 2.

Now, let us prove 2) in Proposition 2.

$$\begin{aligned} A1_{p,q}(\phi_0) &= -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( R_{\underline{0}}(\phi_0), R_{-t_p}(\phi_0) \right) X_{-t_q} \right) \\ &= -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( \varepsilon_{\underline{0}}, \varepsilon_{-t_p} \right) X_{-t_q} \right) \\ &= -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( \varepsilon_{\underline{0}}, \varepsilon_{-t_p} \right) \left( \varepsilon_{-t_q} + \sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j,\tilde{\mathbf{k}}}(\phi_0) \varepsilon_{-t_q-s(j,\tilde{\mathbf{k}})} \right) \right). \end{aligned}$$

Then

$$A1_{p,q}(\phi_0) = \begin{cases} -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \varepsilon' \right) & \text{if } p = q \\ -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \varepsilon' \right) \sum_{(j,\tilde{\mathbf{k}}) \in (T-T)_1(t_p-t_q)} I_{j,\tilde{\mathbf{k}}}(\phi_0) & \text{if } p \neq q \end{cases}, \quad (41)$$

where  $\varepsilon$  and  $\varepsilon'$  are independent random variables with distribution  $F_\varepsilon$ .

$$\begin{aligned} A2_{p,q}(\phi_0) &= -\frac{1}{\sigma_\varepsilon} E \left( \eta_2 \left( R_{\underline{0}}(\phi_0), R_{-t_p}(\phi_0) \right) X_{-t_p-t_q} \right) \\ &= -\frac{1}{\sigma_\varepsilon} E \left( \eta_2 \left( \varepsilon_{\underline{0}}, \varepsilon_{-t_p} \right) \left( \varepsilon_{-t_p-t_q} + \sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j,\tilde{\mathbf{k}}}(\phi_0) \varepsilon_{-t_p-t_q-s(j,\tilde{\mathbf{k}})} \right) \right) = 0. \end{aligned} \quad (42)$$

Now

$$\begin{aligned} A3_{p,q}(j, \tilde{\mathbf{k}}, \phi_0) &= D_q \left( I_{j,\tilde{\mathbf{k}}}(\phi_0) \right) E \left( \eta \left( R_{\underline{0}}(\phi_0), R_{-t_p-s(j,\tilde{\mathbf{k}})}(\phi_0) \right) \right) \\ &= D_q \left( I_{j,\tilde{\mathbf{k}}}(\phi_0) \right) E \left( \eta \left( \frac{\varepsilon_{\underline{0}}}{\sigma_\varepsilon}, \frac{\varepsilon_{-t_p-s(j,\tilde{\mathbf{k}})}}{\sigma_\varepsilon} \right) \right) = 0, \end{aligned} \quad (43)$$

by Assumption 2-(iii). Consider now

$$\begin{aligned} A5_{p,q}(j, \tilde{\mathbf{k}}, \phi_0) &= -\frac{1}{\sigma_\varepsilon} I_{j,\tilde{\mathbf{k}}}(\phi_0) E \left( \eta_2 \left( R_{\underline{0}}(\phi_0), R_{-t_p-s(j,\tilde{\mathbf{k}})}(\phi_0) \right) X_{-t_p-s(j,\tilde{\mathbf{k}})-t_q} \right) \\ &= -\frac{1}{\sigma_\varepsilon} I_{j,\tilde{\mathbf{k}}}(\phi_0) E \left( \eta_2 \left( \frac{\varepsilon_{\underline{0}}}{\sigma_\varepsilon}, \frac{\varepsilon_{-t_p-s(j,\tilde{\mathbf{k}})}}{\sigma_\varepsilon} \right) X_{-t_p-s(j,\tilde{\mathbf{k}})-t_q} \right). \end{aligned}$$

Now, since  $t_q \neq \underline{0}$  and (5) we have that  $\varepsilon_{\underline{0}}$ ,  $\varepsilon_{-t_p-s(j,\tilde{\mathbf{k}})}$  and  $X_{-t_p-s(j,\tilde{\mathbf{k}})-t_q}$  are independent. Then, since  $E \left( X_{-t_p-s(j,\tilde{\mathbf{k}})-t_q} \right) = 0$ , it follows that

$$A5_{p,q}(j, \tilde{\mathbf{k}}, \phi_0) = 0. \quad (44)$$

Finally,

$$\begin{aligned} A4_{p,q}(j, \tilde{\mathbf{k}}, \phi_0) &= -\frac{1}{\sigma_\varepsilon} I_{j, \tilde{\mathbf{k}}}(\phi_0) E \left( \eta_1 \left( R_0(\phi_0), R_{-t_p-s(j, \tilde{\mathbf{k}})}(\phi_0) \right) X_{-t_q} \right) \\ &= -\frac{1}{\sigma_\varepsilon} I_{j, \tilde{\mathbf{k}}}(\phi_0) E \left( \eta_1 \left( \frac{\varepsilon_0}{\sigma_\varepsilon}, \frac{\varepsilon_{-t_p-s(j, \tilde{\mathbf{k}})}}{\sigma_\varepsilon} \right) \varepsilon_{-t_q} \right) + \sum_{l=1}^{\infty} \sum_{\tilde{\mathbf{h}} \in \mathbb{L}^l} F(j, \tilde{\mathbf{k}}, l, \tilde{\mathbf{h}}), \end{aligned}$$

where

$$F(j, \tilde{\mathbf{k}}, l, \tilde{\mathbf{h}}) = -\frac{1}{\sigma_\varepsilon} I_{j, \tilde{\mathbf{k}}}(\phi_0) I_{l, \tilde{\mathbf{h}}}(\phi_0) E \left( \eta_1 \left( \frac{\varepsilon_0}{\sigma_\varepsilon}, \frac{\varepsilon_{-t_p-s(j, \tilde{\mathbf{k}})}}{\sigma_\varepsilon} \right) \varepsilon_{-t_q-s(l, \tilde{\mathbf{h}})} \right).$$

Now, if  $-t_p - s(j, \tilde{\mathbf{k}}) \neq -t_q - s(l, \tilde{\mathbf{h}})$ , then

$$E \left( \eta_1 \left( \frac{\varepsilon_0}{\sigma_\varepsilon}, \frac{\varepsilon_{-t_p-s(j, \tilde{\mathbf{k}})}}{\sigma_\varepsilon} \right) \varepsilon_{-t_q-s(l, \tilde{\mathbf{h}})} \right) = 0,$$

Therefore

$$\sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} A4_{p,q}(j, \tilde{\mathbf{k}}, \phi) = -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \varepsilon' \right) A4_{p,q}^* \quad (45)$$

where  $\varepsilon$  and  $\varepsilon'$  are independent random variables with distribution  $F_\varepsilon$  and

$$A4_{p,q}^* = \left( \sum_{(j, \tilde{\mathbf{k}}) \in (T-T)_1(t_q - t_p)} I_{j, \tilde{\mathbf{k}}}(\phi_0) + \sum_{(j, \tilde{\mathbf{k}}, l, \tilde{\mathbf{h}}) \in (T-T)_2(t_p - t_q)} I_{j, \tilde{\mathbf{k}}}(\phi_0) I_{l, \tilde{\mathbf{h}}}(\phi_0) \right).$$

By (41), (42), (43), (44), (45) and by the definition of  $\mathbf{I}_\infty(\phi_0)$  it follows that

$$\mathbb{D}\mathbf{f}(\phi_0) = -E \left( \eta_1 \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \varepsilon' \right) \mathbf{I}_\infty(\phi_0),$$

where  $\varepsilon$  and  $\varepsilon'$  are independent random variables with distribution  $F_\varepsilon$ .

Now, by the Lemmas 1 and Assumption 2-(iv),  $\mathbb{D}\mathbf{f}(\phi_0)$ , with the exception of a non zero constant, is equal to the covariance matrix of  $(X_m)_{m \in T}$  and, hence, it is invertible. This finishes the proof of the Proposition 2. ■

**Proposition 3.** *There exists a subset  $\Omega_0$  of  $\Omega$  with  $P(\Omega_0) = 1$ , such that*

$$\Omega_0 \subset \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} |\mathbf{F}_M(\omega, \phi) - \mathbf{f}(\phi)| = 0 \text{ and } \lim_{M \rightarrow \infty} \sup_{\phi \in C} |\mathbb{D}\mathbf{F}_M(\omega, \phi) - \mathbb{D}\mathbf{f}(\phi)| = 0 \right\}.$$

**PROOF:** It is divided into two stages, a) and b).

a) There exists a subset  $\Omega_{01}$  of  $\Omega$  with  $P(\Omega_{01}) = 1$ , such that

$$\Omega_{01} \subset \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} |\mathbf{F}_M(\omega, \phi) - \mathbf{f}(\phi)| = 0 \right\}.$$

For all  $1 \leq p \leq L$ , it is enough to apply Lemma 2 under the following setup

$$c_0(\phi) = 1, \quad c_{j, \tilde{\mathbf{k}}}(\phi) = I_{j, \tilde{\mathbf{k}}}(\phi),$$

for all  $j \geq 1$ ,  $\tilde{\mathbf{k}} \in \mathbb{L}^j$  and  $\phi \in \Lambda$ .

$$W(\mathbf{x}, \mathbf{y}, \phi) = \eta \left( \frac{x_1 - \sum_{p'=1}^L \phi_{p'} x_{p'+1}}{\sigma_\varepsilon}, \frac{y_1 - \sum_{p''=1}^L \phi_{p''} y_{p''+1}}{\sigma_\varepsilon} \right), \quad (46)$$

$$\rho(\mathbf{x}, \mathbf{y}) = 1, \quad (47)$$

with  $x = (x_1, \dots, x_{L+1})'$ ,  $y = (y_1, \dots, y_{L+1})'$  and  $\phi = (\phi_1, \dots, \phi_L)'$ . Finally, set

$$\underline{n}_1 = \underline{t}_p. \quad (48)$$

b) There exists a subset  $\Omega_{02}$  of  $\Omega$  with  $P(\Omega_{02}) = 1$ , such that

$$\Omega_{02} \subset \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} |\mathbb{D}\mathbf{F}_M(\omega, \phi) - \mathbb{D}\mathbf{f}(\phi)| = 0 \right\}.$$

For all  $1 \leq p, q \leq L$ , the following notation will be used

$$\begin{aligned} (a1_{\underline{n}}(\omega, \phi))_{p,q} &= -\frac{1}{\sigma_\varepsilon} \eta_1 \left( R_{\underline{n}}(\omega, \phi), R_{\underline{n}-\underline{t}_p}(\omega, \phi) \right) X_{\underline{n}-\underline{t}_q}(\omega), \\ (a2_{\underline{n}}(\omega, \phi))_{p,q} &= -\frac{1}{\sigma_\varepsilon} \eta_2 \left( R_{\underline{n}}(\omega, \phi), R_{\underline{n}-\underline{t}_p}(\omega, \phi) \right) X_{\underline{n}-\underline{t}_p-\underline{t}_q}(\omega), \\ (a3_{\underline{n},j,\tilde{\mathbf{k}}}(\omega, \phi))_{p,q} &= D_q \left( I_{j,\tilde{\mathbf{k}}}(\phi) \eta \left( R_{\underline{n}}(\omega, \phi), R_{\underline{n}-\underline{t}_p-s(j,\tilde{\mathbf{k}})}(\omega, \phi) \right) \right), \\ (a4_{\underline{n},j,\tilde{\mathbf{k}}}(\omega, \phi))_{p,q} &= -\frac{1}{\sigma_\varepsilon} I_{j,\tilde{\mathbf{k}}}(\phi) \eta_1 \left( R_{\underline{n}}(\omega, \phi), R_{\underline{n}-\underline{t}_p-s(j,\tilde{\mathbf{k}})}(\omega, \phi) \right) X_{\underline{n}-\underline{t}_q}(\omega), \\ (a5_{\underline{n},j,\tilde{\mathbf{k}}}(\omega, \phi))_{p,q} &= -\frac{1}{\sigma_\varepsilon} I_{j,\tilde{\mathbf{k}}}(\phi) \eta_2 \left( R_{\underline{n}}(\omega, \phi), R_{\underline{n}-\underline{t}_p-s(j,\tilde{\mathbf{k}})}(\omega, \phi) \right) X_{\underline{n}-\underline{t}_p-s(j,\tilde{\mathbf{k}})-\underline{t}_q}(\omega), \end{aligned}$$

where  $\omega \in \Omega$ ,  $\underline{n} \in Z^2$ ,  $j$  is a positive integer  $\tilde{\mathbf{k}} \in \mathbb{L}^j$ ,  $\phi \in \Lambda$ . Then

$$\begin{aligned} D_q \left( (\mathbb{D}\mathbf{F}_M(\omega, \phi))_p \right) &= \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} (a1_{\underline{m}}(\omega, \phi))_{p,q} + \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} (a2_{\underline{m}}(\omega, \phi))_{p,q} \\ &+ \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \sum_{j=1}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} (a3_{\underline{m},j,\tilde{\mathbf{k}}}(\omega, \phi))_{p,q} + \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \sum_{j=1}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} (a4_{\underline{m},j,\tilde{\mathbf{k}}}(\omega, \phi))_{p,q} \\ &+ \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \sum_{j=1}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} (a5_{\underline{m},j,\tilde{\mathbf{k}}}(\omega, \phi))_{p,q}. \end{aligned}$$

Thus, to prove b) it is enough to show the following five statements:

b1) There exists a subset  $\Omega_G$  of  $\Omega$  with  $P(\Omega_G) = 1$ , such that for all  $1 \leq p, q \leq L$ ,  $\Omega_G$  is contained in  $\mathbb{G}$ , where

$$\mathbb{G} = \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} \left| \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} (a1_{\underline{m}}(\omega, \phi))_{p,q} - A1_{p,q}(\phi) \right| = 0 \right\}.$$

b2) There exists a subset  $\Omega_H$  of  $\Omega$  with  $P(\Omega_H) = 1$ , such that for each  $\underline{s}, \underline{t} \in T$ ,  $\Omega_H$  is contained in  $\mathbb{H}$ , where

$$\mathbb{H} = \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} \left| \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} (a2_{\underline{m}}(\omega, \phi))_{p,q} - A2_{p,q}(\phi) \right| = 0 \right\}.$$

b3) There exists a subset  $\Omega_A$  of  $\Omega$  with  $P(\Omega_A) = 1$ , such that for each  $\underline{s}, \underline{t} \in T$ ,  $\Omega_A$  is included in  $\mathbb{A}$ , where

$$\mathbb{A} = \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} \left| \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \sum_{j=1}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} (a3_{\underline{m},j,\tilde{\mathbf{k}}}(\omega, \phi))_{p,q} - \sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} A3_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| = 0 \right\}.$$

b4) There exists a subset  $\Omega_B$  of  $\Omega$  with  $P(\Omega_B) = 1$ , such that for each  $\underline{s}, \underline{t} \in T$ ,  $\Omega_B$  is included in  $\mathbb{B}$ , where

$$\mathbb{B} = \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} \left| \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \sum_{j=1}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left( a4_{\underline{n}, j, \tilde{\mathbf{k}}}(\omega, \phi) \right)_{p,q} - \sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} A4_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| = 0 \right\}.$$

b5) There exists a subset  $\Omega_D$  of  $\Omega$  with  $P(\Omega_D) = 1$ , such that for each  $\underline{s}, \underline{t} \in T$ ,  $\Omega_D$  is included in  $\mathbb{D}$ , where

$$\mathbb{D} = \left\{ \omega \in \Omega : \lim_{M \rightarrow \infty} \sup_{\phi \in C} \left| \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \sum_{j=1}^{M-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left( a5_{\underline{n}, j, \tilde{\mathbf{k}}}(\omega, \phi) \right)_{p,q} - \sum_{j=1}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} A5_{p,q}(j, \tilde{\mathbf{k}}, \phi) \right| = 0 \right\}.$$

Proof of b1). If we suppose that (a) of Assumption 2-(ii) holds then we apply Lemma 2 under the following setup

$$c_0(\phi) = 1, \quad c_{j, \tilde{\mathbf{k}}}(\phi) = 0, \quad \forall j \geq 1, \quad \tilde{\mathbf{k}} \in \mathbb{L}^j, \quad \phi \in \Lambda, \quad (49)$$

$$W(\mathbf{x}, \mathbf{y}, \phi) = \eta_1 \left( \frac{x_1 - \sum_{p'=1}^L \phi_{p'} x_{p'+1}}{\sigma_\varepsilon}, \frac{y_1 - \sum_{p''=1}^L \phi_{p''} y_{p''+1}}{\sigma_\varepsilon} \right) x_{q+1}, \quad (50)$$

$$\rho(\mathbf{x}, \mathbf{y}) = x_{q+1}, \quad (51)$$

with  $x = (x_1, \dots, x_{L+1})'$ ,  $y = (y_1, \dots, y_{L+1})'$  and  $\phi = (\phi_1, \dots, \phi_L)'$ . Finally,  $\underline{n}_1$  is considered as in (48). Suppose now that (b) of Assumption 2-(ii), then we replace (51) by

$$\rho(\mathbf{x}, \mathbf{y}) = \left| x_{q+1} \frac{y_1}{\sigma_\varepsilon} \right| + \frac{1}{L} \sum_{p'=1}^L \left| x_{q+1} \frac{y_{p'+1}}{\sigma_\varepsilon} \right|. \quad (52)$$

Proof of b2). Suppose that (a) of Assumption 2-(ii) holds. The proof consists of applying Lemma 2 under the following setup:  $c_0$  and  $c_{j, \tilde{\mathbf{k}}}$  as in (49),

$$W(\mathbf{x}, \mathbf{y}, \phi) = \eta_2 \left( \frac{x_1 - \sum_{p'=1}^L \phi_{p'} x_{p'+1}}{\sigma_\varepsilon}, \frac{y_1 - \sum_{p''=1}^L \phi_{p''} y_{p''+1}}{\sigma_\varepsilon} \right) y_{q+1}, \quad (53)$$

$$\rho(\mathbf{x}, \mathbf{y}) = y_{q+1}, \quad (54)$$

with  $x = (x_1, \dots, x_{L+1})'$ ,  $y = (y_1, \dots, y_{L+1})'$  and  $\phi = (\phi_1, \dots, \phi_L)'$ . Finally,  $\underline{n}_1$  is considered as in (48).

The proof is complete if we suppose that (b) of Assumption 2-(ii) holds, and change (54) by

$$\rho(\mathbf{x}, \mathbf{y}) = \left| y_{q+1} \frac{x_1}{\sigma_\varepsilon} \right| + \frac{1}{L} \sum_{p'=1}^L \left| y_{q+1} \frac{x_{p'+1}}{\sigma_\varepsilon} \right|. \quad (55)$$

Proof of b3). The proof consists in applying Lemma 2 under the following setup:  $c_0(\phi) = 0$ ,  $c_{j, \tilde{\mathbf{k}}}(\phi) = D_q(I_{j, \tilde{\mathbf{k}}}(\phi))$ ,  $j \geq 1$ ,  $\tilde{\mathbf{k}} \in \mathbb{L}^j$  and  $\phi \in \Lambda$ ,  $W(x, y, \phi)$  as in (46),  $\rho$  as in (47) and  $\underline{n}_1$  as in (48).

Proof of b4). Suppose that (a) of Assumption 2-(ii) holds. Then we apply Lemma 2 for

$$c_0(\phi) = 0, \quad c_{j, \tilde{\mathbf{k}}}(\phi) = -I_{j, \tilde{\mathbf{k}}}(\phi), \quad (56)$$

for all  $j \geq 1$ ,  $\tilde{\mathbf{k}} \in \mathbb{L}^j$  and  $\phi \in \Lambda$ ,  $W(x, y, \phi)$  as in (50),  $\rho$  as in (51) and  $\underline{n}_1$  as in (48). Suppose that (b) of Assumption 2-(ii) holds then consider  $\rho$  as in (52).

Proof of b5). Suppose (a) of Assumption 2-(ii) holds. Similarly, we apply Lemma 2 under the following conditions:  $c_0$  and  $c_{j, \tilde{\mathbf{k}}}$  as in (56),  $W(x, y, \phi)$  as in (53),  $\rho$  as in (54). Consider  $\underline{n}_1$  as in (48). To complete the proof, under (b) of Assumption 2-(ii), consider  $\rho$  as in (55). This completes the proof of Proposition 3. ■

### PROOF OF THEOREM 1

Let  $\Omega_0$  be as in Proposition 3. Let  $\omega \in \Omega_0$ . We consider the Zeros Lemma (Lemma 3) under the following conditions:  $U = C^\circ$  (the interior of  $C$ ),  $\lambda_0 = \phi_0$ , for all  $M$ ,  $\mathbf{q}_M$  is the function  $\phi \mapsto \mathbf{F}_M(\omega, \phi)$ ,  $\mathbf{q}$  is the function  $\mathbf{f}$  defined in (17). Then, 1) in the Zeros Lemma follows from (18). 2) in the Zeros Lemma follows from 2) in Proposition 2. Finally, 3) in

Zeros Lemma follows from Proposition 3. Hence  $\omega \in \Omega'''$ . Setting  $\Omega'' = \Omega_0$ , the proof of the Theorem 1 is completed.  $\blacksquare$

## 6.2 Proofs of the Asymptotic Normality of the RA Estimator

We will start introducing some additional notation.

Consider  $1 \leq M, N \leq \infty$  integers,  $\underline{n}, \underline{m} \in \mathbb{Z}^2$ ,  $\omega \in \Omega$ ,  $\phi \in \Lambda$  and  $1 \leq p \leq L$ . Denoting

$$\begin{aligned} \left( \mathbf{Y}_{\underline{n}, N}^n(\omega, \phi) \right)_p &= \sum_{j=N}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j, \tilde{\mathbf{k}}}(\phi) \eta \left( R_{\underline{n}}(\omega, \phi), R_{\underline{n} - \underline{t}_p - s(j, \tilde{\mathbf{k}})}(\omega, \phi) \right), \\ \mathbf{Y}_{M, N}(\omega, \phi) &= \frac{1}{\#(W_M)} \sum_{\underline{m} \in W_M} \mathbf{Y}_{\underline{m}, N}^n(\omega, \phi), \end{aligned}$$

one has

$$\sqrt{\#(W_M)} \mathbf{G}_{M, \infty}(\phi) = \sqrt{\#(W_M)} \mathbf{G}_{M, N}(\phi) + \sqrt{\#(W_M)} \mathbf{Y}_{M, N}(\phi). \quad (57)$$

In particular,

$$\sqrt{\#(W_M)} \mathbf{G}_{M, \infty}(\hat{\phi}_M^{RA}) = \sqrt{\#(W_M)} \mathbf{F}_M(\hat{\phi}_M^{RA}) + \sqrt{\#(W_M)} \mathbf{Y}_{M, M}(\hat{\phi}_M^{RA}). \quad (58)$$

For each  $N \geq 2$  integer, let  $\mathbf{G}_N$  be the  $L \times L$  matrix given by

$$\mathbf{G}_N = \mathbf{I}_N(\phi_0) \cdot E \left( \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right)^2 \right),$$

where  $\mathbf{I}_N(\phi_0)$  is the  $L \times L$  matrix given by

$$\mathbf{I}_N(\phi_0)_{p, p} = 1 + \sum_{(j, \tilde{\mathbf{k}}, l, \tilde{\mathbf{h}}) \in (T-T)_2^N(\underline{0})} I_{j, \tilde{\mathbf{k}}}(\phi_0) I_{l, \tilde{\mathbf{h}}}(\phi_0),$$

for all  $1 \leq p \leq L$ ,

$$\begin{aligned} \mathbf{I}_N(\phi_0)_{p, q} &= \sum_{(j, \tilde{\mathbf{k}}) \in (T-T)_1^N(\underline{t}_p - \underline{t}_q)} I_{j, \tilde{\mathbf{k}}}(\phi_0) + \sum_{(j, \tilde{\mathbf{k}}) \in (T-T)_1^N(\underline{t}_q - \underline{t}_p)} I_{j, \tilde{\mathbf{k}}}(\phi_0) \\ &+ \sum_{(j, \tilde{\mathbf{k}}, l, \tilde{\mathbf{h}}) \in (T-T)_2^N(\underline{t}_p - \underline{t}_q)} I_{j, \tilde{\mathbf{k}}}(\phi_0) \cdot I_{l, \tilde{\mathbf{h}}}(\phi_0), \end{aligned}$$

for  $p \neq q$  with  $1 \leq p, q \leq L$ , where

$$\begin{aligned} (T-T)_1^N(\underline{v}) &= \left\{ (j, \tilde{\mathbf{k}}) : N-1 \geq j \geq 1, \tilde{\mathbf{k}} \in \mathbb{L}^j \text{ and } s(j, \tilde{\mathbf{k}}) = \underline{v} \right\}, \\ (T-T)_2^N(\underline{v}) &= \left\{ (j, \tilde{\mathbf{k}}, l, \tilde{\mathbf{h}}) : N-1 \geq j, l \geq 1, \tilde{\mathbf{k}} \in \mathbb{L}^j, \tilde{\mathbf{h}} \in \mathbb{L}^l \text{ and } s(j, \tilde{\mathbf{k}}) - s(l, \tilde{\mathbf{h}}) = \underline{v} \right\}, \end{aligned}$$

with  $\underline{v} \in (T-T)$  and  $(T-T)$  as in (12).

Using the mean value theorem we have that

$$\sqrt{\#(W_M)} \mathbf{G}_{M, \infty}(\hat{\phi}_M^{RA}) = \sqrt{\#(W_M)} \mathbf{G}_{M, \infty}(\phi_0) + \mathbb{D} \mathbf{G}_{M, \infty}(\hat{\phi}_M^*) \cdot \sqrt{\#(W_M)} (\hat{\phi}_M^{RA} - \phi_0), \quad (59)$$

where, for each  $M$ ,  $\hat{\phi}_M^*$  is a random vector belonging to the segment in  $\mathbb{R}^L$  that connects  $\hat{\phi}_M^{RA}$  and  $\phi_0$ .

The following lemmas and propositions are necessary to prove Theorem 2, which is proved at the end of this subsection.

**Lemma 4.** *Let  $1 \leq M, N$  integers. For all  $1 \leq p \leq L$  we have that*

$$\lim_{N \rightarrow \infty} \limsup_{M \rightarrow \infty} \left( \sqrt{\#(W_M)} \mathbf{Y}_{M, N}(\phi_0) \right) = 0,$$

in probability.

**PROOF:** Let  $M \geq 1$  be an integer and  $\delta > 0$ . By Chebyshev's inequality

$$\begin{aligned} P\left(\left|\left(\sqrt{\#(W_M)}\mathbf{Y}_{M,N}(\phi_0)\right)_p\right|\geq\delta\right) &= P\left(\left|\left(\frac{1}{\sqrt{\#(W_M)}}\sum_{\underline{m}\in W_M}\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p\right|\geq\delta\right) \\ &\leq\frac{1}{\delta^2\#(W_M)}\text{Var}\left(\sum_{\underline{m}\in W_M}\left(\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p\right). \end{aligned}$$

Considering that,

$$\begin{aligned} \left(\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p &= \sum_{j=N}^{\infty}\sum_{\tilde{\mathbf{k}}\in T^j}I_{j,\tilde{\mathbf{k}}}(\phi_0)\cdot\eta\left(R_{\underline{m}}(\phi_0),R_{\underline{m}-\underline{t}_p-s(j,\tilde{\mathbf{k}})}(\phi_0)\right) \\ &= \sum_{j=N}^{\infty}\sum_{\tilde{\mathbf{k}}\in L^j}I_{j,\tilde{\mathbf{k}}}(\phi_0)\cdot\eta\left(\frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon},\frac{\varepsilon_{\underline{m}-\underline{t}_p-s(j,\tilde{\mathbf{k}})}}{\sigma_\varepsilon}\right), \end{aligned}$$

we have that

$$E\left(\left(\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p\right)=0.$$

and, therefore,

$$\begin{aligned} \text{Var}\left(\sum_{\underline{m}\in W_M}\left(\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p\right) &= E\left(\left(\sum_{\underline{m}\in W_M}\left(\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p\right)^2\right) \\ &= \sum_{\underline{m}\in W_M}\sum_{\underline{m}'\in W_M}E\left(\left(\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p\left(\mathbf{r}_{\underline{m}',N}^\eta(\phi_0)\right)_p\right). \end{aligned} \quad (60)$$

If  $\underline{m} \neq \underline{m}'$ , by the Assumption 3, it follows that

$$\begin{aligned} &E\left(\left(\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p\left(\mathbf{r}_{\underline{m}',N}^\eta(\phi_0)\right)_p\right) = \\ &= \sum_{j_1=N}^{\infty}\sum_{\tilde{\mathbf{k}}_1\in L^{j_1}}\sum_{j_2=N}^{\infty}\sum_{\tilde{\mathbf{k}}_2\in L^{j_2}}I_{j_1,\tilde{\mathbf{k}}_1}(\phi_0)I_{j_2,\tilde{\mathbf{k}}_2}(\phi_0)E\left(\eta\left(\frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon},\frac{\varepsilon_{\underline{m}-\underline{t}_p-s(j_1,\tilde{\mathbf{k}}_1)}}{\sigma_\varepsilon}\right)\eta\left(\frac{\varepsilon_{\underline{m}'}}{\sigma_\varepsilon},\frac{\varepsilon_{\underline{m}'-\underline{t}_p-s(j_2,\tilde{\mathbf{k}}_2)}}{\sigma_\varepsilon}\right)\right) \\ &= 0. \end{aligned}$$

Hence, for any  $\underline{m}$

$$\begin{aligned} &E\left(\left(\left(\mathbf{r}_{\underline{m},N}^\eta(\phi_0)\right)_p\right)^2\right) = \\ &= \sum_{j_1=N}^{\infty}\sum_{\tilde{\mathbf{k}}_1\in L^{j_1}}\sum_{j_2=N}^{\infty}\sum_{\tilde{\mathbf{k}}_2\in L^{j_2}}I_{j_1,\tilde{\mathbf{k}}_1}(\phi_0)I_{j_2,\tilde{\mathbf{k}}_2}(\phi_0)E\left(\eta\left(\frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon},\frac{\varepsilon_{\underline{m}-\underline{t}_p-s(j_1,\tilde{\mathbf{k}}_1)}}{\sigma_\varepsilon}\right)\eta\left(\frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon},\frac{\varepsilon_{\underline{m}-\underline{t}_p-s(j_2,\tilde{\mathbf{k}}_2)}}{\sigma_\varepsilon}\right)\right) \\ &= \sum_{(j_1,\tilde{\mathbf{k}}_1,j_2,\tilde{\mathbf{k}}_2)\in(T-T)_2^{N,\infty}(0)}I_{j_1,\tilde{\mathbf{k}}_1}(\phi_0)I_{j_2,\tilde{\mathbf{k}}_2}(\phi_0)E\left(\eta\left(\frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon},\frac{\varepsilon_{\underline{m}-\underline{t}_p-s(j_1,\tilde{\mathbf{k}}_1)}}{\sigma_\varepsilon}\right)\eta\left(\frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon},\frac{\varepsilon_{\underline{m}-\underline{t}_p-s(j_2,\tilde{\mathbf{k}}_2)}}{\sigma_\varepsilon}\right)\right), \end{aligned}$$

where

$$(T-T)_2^{N,\infty}(0)=\left\{(j_1,\tilde{\mathbf{k}}_1,j_2,\tilde{\mathbf{k}}_2):j_1,j_2\geq N,\tilde{\mathbf{k}}_1\in L^{j_1},\tilde{\mathbf{k}}_2\in L^{j_2},s(j_1,\tilde{\mathbf{k}}_1)=s(j_2,\tilde{\mathbf{k}}_2)\right\}.$$

So,

$$\begin{aligned} & \text{Var} \left( \sum_{\underline{m} \in W_M} \left( \mathbf{r}_{\underline{m}, N}^\eta(\phi_0) \right)_p \right) \\ &= \#(W_M) \sum_{(j_1, \tilde{\mathbf{k}}_1, j_2, \tilde{\mathbf{k}}_2) \in (T-T)_2^{N, \infty}(0)} I_{j_1, \tilde{\mathbf{k}}_1}(\phi_0) I_{j_2, \tilde{\mathbf{k}}_2}(\phi_0) E \left( \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_{\varepsilon'}} \right)^2 \right), \end{aligned}$$

where  $\varepsilon$  and  $\varepsilon'$  are independent random variables with distribution  $F_\varepsilon$ .

Using this expression for the variance and taking lim sup in the Chebyshev inequality, we have that

$$\begin{aligned} & \limsup_{M \rightarrow \infty} P \left( \left| \left( \frac{1}{\sqrt{\#(W_M)}} \sum_{\underline{m} \in W_M} \mathbf{r}_{\underline{m}, N}^\eta(\phi_0) \right)_p \right| \geq \delta \right) \\ & \leq \frac{1}{\delta^2} \sum_{(j_1, \tilde{\mathbf{k}}_1, j_2, \tilde{\mathbf{k}}_2) \in (T-T)_2^{N, \infty}(0)} I_{j_1, \tilde{\mathbf{k}}_1}(\phi_0) I_{j_2, \tilde{\mathbf{k}}_2}(\phi_0) E \left( \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_{\varepsilon'}} \right)^2 \right). \end{aligned}$$

Notice that  $(T-T)_2^{N, \infty}(0) \subset (T-T)_2(0)$ ,  $((T-T)_2^{N, \infty}(0))_N$  is a decreasing sequence of subsets of  $(T-T)_2(0)$  such that  $\bigcap_{N \geq 1} (T-T)_2^{N, \infty}(0) = \emptyset$  and that  $|\sum_{(j, \tilde{\mathbf{k}}, l, \tilde{\mathbf{h}}) \in (T-T)_2(0)} I_{j, \tilde{\mathbf{k}}}(\phi_0) \cdot I_{l, \tilde{\mathbf{h}}}(\phi_0)| < \infty$ . Then, one has that  $\sum_{(j_1, \tilde{\mathbf{k}}_1, j_2, \tilde{\mathbf{k}}_2) \in (T-T)_2^{N, \infty}(0)} I_{j_1, \tilde{\mathbf{k}}_1}(\phi_0) I_{j_2, \tilde{\mathbf{k}}_2}(\phi_0) \xrightarrow{N \rightarrow \infty} 0$ . Thus,

$$\lim_{N \rightarrow \infty} \limsup_{M \rightarrow \infty} P \left( \left| \left( \frac{1}{\sqrt{\#(W_M)}} \sum_{\underline{m} \in W_M} \mathbf{r}_{\underline{m}, N}^\eta(\phi_0) \right)_p \right| \geq \delta \right) = 0. \blacksquare$$

The following Central Limit Theorem for Random Fields is needed. The proof can be found in Guyon (1993), page 99.

**Lemma 5** (Central Limit Theorem for Random Fields). *Let  $X = \{X_i : i \in \mathbb{Z}^d\}$  be a real valued random field such that  $E(X_i) = 0$ , for all  $i \in \mathbb{Z}^d$ . For each  $\mathbb{S} \subset \mathbb{Z}^d$ , let  $F(X, \mathbb{S})$  be the  $\sigma$ -algebra generated by  $\{X_i^{-1}(C) : C \text{ is a Borel set}, i \in \mathbb{S}\}$ ; for each  $k, l \in \mathbb{N} \cup \{\infty\}$  and  $n \in \mathbb{N}$  let*

$$\alpha_{k, l}(n) = \sup \{ |P(A \cap B) - P(A)P(B)| : A, B \in \mathcal{C}_{k, l}(n) \}$$

where

$$\mathcal{C}_{k, l}(n) = \{A \in F(X, \mathbb{S}_1), B \in F(X, \mathbb{S}_2) : \#\mathbb{S}_1 \leq k, \#\mathbb{S}_2 \leq l, \text{dist}(\mathbb{S}_1, \mathbb{S}_2) \geq n\}.$$

Let  $(D_n)_{n \geq 1}$  be a decreasing sequence of finite subsets of  $\mathbb{Z}^d$  and  $S_n = \sum_{i \in D_n} X_i$  with variance  $\sigma_n^2$ . If the following two conditions hold

(i)  $\sum_{m \geq 1} m^{d-1} \alpha_{k, l}(m) < \infty$  if  $k + l \leq 4$  and  $\alpha_{1, \infty}(m) = o(m^{-d})$ ;

(ii) there exists  $\delta > 0$  such that  $\sup_i \|X_i\|_{\delta+2} < \infty$  and  $\sum_{m \geq 1} m^{d-1} (\alpha_{1, 1}(m))^{\frac{\delta}{\delta+2}} < \infty$ , then

$$\limsup_n \frac{1}{\#(D_n)} \sum_{i, j \in D_n} |\text{Cov}(X_i, X_j)| < \infty.$$

If, further, we assume that

(iii)

$$\liminf_{n \rightarrow \infty} \frac{1}{\#(D_n)} \sigma_n^2 > 0,$$

then

$$\sigma_n^{-1} S_n \xrightarrow{D} N(0, 1).$$



If, for each  $i \in \mathbb{Z}^d$ ,  $X_i$  takes values in  $\mathbb{R}^p$  and if (iii) is replaced by

$$\liminf_{n \rightarrow \infty} \frac{1}{\#(D_n)} \Sigma_n \geq I_0 > 0,$$

with  $I_0$  a symmetric and positive definite matrix, then

$$\Sigma_n^{-1/2} S_n \xrightarrow{D} N(\mathbf{0}, \mathbf{I}_p),$$

where  $\Sigma_n$  is the covariance matrix of  $S_n$  and  $\mathbf{I}_p$  is the identity matrix of order  $p$ .

**Lemma 6.** *There exists  $N_0$  such that, for each  $N \geq N_0$ ,  $\sqrt{\#(W_M)} \mathbf{G}_{M,N}(\phi_0) \xrightarrow{D} N(0, \mathbf{G}_N)$ , as  $M \rightarrow \infty$ .*

**PROOF:** Let  $N \geq 2$ . For each  $M \geq 1$ , let us define

$$\mathbf{S}_{M,N} = \#(W_M) \mathbf{G}_{M,N}(\phi_0) = \sum_{\underline{m} \in W_M} \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0). \quad (61)$$

For each  $\underline{m} \in \mathbb{Z}^2$ , let

$$\mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) = \left( \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_1, \dots, \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_L \right)',$$

with

$$\begin{aligned} \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_p &= \eta \left( R_{\underline{m}}(\phi_0), R_{\underline{m}-\underline{t}_p}(\phi_0) \right) + \sum_{j=1}^{N-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j,\tilde{\mathbf{k}}}(\phi_0) \cdot \eta \left( R_{\underline{m}}(\phi_0), R_{\underline{m}-\underline{t}_p-s(j,\tilde{\mathbf{k}})}(\phi_0) \right) \\ &= \eta \left( \frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon}, \frac{\varepsilon_{\underline{m}-\underline{t}_p}}{\sigma_\varepsilon} \right) + \sum_{j=1}^{N-1} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} I_{j,\tilde{\mathbf{k}}}(\phi_0) \cdot \eta \left( \frac{\varepsilon_{\underline{m}}}{\sigma_\varepsilon}, \frac{\varepsilon_{\underline{m}-\underline{t}_p-s(j,\tilde{\mathbf{k}})}}{\sigma_\varepsilon} \right). \end{aligned}$$

We will apply Lemma 5 to the  $\mathbb{R}^L$ -valued random field  $(\mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0))_{\underline{m} \in \mathbb{Z}^2}$ . Since  $(\varepsilon(\underline{m}))_{\underline{m} \in \mathbb{Z}^2}$  is a white noise, it is straightforward to verify that  $(\mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0))_{\underline{m} \in \mathbb{Z}^2}$  satisfies the conditions (i) and (ii) of this lemma.

Now, let us compute the covariance matrix of  $\mathbf{S}_{M,N}$ ,  $\Sigma(\mathbf{S}_{M,N})$ . Since  $E(\mathbf{S}_{M,N}) = \mathbf{0}$ , then  $\Sigma(\mathbf{S}_{M,N})$  is the  $L \times L$  matrix given by

$$\left[ \sum_{\underline{m} \in W_M} \sum_{\underline{m}' \in W_M} E \left( \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_p \left( \mathbf{\Gamma}_{\underline{m}',N}^\eta(\phi_0) \right)_q \right) \right]_{1 \leq p, q \leq L}.$$

By the Assumption 3, if  $\underline{m} \neq \underline{m}'$ , then  $E \left( \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_p \left( \mathbf{\Gamma}_{\underline{m}',N}^\eta(\phi_0) \right)_q \right) = 0$ , for any  $p$  and  $q$  such that  $1 \leq p, q \leq L$ .

Hence

$$\Sigma(\mathbf{S}_{M,N}) = \left[ \sum_{\underline{m} \in W_M} E \left( \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_p \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_q \right) \right]_{1 \leq p, q \leq L}. \quad (62)$$

Using Assumption 3, a simple calculation shows that

$$\left[ E \left( \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_p \left( \mathbf{\Gamma}_{\underline{m},N}^\eta(\phi_0) \right)_q \right) \right]_{1 \leq p, q \leq L} = \mathbf{G}_N,$$

for all  $\underline{m} \in W_M$ . Therefore, by (62)

$$\frac{1}{\#(W_M)} \Sigma(\mathbf{S}_{M,N}) = \mathbf{G}_N. \quad (63)$$

Now,  $\mathbf{I}_N(\phi_0) \rightarrow_{N \rightarrow \infty} \mathbf{I}_\infty(\phi_0)$  and  $\mathbf{I}_\infty(\phi_0)$  is positive definite. Then, there exists  $N_0$  such that for each  $N \geq N_0$   $\mathbf{I}_N(\phi_0)$  is positive definite. Hence, by (63), for each  $N \geq N_0$  (iii) of Lemma 5 is satisfied with  $D_M = W_M$ ; hence

$$(\Sigma(\mathbf{S}_{M,N}))^{-1/2} \mathbf{S}_{M,N} \rightarrow_{M \rightarrow \infty}^D N(0, \mathbf{I}_L), \quad (64)$$

where  $\mathbf{I}_L$  is the  $L \times L$  identity matrix. Therefore by (63) and (61),

$$(\Sigma(\mathbf{S}_{M,N}))^{-1/2} \mathbf{S}_{M,N} = \frac{1}{\sqrt{\#(W_M)}} (\mathbf{G}_N)^{-1/2} \#(W_M) \mathbf{G}_{M,N}(\phi_0) = (\mathbf{G}_N)^{-1/2} \sqrt{\#(W_M)} \mathbf{G}_{M,N}(\phi_0),$$

for all  $M$  and  $N \geq N_0$ . Consequently, by (64) the proof is completed. ■

**Lemma 7.**

$$\mathbb{D}\mathbf{G}_{M,\infty}(\widehat{\phi}_M^*) \xrightarrow{P} \mathbb{D}\mathbf{f}(\phi_0), \text{ as } M \rightarrow \infty,$$

where  $\widehat{\phi}_M^*$  is as in (59).

**PROOF:** Let  $\delta$  and  $\delta'$  be any positive numbers. From the proof of Proposition 3, we see that

$$\sup_{\phi \in \mathcal{C}} |\mathbb{D}\mathbf{G}_{M,\infty}(\phi) - \mathbb{D}\mathbf{f}(\phi)| \xrightarrow{P} \mathbf{0}, \text{ as } M \rightarrow \infty.$$

Then, there exists  $M_1$  such that

$$M \geq M_1 \implies P(\Omega_M(\delta/2)) \geq 1 - \delta'/2,$$

where

$$\Omega_M(\delta/2) = \left\{ \omega \in \Omega : \sup_{\phi \in \mathcal{C}} |\mathbb{D}\mathbf{G}_{M,\infty}(\omega, \phi) - \mathbb{D}\mathbf{f}(\phi)| < \delta/2 \right\}.$$

Hence, for each  $\omega \in \Omega_M(\delta/2)$  and  $M \geq M_1$ ,

$$\left| \mathbb{D}\mathbf{G}_{M,\infty}(\omega, \widehat{\phi}_M^*(\omega)) - \mathbb{D}\mathbf{f}(\widehat{\phi}_M^*(\omega)) \right| < \delta/2. \quad (65)$$

Because of the continuity of the function  $\phi \mapsto \mathbb{D}\mathbf{f}(\phi)$ , there exists  $\delta'' > 0$ , such that

$$|\phi - \phi_0| < \delta'' \implies |\mathbb{D}\mathbf{f}(\phi) - \mathbb{D}\mathbf{f}(\phi_0)| < \delta/2. \quad (66)$$

By (16) and (6.2), it follows that there exists  $M_2$  such that

$$M \geq M_2 \implies P(\Omega'_M(\delta'')) \geq 1 - \delta'/2,$$

with

$$\Omega'_M(\delta'') = \left\{ \omega \in \Omega : \left| \widehat{\phi}_M^*(\omega) - \phi_0 \right| < \delta'' \right\}.$$

Consequently

$$\Omega'_M(\delta'') \subseteq \left\{ \omega \in \Omega : \left| \mathbb{D}\mathbf{f}(\widehat{\phi}_M^*(\omega)) - \mathbb{D}\mathbf{f}(\phi_0) \right| < \delta/2 \right\}. \quad (67)$$

Then, for  $M \geq \max(M_1, M_2)$ , using (65) and (67) we have that

$$\left| \mathbb{D}\mathbf{G}_{M,\infty}(\omega, \widehat{\phi}_M^*(\omega)) - \mathbb{D}\mathbf{f}(\phi_0) \right| < \delta, \text{ if } \omega \in \Omega_M(\delta/2) \cap \Omega'_M(\delta'').$$

Thus

$$P\left(\left\{\omega \in \Omega : \left| \mathbb{D}\mathbf{G}_{M,\infty}(\omega, \widehat{\phi}_M^*(\omega)) - \mathbb{D}\mathbf{f}(\phi_0) \right| < \delta\right\}\right) \geq 1 - \delta'. \blacksquare$$

**Proposition 4.**

$$\sqrt{\#(W_M)} \mathbf{G}_{M,\infty}(\widehat{\phi}_M^{RA}) \xrightarrow{P} \mathbf{0}, \text{ as } M \rightarrow \infty.$$

**PROOF:** Since (16) and (58) it is enough to show that for each  $1 \leq p \leq L$ ,

$$\left(\sqrt{\#(W_M)} \mathbf{Y}_{M,M}(\widehat{\phi}_M^{RA})\right)_p \xrightarrow{P} 0, \text{ as } M \rightarrow \infty.$$

Now, for each  $\omega \in \Omega$  and  $\phi \in \Lambda$

$$\begin{aligned}
& \left| \left( \sqrt{\#(W_M)} \mathbf{Y}_{M,M}(\omega, \phi) \right)_p \right| \\
& \leq \frac{1}{\sqrt{\#(W_M)}} \sum_{j=M}^{\infty} \sum_{\tilde{\mathbf{k}} \in \mathbb{L}^j} \left| I_{j, \tilde{\mathbf{k}}}(\phi) \right| \sum_{m \in W_M} \left| \eta \left( R_m(\omega, \phi), R_{\underline{m} - \underline{t}_p - s(j, \tilde{\mathbf{k}})}(\omega, \phi) \right) \right| \\
& \leq \frac{1}{\sqrt{\#(W_M)}} \sum_{j=M}^{\infty} b^j \#(W_M) K_\eta \\
& = K_\eta b \sqrt{\#(W_M)} \sum_{j=M}^{\infty} b^{j-1} = K_\eta b \frac{\sqrt{\#(W_M)}}{M} \sum_{j=M}^{\infty} M b^{j-1} \\
& \leq K_\eta b \frac{\sqrt{\#(W_M)}}{M} \sum_{j=M}^{\infty} j b^{j-1} \rightarrow 0, \text{ as } M \rightarrow \infty,
\end{aligned}$$

since  $\frac{\sqrt{\#(W_M)}}{M} = \frac{\sqrt{(M+1)^2 - 1}}{M} = \sqrt{1 + \frac{2}{M}}$ , and  $\sum_{j=1}^{\infty} j b^{j-1}$  is the derivative of  $b \mapsto \frac{1}{1-b}$  with  $0 < b < 1$ . ■

**Lemma 8** (Theorem 3.2 in Billingsley (1999), page 27)). *Suppose that for each  $N, M \in \mathbb{N}$ ,  $(X_{N,M}, X_M)$  are random elements of  $S \times S$  where  $(S, \rho)$  is a metric space. If for all  $N$*

$$\begin{aligned}
& X_{N,M} \xrightarrow{D} Z_N, \text{ as } M \rightarrow \infty \\
& Z_N \xrightarrow{D} X, \text{ as } N \rightarrow \infty \\
& \text{for all } \varepsilon > 0: \lim_{N \rightarrow \infty} \limsup_{M \rightarrow \infty} P[\rho(X_{N,M}, X_M) \geq \varepsilon] = 0,
\end{aligned}$$

then  $X_M \xrightarrow{D} X$  as  $M \rightarrow \infty$ .

**Proposition 5.** *Let  $\mathbf{G}$  be the  $L \times L$  matrix given by*

$$\mathbf{G} = \mathbf{I}_\infty(\phi_0) \cdot E \left( \left( \eta \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_{\varepsilon'}} \right) \right)^2 \right), \quad (68)$$

where  $\varepsilon$  and  $\varepsilon'$  are independent random variables with distribution  $F_\varepsilon$ . Then  $\sqrt{\#(W_M)} \mathbf{G}_{M,\infty}(\phi_0) \xrightarrow{D} N(\mathbf{0}, \mathbf{G})$ , as  $M \rightarrow \infty$ .

**PROOF:** Let  $N_0$  be as in Lemma 6 and for each  $N \geq N_0$ , let

$$\begin{aligned}
\mathbf{X}_{N,M} &= \sqrt{\#(W_M)} \mathbf{G}_{M,N}(\phi_0), \\
\mathbf{X}_M &= \sqrt{\#(W_M)} \mathbf{G}_{M,\infty}(\phi_0).
\end{aligned}$$

For each  $N \geq N_0$ , let  $Z_N$  be a random vector variable with distribution  $N(\mathbf{0}, \mathbf{G}_N)$  and  $X$  a random vector variable with distribution  $N(\mathbf{0}, \mathbf{G})$ . By Lemma 6, for each  $N \geq N_0$ ,

$$\mathbf{X}_{N,M} \xrightarrow{D} \mathbf{Z}_N, \text{ as } M \rightarrow \infty.$$

Since  $\mathbf{G}_N \rightarrow \mathbf{G}$ , as  $N \rightarrow \infty$ , then

$$\mathbf{Z}_N \xrightarrow{D} \mathbf{X}, \text{ as } N \rightarrow \infty.$$

By (57),  $\mathbf{X}_M - \mathbf{X}_{N,M} = \sqrt{\#(W_M)} \mathbf{Y}_{M,N}(\phi)$  and, by Lemma 4,

$$\lim_{N \rightarrow \infty} \limsup_{M \rightarrow \infty} \left( \sqrt{\#(W_M)} \mathbf{Y}_{M,N}(\phi_0) \right) = 0.$$

So, the assumptions of Lemma 8 are satisfied; hence  $X_M \xrightarrow{D} X$  as  $M \rightarrow \infty$ , that is,

$$\sqrt{\#(W_M)} \mathbf{G}_{M,\infty}(\phi_0) \xrightarrow{D} N(\mathbf{0}, \mathbf{G}), \text{ as } M \rightarrow \infty. \blacksquare$$

**PROOF OF THEOREM 2:** By Lemma 7, (59), Proposition 4, Proposition 5, and Slutsky's Theorem we have that

$$\sqrt{\#(W_M)} \left( \hat{\phi}_M^{RA} - \phi_0 \right) \xrightarrow{D} N \left( 0, \mathbb{Df}(\phi_0)^{-1} \mathbf{G} \mathbb{Df}(\phi_0)^{-1} \right), \text{ as } M \rightarrow \infty. \quad (69)$$

Now, notice (see (68)) that  $\mathbb{Df}(\phi_0) = -\frac{1}{\sigma_\varepsilon} E \left( \eta_1 \left( \frac{\varepsilon}{\sigma_\varepsilon}, \frac{\varepsilon'}{\sigma_\varepsilon} \right) \cdot \varepsilon' \right) \cdot \mathbf{I}_\infty(\phi_0)$ , where  $\varepsilon$  and  $\varepsilon'$  are independent random variables with distribution  $F_\varepsilon$ . From the results proved in the previous Section and (69) the proof is completed. ■

## 7 CONCLUDING REMARKS

The following comments give a brief summary of the results that we obtained in this paper.

- ✓ Under mild regularity conditions, we established the asymptotic normality and consistency of a class of robust estimators (the RA estimators) for the parameter  $\phi$  of a two dimensional autoregressive unilateral process.
- ✓ The results we proved extend the asymptotic theory of the RA estimators, available only for one-dimensional time series (see Bustos et al. (1984)).
- ✓ Although in the literature several reasonable classes of estimators for the parameter  $\phi$  have been proposed, such as M and GM estimators, their asymptotic behavior are still open problems. Moreover, the advantage of the RA estimator over the other classes is that they are less sensitive to the presence of additive outliers (see Ojeda et al. (2002)).

The following proposals outline some directions for future work:

- ✓ The extension of the results proved in this paper to the model with colored noise (that is, with non null autocorrelation), instead of white noise  $\tilde{\varepsilon} = (\varepsilon_m)_{m \in \mathbb{Z}^2}$ , would be of importance in signal and image processing.
- ✓ In the case of causal AR-2D processes of order two or three, algorithms for the computation of the RA estimators have been proposed (see Ojeda (1999), Ojeda et al. (2002) and Vallejos et al. (2006)). It would be important to develop efficient algorithms when the order of the process is greater than three.
- ✓ To analyze the behaviour of the RA estimator in combination with image restoration techniques.
- ✓ To study properties of RA estimator, in particular, and robust estimators, in general, as alternatives to the least squares estimators under not causal and semi causal AR-2D.

## REFERENCES

- Allende, H., Galbiati, J. and Vallejos, R. (1998). Digital image restoration using autoregressive time series type models. *ESA*. 434. 53–59.
- Allende, H., Galbiati, J. and Vallejos, R. (2001). Robust image modeling on image processing. *Pattern Recognition Letters*. 22. 1219–1231.
- Allende, H. and Pizarro, L. (2003). Robust estimation of roughness parameter in SAR amplitude images. *Progress in Pattern Recognition, Speech and Image Analysis*. 2905. 129–136.
- Aysal, T. C. and Barner, K. E. (2006). Quadratic weighted median filters for edge enhancement of noisy images. *IEEE Transactions on Image Processing*. 15. 3294–3310.
- Basu, S. and Reinsel, G. (1993). Properties of the spatial unilateral first-order ARMA model. *Adv. Appl. Prob.* 25. 631–648.
- Ben Hamza, A. and Krim, H. (2001). Image denoising: A nonlinear robust statistical approach. *IEEE Transactions on Signal Processing*. 49. 3045–3054.
- Black, M. J. and Rangarajan, A. (1996). On the unification of line processes, outlier rejection, and robust statistics with applications in early vision. *International Journal of Computer Vision*. 19. 57–91.

- Black, M. J., Sapiro, G., Marimont, D. and Heeger, D. (1997). Robust anisotropic diffusion: Connections between robust statistics, line processing, and anisotropic diffusion. *Scale-Space Theory in Computer Vision*. 1252. 323–326.
- Billingsley, P. (1999). *Convergence of probability measures*. Second Edition. Wiley. New York.
- Bouzouba, K. and Radouane, L. (2000). Image identification and estimation using the maximum entropy principle. *Pattern Recognition Letters*. 21. 691–700.
- Brandle, N., Bischof, H. and Lapp, H. (2003). Robust DNA microarray image analysis. *Machine Vision and Applications*. 15. 11–28.
- Brunelli, R. and Messelodi, S. (1995). Robust estimation of correlation with applications to computer vision. *Pattern Recognition*. 28. 833–841.
- Bustos O., Fraiman, R. and Yohai, V. (1984). Asymptotic behavior of the estimates based on residual autocovariances for ARMA models. In *Robust and nonlinear time series*. Franke, Hardle and Martin (eds). *Lectures Notes in Statistics*, 29, 26–49, Springer Verlag, New York.
- Bustos, O. H., Lucini, M. M. and Frery, A. C. (2002). M-estimators of roughness and scale for GA0-modelled SAR imagery. *Eurasip Journal on Applied Signal Processing*. 2002. 105–114.
- Bustos, O. and Yohai, V. (1986). Robust estimates for ARMA models. *Journal of the American Statistical Society*. 81 155–168.
- Butler, N. (1998). A frequency domain comparison of estimation methods for gaussian fields. *Communications in Statistics-Theory and Methods*. 27. 2325–2342.
- Chen, J. H., Chen, C. S. and Chen, Y. S. (2003). Fast algorithm for robust template matching with M-estimators. *IEEE Transactions on Signal Processing*. 51. 230–243.
- Chu, C. K., Glad, I. K., Godtliebsen, F. and Marron, J. S. (1998). Edge-preserving smoothers for image processing. *Journal of the American Statistical Association*. 93. 526–541.
- Comport, A. I., Marchand, E. and Chaumette, F. (2006). Statistically robust 2-D visual servoing. *IEEE Transactions on Robotics*. 22. 416–421.
- Dryden, I., Ippoliti, L. and Romagnoli, L. (2002). Adjusted maximum likelihood and pseudo-likelihood estimation for noisy Gaussian Markov random fields. *Journal of Computational and Graphical Statistics*. 11. 370–388.
- Frery, A. C., Sant’Anna, S. J. S., Mascarenhas, N. D. A. and Bustos, O. H. (1997). Robust inference techniques for speckle noise reduction in 1-look amplitude SAR images. *Applied Signal Processing*. 4. 61–76.
- Glendinning, R. H. (1999). Robust shape classification. *Signal Processing*. 77. 121–138.
- Gottardo, R., Raftery, A. E., Yeung, K. Y. and Bumgarner, R. E. (2006). Bayesian robust inference for differential gene expression in microarrays with multiple samples. *Biometrics*. 62. 10–18.
- Guyon, X. (1993). *Champs aléatoires sur un réseau, modélisations, statistique et applications*. Masson. Paris.
- Hasler, D., Sbaiz, L., Susstrunk, S. and Vetterli, M. (2003). Outlier modeling in image matching. *IEEE Transactions on Pattern Analysis and Machine Intelligence*. 25. 301–315.
- Huang, H. C. and Lee, T. C. A. (2006). Data adaptive median filters for signal and image denoising using a generalized SURE criterion. *IEEE Signal Processing Letters*. 13. 561–564.
- Kashyap, R. L. and Eom, K. B. (1988). Robust Image Techniques with an image restoration application. *IEEE Trans. on Acoustics, Speech, and Signal Processing*. 36. 1313–1325.
- Kim, J. H. and Han, J. H. (2006). Outlier correction from uncalibrated image sequence using the triangulation method. *Pattern Recognition*. 39. 394–404.
- Koivunen, V. (1995). A robust nonlinear filter for image-restoration. *IEEE Transactions on Image Processing*. 4. 569–578.

- Li, S. Z., Wang, H. and Soh, W. Y. C. (1998). Robust estimation of rotation angles from image sequences using the annealing M-estimator. *Journal of Mathematical Imaging and Vision*. 8. 181–192.
- Marroquin, J. L., Servin, M. and Rodriguez-Vera, R. (1998). Robust filters for low-level vision. *Expert Systems with Applications*. 14. 169–177.
- Meer, P., Mintz, D., Rosenfeld, A. and Kim, D. Y. (1991). Robust regression methods for computer vision: a review. *International Journal of Computer Vision*. 6. 59–70.
- Mirza, M. J. and Boyer, K. L. (1993). Performance evaluation of a class of M-estimators for surface parameter-estimation in noisy range data. *IEEE Transactions on Robotics and Automation*. 9. 75–85.
- Nirel, R., Mugglestone, M. A. and Barnett, V. (1998). Outlier-robust spectral estimation for spatial lattice processes. *Communications in Statistics-Theory and Methods*. 27. 3095–3111.
- Ojeda, S. (1999). Estimadores robustos RA en el modelo AR-2D aplicado al procesamiento estadístico de imágenes satelitales. Tesis doctoral. Facultad de Matemática, Astronomía y Física. Universidad Nacional de Córdoba, Argentina.
- Ojeda, S., Vallejos, R. and Lucini, M. (2002). Performance of robust RA estimator for bidimensional autoregressive models. *Journal of Statistical Computation and Simulation*. 72(1). 47–62.
- Palenichka, R. M., Zinterhof, P. and Ivasenko, I. B. (2000). Adaptive image filtering and segmentation using robust estimation of intensity. *Advances in Pattern Recognition*. 1876. 888–897.
- Palenichka, R. M., Zinterhof, P., Rytsar, Y. B. and Ivasenko, I. B. (1998). Structure-adaptive image filtering using order statistics. *Journal of Electronic Imaging*. 7. 339–349.
- Prastawa, M., Bullitt, E., Ho, S. and Gerig, G. (2004). A brain tumor segmentation framework based on outlier detection. *Medical Image Analysis*. 8. 275–283.
- Rabie, T. (2005). Robust estimation approach for blind denoising. *IEEE Transactions On Image Processing*. 14. 1755–1765.
- Roth, V. (2006). Kernel fisher discriminants for outlier detection. *Neural Computation*. 18. 942–960.
- Ruskin, D.M. (1978). M-estimates of nonlinear regression parameters and their jackknife constructed confidence intervals. Ph.D. Dissertation, University of California, Los Angeles.
- Sim, K. S., Cheng, Z. and Chuah, H. T. (2004). Robust image signal-to-noise ratio estimation using mixed Lagrange time delay estimation autoregressive model. *Scanning*. 26. 287–295.
- Singh, M., Arora, H. and Ahuja, N. (2004). A robust probabilistic estimation framework for parametric image models. *Computer Vision ECCV*. 3021. 508–522.
- Steland, A. (2005). On the distribution of the clipping median under a mixture model. *Statistics and Probability Letters*. 71. 1–13.
- Stewart, C. V. (1999). Robust parameter estimation in computer vision. *Siam Review*. 41. 513–537.
- Tarel, J. P., Ieng, S. S. and Charbonnier, P. (2002). Using robust estimation algorithms for tracking explicit curves. *Computer Vision ECCV*. 2350. 492–507.
- Tjøstheim, D. (1978). Statistical spatial series modelling. *Adv. Appl. Prob.* 10. 130–154.
- Tohka, J., Zijdenbos, A. and Evans, A. (2004). Fast and robust parameter estimation for statistical partial volume models in brain MRI. *Neuroimage*. 23. 84–97.
- Torr, P. H. S. and Zisserman, A. (2000). MLESAC: a new robust estimator with application to estimating image geometry. *Computer Vision and Image Understanding*. 78. 138–156.
- Vallejos, R. O. and Mardesic, T. J. (2004). A recursive algorithm to restore images based on robust estimation of NSHP autoregressive models. *Journal of Computational and Graphical Statistics*. 13. 674–682.

- Vallejos, R., Ojeda, S. and Bustos, O. (2006). On RA and GM estimates for first order spatial autoregressive models. (Submitted)
- Van de Weijer, J. and Van den Boomgaard, R. (2005). Least squares and robust estimation of local image structure. *International Journal of Computer Vision*. 64. 143–155.
- Voloshynovskiy, S. V., Allen, A. R. and Hrytskiv, Z. D. (2000). Robust edge-preserving image restoration in the presence of non-Gaussian noise. *Electronics Letters*. 36. 2006–2007.
- Wang, H. Z. and Suter, D. (2004a). MDPE: a very robust estimator for model fitting and range image segmentation. *International Journal of Computer Vision*. 59. 139–166.
- Wang, H. Z. and Suter, D. (2004b). Robust adaptive-scale parametric model estimation for computer vision. *IEEE Transactions on Pattern Analysis and Machine Intelligence*. 26. 1459–1474.
- Whittle, P. (1954). On stationary processes in the plane. *Biometrika*. 41. 434–449.
- Xu, P. L. (2005). Sign-constrained robust least squares, subjective breakdown point and the effect of weights of observations on robustness. *Journal of Geodesy*. 79. 146–159.
- Zervakis, M. E. and Kwon, T. M. (1992). Robust estimation techniques in regularized image-restoration. *Optical Engineering*. 31. 2174–2190.
- Zervakis, M. E., Katsaggelos, A. K. and Kwon, T. M. (1995). A class of robust entropic functionals for image-restoration. *IEEE Transactions on Image Processing*. 4. 752–773.