

WEIGHTED INEQUALITIES FOR INTEGRAL OPERATORS
WITH SOME HOMOGENEOUS KERNELS

MARÍA SILVINA RIVEROS, MARTA URUIOLO, Córdoba

(Received August 1, 2002)

Abstract. In this paper we study integral operators of the form

$$Tf(x) = \int |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} f(y) \, dy,$$

$\alpha_1 + \dots + \alpha_m = n$. We obtain the $L^p(w)$ boundedness for them, and a weighted $(1, 1)$ inequality for weights w in A_p satisfying that there exists $c \geq 1$ such that $w(a_i x) \leq cw(x)$ for a.e. $x \in \mathbb{R}^n$, $1 \leq i \leq m$. Moreover, we prove $\|Tf\|_{\text{BMO}} \leq c\|f\|_\infty$ for a wide family of functions $f \in L^\infty(\mathbb{R}^n)$.

Keywords: weights, integral operators

MSC 2000: 42B25, 42A50, 42B20

1. INTRODUCTION

In [7] the authors study the boundedness on $L^2(\mathbb{R})$ of the operator

$$Tf(x) = \int |x - y|^{-\alpha} |x + y|^{\alpha-1} f(y) \, dy,$$

$0 < \alpha < 1$.

In [3] the authors study integral operators of the form

$$Tf(x) = \int_{\mathbb{R}^n} |x - y|^{-\alpha} |x + y|^{-n+\alpha} f(y) \, dy,$$

$0 < \alpha < n$. They obtain the $L^p(\mathbb{R}^n, dx)$ boundedness and the weak type $(1, 1)$ of them.

Partially supported by CONICET, Agencia Córdoba Ciencia and SECYT-UNC.

In this paper we consider integral operators defined for f belonging to the Schwartz class $S(\mathbb{R}^n)$ by

$$(1.1) \quad Tf(x) = \int_{\mathbb{R}^n} |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} f(y) dy,$$

$\alpha_1 + \dots + \alpha_m = n$, $\alpha_i > 0$ and $a_i \in \mathbb{R} - \{0\}$ for $i = 1, \dots, m$.

We take the Hardy-Littlewood maximal function as

$$Mf(x) = \sup_Q \frac{1}{|Q|} \int_Q |f(x)| dx$$

where the supremum is taken along all cubes Q such that x belongs to Q . We recall that a weight w is a measurable, non negative and locally integrable function. It is well known that, for $p > 1$, M is bounded on $L^p(w)$ if and only if there exists $c > 0$ such that

$$(1.2) \quad \sup_Q \left(\frac{1}{|Q|} \int_Q w \right) \left(\frac{1}{|Q|} \int_Q w^{-1/(p-1)} \right)^{p-1} \leq c.$$

The class of functions that satisfy (1.2) is denoted by A_p . For $p = 1$, the class A_1 is defined by

$$Mw(x) \leq cw(x)$$

for a.e. $x \in \mathbb{R}^n$ and for some positive constant $c > 0$. The weak type (1,1) of the maximal function is equivalent to $w \in A_1$. These classes A_p have been defined by Muckenhoupt (see [6]) in the one dimensional case and for higher dimensions by Coifmann and Fefferman (see [1]).

In this paper we obtain the boundedness of T on $L^p(\mathbb{R}^n, w)$ and a weighted (1,1) inequality for a wide class of weights w in A_p . We prove the following result:

Theorem 1. *Let T be defined by (1.1). Suppose there exists $c \geq 1$ such that $w(a_i x) \leq cw(x)$ for $1 \leq i \leq m$ and for almost every $x \in \mathbb{R}^n$.*

- a) *If $w \in A_p$, $1 < p < \infty$ then T is bounded on $L^p(\mathbb{R}^n, w)$.*
- b) *If $w \in A_1$ then there exists $k > 0$ such that, for $\lambda > 0$ and $f \in S(\mathbb{R}^n)$,*

$$w(\{x: |Tf(x)| > \lambda\}) \leq \frac{k}{\lambda} \int |f(x)|w(x) dx.$$

We also analyze the boundedness of the operator T from L^∞ into BMO, the classical space consisting of functions with bounded mean oscillation, defined by

John and Nirenberg in [5]. Precisely, we say that $f \in L^1_{\text{loc}}$ belongs to BMO if there exist $c > 0$ such that

$$\frac{1}{|Q|} \int \left| f(x) - \frac{1}{|Q|} \int f \right| dx \leq c$$

for all cubes $Q \subset \mathbb{R}^n$. The smallest bound c for which the above inequality holds is called $\|f\|_*$. From the techniques used, the following result follows immediately:

Theorem 2. *Let T be defined by (1.1). Then there exists $c > 0$ such that*

$$\|Tf\|_* \leq c\|f\|_\infty$$

for all $f \in S(\mathbb{R}^n)$.

If f is a positive constant then $Tf(x) = \infty$ for all $x \in \mathbb{R}^n$, so we cannot expect a general boundedness from L^∞ into BMO. With techniques similar to those developed in [8], we obtain

Theorem 3. *Let T be defined by (1.1).*

- a) *If $f \in L^\infty$ and $T|f|(x_0) < \infty$ for some $x_0 \in \mathbb{R}^n$ then $Tf(x)$ is well defined for all $x \neq 0$ and $Tf \in L^1_{\text{loc}}(\mathbb{R}^n)$.*
- b) *There exists $c > 0$ such that*

$$\|Tf\|_* \leq c\|f\|_\infty$$

for all f as in a).

By c we denote a positive constant, not the same at each occurrence.

PROOF OF THE MAIN RESULTS

We follow the argument developed in [2, p. 144] where the case of the Calderón-Zygmund operators is treated. As there we define, for $f \in L^1_{\text{loc}}(\mathbb{R}^n)$, the sharp maximal function by

$$M^\# f(x) = \sup_{Q: x \in Q} \frac{1}{|Q|} \int_Q |f - f_Q|(y) dy$$

with $f_Q = \frac{1}{|Q|} \int_Q f$.

We denote $D = \max_{1 \leq i \leq m} |a_i^{-1}|$ and $d = \min_{1 \leq i \leq m} |a_i^{-1}|$. We need the following result:

Lemma 1.3. *If T is defined by (1.1) and $s > 1$ then there exists $c > 0$ such that for all $f \in S(\mathbb{R}^n)$,*

$$M^\#Tf(x) \leq c[(Mf^s(a_1^{-1}x))^{1/s} + \dots + (Mf^s(a_m^{-1}x))^{1/s}].$$

Proof. We first observe that T is a bounded operator on $L^p(\mathbb{R}^n, dx)$, $1 < p < \infty$ (see [4]), so for $f \in S(\mathbb{R}^n)$, $Tf \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $M^\#Tf(x)$ is well defined for all $x \in \mathbb{R}^n$. We take $x \in \mathbb{R}^n$ such that $T|f|(x) < \infty$ and Q a cube that contains x . We set $l(Q)$ as the length of the side of Q , denote by \overline{Q} the cube with the same center as Q , such that $l(\overline{Q}) \geq 2D/d \cdot l(Q)$ and, for $1 \leq i \leq m$, we also set $\overline{Q}_i = a_i^{-1}\overline{Q}$. We decompose $f = f_1 + f_2$, $f_1 = f\chi_{(\cup_{1 \leq k \leq m} \overline{Q}_k)^c}$ and take $a = Tf_2(x)$. Then

$$\frac{1}{|Q|} \int_Q |Tf(y) - a| dy \leq \frac{1}{|Q|} \int_Q |Tf_1(y)| dy + \frac{1}{|Q|} \int_Q |Tf_2(y) - Tf_2(x)| dy.$$

If $s > 1$ then T is bounded on $L^s(\mathbb{R}^n, dx)$ (see [4]), so

$$\begin{aligned} \frac{1}{|Q|} \int_Q |Tf_1(y)| dy &\leq \left(\frac{1}{|Q|} \int_Q |Tf_1(y)|^s dy \right)^{1/s}, \\ \left(\frac{1}{|Q|} \int_{\overline{Q}_1} |f(y)|^s dy \right)^{1/s} + \dots + \left(\frac{1}{|Q|} \int_{\overline{Q}_m} |f(y)|^s dy \right)^{1/s} \\ &\leq c'[(Mf^s(a_1^{-1}x))^{1/s} + \dots + (Mf^s(a_m^{-1}x))^{1/s}]. \end{aligned}$$

On the other hand,

$$\frac{1}{|Q|} \int_Q |Tf_2(y) - Tf_2(x)| dy \leq \frac{1}{|Q|} \int_Q \left| \int_{(\cup_{1 \leq k \leq m} \overline{Q}_k)^c} (K(y, z) - K(x, z))f(z) dz \right| dy$$

where we denote by $K(x, y)$ the kernel $|x - a_1y|^{-\alpha_1} \dots |x - a_my|^{-\alpha_m}$. □

We now estimate $|K(y, z) - K(x, z)|$.

Case $l(Q) \geq 2|x|$. In this situation $\bigcup_{1 \leq k \leq m} \overline{Q}_k \supset \{y: |y| < 3D|x|\}$. Indeed, if $z \in (\bigcup_{1 \leq k \leq m} \overline{Q}_k)^c$, then $|z| \geq |z - a_1^{-1}x| - |a_1^{-1}x| \geq l(\overline{Q}_1) - D|x| \geq dl(\overline{Q}) - D|x| \geq 3D|x|$. Moreover, in this case $|x - a_1z| \leq |x| + |a_1z| \leq (|a_1| + \frac{1}{3D})|z|$ then

$$\begin{aligned} (1.4) \quad |x - a_i z| &\geq |a_i z| - |x| \geq \left(|a_i| - \frac{1}{3d} \right) |z| \\ &\geq \left(\frac{3|a_i|D - 1}{3|a_1|D + 1} \right) \frac{1}{2} |x - a_1 z|. \end{aligned}$$

Thus we apply the mean value theorem to obtain, for $x, y \in Q$ and $z \in \left(\bigcup_{1 \leq k \leq m} \overline{Q}_k \right)^c$,

$$|K(y, z) - K(x, z)| \leq |x - y| \sum_{i=1}^m \frac{\alpha_i}{|\xi - a_i z|^{\alpha_i+1} \prod_{l \neq i} |\xi - a_l z|^{\alpha_l}}$$

for some ξ between x and y . But $|a_i^{-1}\xi - z| \geq |a_i^{-1}x - z| - |a_i^{-1}\xi - a_i^{-1}x| \geq \frac{1}{2}|a_i^{-1}x - z|$, so (1.4) implies

$$(1.5) \quad |K(y, z) - K(x, z)| \leq c \frac{|x - y|}{|x - a_1 z|^{n+1}}.$$

Thus

$$\begin{aligned} & \frac{1}{|Q|} \int_Q \left| \int_{\left(\bigcup_{1 \leq k \leq m} \overline{Q}_k \right)^c} (K(y, z) - K(x, z)) f(z) dz \right| dy \\ & \leq \frac{c}{|Q|} \int_Q \sum_{k=1}^{\infty} \int_{2^k D l(Q) \leq |a_1^{-1}x - z| < 2^{k+1} D l(Q)} \frac{|x - y|}{|a_1^{-1}x - z|^{n+1}} |f(z)| dz dy \\ & \leq c l(Q) \sum_{k=1}^{\infty} \frac{1}{2^k D l(Q)} \frac{1}{(2^k D l(Q))^n} \int_{|a_1^{-1}x - z| < 2^{k+1} D l(Q)} |f(z)| dz \\ & \leq c M f(a_1^{-1}x) \leq c (M f^s(a_1^{-1}x))^{1/s}. \end{aligned}$$

Case $l(Q) < 2|x|$. We decompose

$$\int_{\left(\bigcup_{1 \leq k \leq m} \overline{Q}_k \right)^c} (K(y, z) - K(x, z)) f(z) dz = \int_{|z| \geq 3D|x|} + \int_{\{|z| < 3D|x|\} \cap \left(\bigcup_{1 \leq k \leq m} \overline{Q}_k \right)^c}.$$

To estimate the first integral, we proceed as before and we obtain (1.5) for $x, y \in Q$ and $|z| \geq 3D|x|$, then

$$\frac{1}{|Q|} \int_Q \left| \int_{|z| \geq 3D|x|} (K(y, z) - K(x, z)) f(z) dz \right| dy \leq c (M f^s(a_1^{-1}x))^{1/s}.$$

We now study the second integral. For $1 \leq i \leq m$, $x, y \in Q$ and $z \in \{z: |z| < 3D|x|\} \cap \left(\bigcup_{1 \leq k \leq m} \overline{Q}_k \right)^c$, we have

$$|a_i^{-1}y - z| \geq |a_i^{-1}x - z| - |a_i^{-1}y - a_i^{-1}x| \geq \frac{|a_i^{-1}x - z|}{2},$$

hence

$$|K(y, z) - K(x, z)| \leq c |K(x, z)|.$$

So

$$\begin{aligned} & \int_{\{|z| < 3D|x|\} \cap (\bigcup_{1 \leq k \leq m} \overline{Q}_k)^c} (K(y, z) - K(x, z)) f(z) \, dz \\ & \leq c \int_{\{|z| < 3D|x|\}} \frac{|f(z)|}{|x - a_1 z|^{\alpha_1} \dots |x - a_m z|^{\alpha_m}} \, dz. \end{aligned}$$

We define $b = \frac{1}{2} \min_{1 \leq l, j \leq m} (|a_l^{-1} - a_j^{-1}|)$. We set $A_i = \{z: |a_i^{-1}x - z| \leq b|x|\}$, $1 \leq i \leq m$, and $A_{m+1} = \left(\bigcup_{i=1}^m A_i\right)^c$ and decompose

$$\begin{aligned} & \int_{\{|z| < 3D|x|\}} \frac{|f(z)|}{|x - a_1 z|^{\alpha_1} \dots |x - a_m z|^{\alpha_m}} \, dz \\ & = \int_{A_1} + \dots + \int_{A_m} + \int_{A_{m+1} \cap \{|z| < 3D|x|\}}. \end{aligned}$$

For $z \in A_i$ and $l \neq i$ we have $|a_l^{-1}x - z| \geq b|x|$, hence

$$\begin{aligned} & \int_{A_i} \frac{|f(z)|}{|x - a_1 z|^{\alpha_1} \dots |x - a_m z|^{\alpha_m}} \, dz \\ & \leq \frac{c}{|x|^{n-\alpha_i}} \sum_{j=0}^{\infty} \int_{2^{-j-1}b|x| \leq |a_i^{-1}x - z| \leq 2^{-j}b|x|} \frac{|f(z)|}{|a_i^{-1}x - z|^{\alpha_i}} \, dz \\ & \leq c \sum_{j=1}^{\infty} 2^{j(\alpha_i - n)} \frac{1}{(2^{-j}b|x|)^n} \\ & \quad \times \int_{|z - a_i^{-1}x| \leq 2^{-j}b|x|} |f(z)| \, dz \leq cMf(a_i^{-1}x) \leq c(Mf^s(a_i^{-1}x))^{1/s}. \end{aligned}$$

Now

$$\begin{aligned} & \int_{A_{m+1} \cap \{|z| < 3D|x|\}} \frac{|f(z)|}{|x - a_1 z|^{\alpha_1} \dots |x - a_m z|^{\alpha_m}} \, dz \leq c|x|^{-n} \int_{\{|z| < 3D|x|\}} |f(z)| \, dz \\ & \leq cMf(a_1^{-1}x) \leq c(Mf^s(a_1^{-1}x))^{1/s}, \end{aligned}$$

and the lemma follows. \square

Lemma 1.6. *Let T be defined by (1.1), $1 < p < \infty$, $w \in A_p$ and $f \in L^p(w)$. Then $Tf \in L^p(w)$.*

Proof. If $\text{supp } f \subset B(0, R)$ and $|x| > 2R$ then $|K(x, y)| \leq c/|x|^n$ and so in this case $|Tf(x)| \leq c_R/|x|^n$. The proof follows as in Theorem 7.18 in [2], since T is a bounded operator on $L^p(\mathbb{R}^n, dx)$ (see [4]). \square

Proof of Theorem 1. a) Taking account of Lemmas 1.3 and 1.6, we proceed as in the proof of Theorem 7.18 in [2] to obtain, for $f \in S(\mathbb{R}^n)$,

$$\begin{aligned} & \int |Tf(x)|^p w(x) \, dx \\ & \leq c \int |(Mf^s(a_1^{-1}x))^{1/s} + \dots + (Mf^s(a_m^{-1}x))^{1/s}|^p w(x) \, dx \\ & \leq c \int |Mf^s(x)|^{p/s} w(a_1x) \, dx + \dots + \int |Mf^s(x)|^{p/s} w(a_mx) \, dx \\ & \leq c \int |Mf^s(x)|^{p/s} w(x) \, dx. \end{aligned}$$

The last inequality follows from the hypothesis about the weight w . The rest of the proof is as in Theorem 7.18 in [2].

b) For $\lambda > 0$ we perform the Calderón-Zygmund decomposition for f to obtain a sequence of disjoint $\{Q_j\}_{j \in \mathbb{N}}$ such that $f(x) \leq \lambda$ for almost every $x \notin \bigcup_{j \in \mathbb{N}} Q_j$. We take

$$g(x) = \begin{cases} f(x) & \text{if } x \notin \bigcup_{j \in \mathbb{N}} Q_j, \\ \frac{1}{|Q_j|} \int_{Q_j} f & \text{if } x \in Q_j \end{cases}$$

and write $f = g + b$.

As usual, from a), we obtain

$$w\{x: |Tg(x)| > \lambda\} \leq \frac{c}{\lambda} \int |f(x)| w(x) \, dx.$$

For each $i = 1, \dots, m$ and $j \in \mathbb{N}$ we denote by $\overline{Q_j}$ the cube with the same center as Q_j and such that $l(\overline{Q_j}) \geq 2D/d \cdot l(Q_j)$, and $\overline{Q_{j,i}} = a_i \overline{Q_j}$. We obtain

$$\begin{aligned} w\left(\bigcup_{j \in \mathbb{N}} \overline{Q_{j,i}}\right) & \leq \sum_{j \in \mathbb{N}} w(\overline{Q_{j,i}}) \leq c \sum_{j \in \mathbb{N}} \frac{w(\overline{Q_{j,i}})}{|\overline{Q_{j,i}}|} |\overline{Q_{j,i}}| \\ & \leq c \sum_{j \in \mathbb{N}} |Q_j| \frac{w(\overline{Q_{j,i}})}{|\overline{Q_{j,i}}|} \leq \sum_{j \in \mathbb{N}} \frac{c}{\lambda} \int_{Q_j} |f| \frac{w(\overline{Q_{j,i}})}{|\overline{Q_{j,i}}|} \\ & \leq \frac{c}{\lambda} \sum_{j \in \mathbb{N}} \int_{Q_j} |f(y)| Mw(a_i y) \, dy \\ & \leq \frac{c}{\lambda} \int |f(y)| w(a_i y) \, dy \leq \frac{c}{\lambda} \int |f(y)| w(y) \, dy. \end{aligned}$$

Then

$$w\left(\bigcup_{j \in \mathbb{N}, i=1, \dots, m} \overline{Q_{j,i}}\right) \leq \frac{c}{\lambda} \int |f(y)| w(y) \, dy.$$

Now for each fixed $i = 1, \dots, m$, if c_j denotes the center of Q_j , we have

$$\begin{aligned} & w\left(\{x: |Tb(x)| > \lambda\} \cap \left(\bigcup_{j \in \mathbb{N}} \overline{Q_{j,i}}\right)^c\right) \\ & \leq \frac{c}{\lambda} \sum_{j \in \mathbb{N}} \int_{(\overline{Q_{j,i}})^c} \left| \int_{Q_j} b_j(y) (K(x,y) - K(x,c_j)) dy \right| w(x) dx \\ & \leq \frac{c}{\lambda} \sum_{j \in \mathbb{N}} \int_{Q_j} |b_j(y)| \int_{(\overline{Q_{j,i}})^c} |K(x,y) - K(x,c_j)| w(x) dx dy. \end{aligned}$$

Now we observe that $K(x,y) = c\tilde{K}(y,x)$ where $\tilde{K}(x,y) = |x - a_1^{-1}y|^{-\alpha_1} \dots |x - a_m^{-1}y|^{-\alpha_m}$. Reasoning as in a) with \tilde{K} instead of K and using the hypothesis on w , we get

$$\int_{(\overline{Q_{j,i}})^c} |K(x,y) - K(x,c_j)| w(x) dx \leq cMw(a_j y) \leq cw(y).$$

So

$$\begin{aligned} & w\left(\{x: |Tb(x)| > \lambda\} \cap \left(\bigcup_{j \in \mathbb{N}, i=1, \dots, m} \overline{Q_{j,i}}\right)^c\right) \\ & \leq \frac{c}{\lambda} \int |b(y)| w(y) dy \leq \frac{c}{\lambda} \int |f(y)| w(y) dy. \end{aligned}$$

□

Proof of Theorem 2. It follows straightforward from Lemma 1.3. □

Proof of Theorem 3. a) Let $f \in L^\infty(\mathbb{R}^n)$ and let x_0 be such that $T|f|(x_0) < \infty$. We take $R = 4D|x_0|$, denote $B = B(0, R) = \{x \in \mathbb{R}^n: |x| \leq R\}$, define $f_1 = |f|\chi_B$ and decompose $|f| = f_1 + f_2$. Then

$$\begin{aligned} Tf_1(x) & \leq \int_B |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} f(y) dy \\ & \leq \|f\|_\infty \int_B |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} dy. \end{aligned}$$

If $x \neq 0$ we choose $r > 0$ such that $r = \frac{1}{4} \min_{1 \leq i, k \leq m} |a_i^{-1} - a_k^{-1}| |x|$. For $1 \leq i \leq m$, we define $B_i = B(a_i^{-1}x, r)$. We have

$$\begin{aligned} & \int_B |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} dy \\ & \leq \sum_{1 \leq i \leq m} \int_{B_i} |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} dy \\ & \quad + \int_{B \cap (\bigcup_{1 \leq i \leq m} B_i)^c} |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} dy. \end{aligned}$$

Now

$$\begin{aligned} & \int_{B_i} |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} dy \\ & \leq c \prod_{k \neq i} r^{-\alpha_k} \int_{B_i} |x - a_i y|^{-\alpha_i} dy \leq c \prod_{k \neq i} r^{-\alpha_k} r^{-\alpha_i + n} = c. \end{aligned}$$

If $|a_i^{-1}x| < 2R$ for some $1 \leq i \leq m$, then, for $y \in B \cap (B_i)^c$, we have $r < |a_i^{-1}x - y| \leq 3R$ and so

$$\begin{aligned} & \int_{B \cap (\bigcup_{1 \leq i \leq m} B_i)^c} |x - a_1 y|^{-\alpha_1} \dots |x - a_m y|^{-\alpha_m} dy \\ & \leq c \prod_{k \neq i} r^{-\alpha_k} \int_{B \cap (B_i)^c} |x - a_i y|^{-\alpha_i} dy \\ & \leq c \prod_{k \neq i} r^{-\alpha_k} \int_r^{3R} t^{-\alpha_i + n - 1} dt \\ & = c \prod_{k \neq i} r^{-\alpha_k} [(3R)^{-\alpha_i + n} - r^{-\alpha_i + n}] = c \left(|x|^{\sum_{k \neq i} -\alpha_k} + 1 \right), \end{aligned}$$

so for $x \neq 0$ and such that $|a_i^{-1}x| < 2R$ we obtain

$$(1.7) \quad |Tf_1(x)| \leq c \|f\|_\infty \left(1 + |x|^{\sum_{k \neq i} -\alpha_k} \right).$$

Now if $|a_i^{-1}x| \geq 2R$ for all $1 \leq i \leq m$, then $|a_i^{-1}x - y| \geq R$ for $y \in B(0, R)$ and so

$$|Tf_1(x)| \leq \|f\|_\infty.$$

So (1.7) holds for all $x \neq 0$. Then $Tf_1(x) < \infty$ for all $x \neq 0$ and it belongs to $L_{\text{loc}}^1(\mathbb{R}^n)$.

Now $Tf_2(x_0) < \infty$ so we write, for $x \in \mathbb{R}^n$, $Tf_2(x) = Tf_2(x) - Tf_2(x_0) + Tf_2(x_0)$. Then we have to study

$$\int_{B^c} |K(x, y) - K(x_0, y)| |f|(y) dy.$$

For $x \neq 0$ we have

$$\begin{aligned} \int_{B^c} |K(x, y) - K(x_0, y)| |f|(y) dy & \leq \int_{B^c \cap B(0, 4D|x|)^c} |K(x, y) - K(x_0, y)| |f|(y) dy \\ & \quad + \int_{B^c \cap B(0, 4D|x|)} |K(x, y)| |f|(y) dy + c. \end{aligned}$$

To estimate the first integral, we proceed as in the proof of Lemma 1.3 to obtain that, for $y \in B^c \cap B(0, 4D|x|)^c$,

$$|K(x, y) - K(x_0, y)| \leq c \frac{|x - x_0|}{|x - a_1 y|^{n+1}},$$

so

$$\begin{aligned} \int_{B^c \cap B(0, 4D|x|)^c} |K(x, y) - K(x_0, y)| |f|(y) \, dy &\leq c|x - x_0| \int_{B^c} \frac{|f|(y)}{|x - a_1 y|^{n+1}} \, dy \\ &\leq c|x - x_0| \|f\|_\infty. \end{aligned}$$

To study the second integral, we observe that it appears only if $D|x| \geq R/4$, so we proceed as in the previous estimate for Tf_1 to obtain that, for x in this region,

$$\int_{B^c \cap B(0, 4D|x|)} |K(x, y)| |f|(y) \, dy \leq c \|f\|_\infty.$$

So, for $x \neq 0$, $Tf_2(x) < \infty$ and it belongs to $L^1_{\text{loc}}(\mathbb{R}^n)$.

b) If f satisfies the hypothesis of a) we obtain that $M^\#Tf(x)$ is well defined for all $x \in \mathbb{R}^n$, so Lemma 1.3 still holds for these functions, and b) follows. \square

References

- [1] *R. Coifmann, C. Fefferman*: Weighted norm inequalities for maximal functions and singular integrals. *Studia Math.* 51 (1974), 241–250.
- [2] *J. Duoandikoetxea*: Análisis de Fourier. Ediciones de la Universidad Autónoma de Madrid, Editorial Siglo XXI, 1990.
- [3] *T. Godoy, M. Urciuolo*: About the L^p boundedness of some integral operators. *Revista de la UMA* 38 (1993), 192–195.
- [4] *T. Godoy, M. Urciuolo*: On certain integral operators of fractional type. *Acta Math. Hungar.* 82 (1999), 99–105.
- [5] *F. John, L. Nirenberg*: On functions of bounded mean oscillation. *Comm. Pure Appl. Math.* 14 (1961), 415–426.
- [6] *B. Muckenhoupt*: Weighted norm inequalities for the Hardy maximal function. *Trans. Amer. Math. Soc.* 165 (1972), 207–226.
- [7] *F. Ricci, P. Sjögren*: Two parameter maximal functions in the Heisenberg group. *Math. Z.* 199 (1988), 565–575.
- [8] *A. de la Torre, J.L. Torrea*: One-sided discrete square function. *Studia Math.* 156 (2003), 243–260.

Authors' address: María Silvana Riveros, Marta Urciuolo, FaMAF Universidad Nacional de Córdoba, Ciem-CONICET, Ciudad Universitaria 5000 Córdoba, e-mails: sriveros@mate.uncor.edu, urciuolo@mate.uncor.edu.