

ON THE BEST RANGES FOR A_p^+ AND RH_r^+

M. S. RIVEROS, Córdoba, and A. DE LA TORRE, Málaga

(Received May 28, 1998)

Abstract. In this paper we study the relationship between one-sided reverse Hölder classes RH_r^+ and the A_p^+ classes. We find the best possible range of RH_r^+ to which an A_1^+ weight belongs, in terms of the A_1^+ constant. Conversely, we also find the best range of A_p^+ to which a RH_∞^+ weight belongs, in terms of the RH_∞^+ constant. Similar problems for A_p^+ , $1 < p < \infty$ and RH_r^+ , $1 < r < \infty$ are solved using factorization.

Keywords: one-sided weights, one-sided reverse Hölder, factorization

MSC 2000: 42B25

1. INTRODUCTION

It is well known that there is a relationship between the A_p classes and the so called reverse Hölder classes RH_r . C. J. Neugebauer [8] studied the following problems:

- (1) For $w \in A_p$, find the precise range of r 's such that $w \in RH_r$, the precise range of $q < p$ for which $w \in A_q$, and the precise range of $s > 1$ such that $w^s \in A_p$.
- (2) Conversely, for a fixed $w \in RH_r$, find the precise range of p 's such that $w \in A_p$, and the precise range of $q > r$ for which $w \in RH_q$.

For the one-sided Hardy-Littlewood maximal operator

$$M^+ f(x) = \sup_{h>0} \frac{1}{h} \int_x^{x+h} |f|,$$

the A_p^+ classes were introduced by E. Sawyer [9]. He proved that M^+ is bounded in $L^p(w)$ ($p > 1$) if, and only if, the weight satisfies A_p^+ , i.e., there exists a constant C

Research supported by D.G.I.Y.T. (PB94-1496), Junta de Andalucía and Universidad Nacional de Córdoba.

such that for any three points $a < b < c$ we have

$$\int_a^b w \left(\int_b^c w^{1-p'} \right)^{p-1} \leq C(c-a)^p.$$

The smallest constant for which this is satisfied will be called the A_p^+ constant of w and will be denoted by $A_p^+(w)$. For $p = 1$ the weak type of the operator holds if, and only if, the weight w satisfies A_1^+ , i.e., there exists C such that for any a and almost every $b > a$,

$$\int_a^b w \leq C(b-a)w(b).$$

The smallest such constant will be called the A_1^+ constant of w and will be denoted by $A_1^+(w)$. For later reference we point out that it is an easy consequence of Lebesgue's differentiation theorem that the constant in the definition of A_1^+ is always greater than, or equal to, one.

These classes are of interest, not only because they control the boundedness of the one-sided Hardy-Littlewood maximal operator, but they are the right classes for the weighted estimates for one-sided singular integrals [1] and they also appear in PDE [4]. In contrast to the Muckenhoupt weights, the one-sided weights are not doubling, but they possess a one-sided doubling property, namely if $w \in A_p^+$ then there exists C such that for any $a \in \mathbb{R}$ and $h > 0$, $\int_a^{a+2h} w \leq C \int_{a+h}^{a+2h} w$. The reverse Hölder property is not satisfied by these weights, either, but nevertheless, Martín-Reyes [5] proved that there is a weak substitute of this notion, that we will denote by RH_r^+ , which is good enough to prove the " $p - \varepsilon$ " property. In [7] the class A_∞^+ was introduced and it was proved that $A_\infty^+ = \bigcup_{p < \infty} A_p^+ = \bigcup_{1 < r} RH_r^+$.

In this note we solve the problems of the Neugebauer paper in this context. In the proofs we will make essential use of the one-sided minimal operator introduced by Cruz-Uribe, Neugebauer and Olesen [3]. It is defined as $m^+f(x) = \inf_{c > x} \frac{1}{c-x} \int_x^c |f|$. We will also use the fact that for any positive function g , the maximal operator $M_g f(x) = \sup_{x \in I} \frac{1}{g(I)} \int_I |f|g dx$ is of weak type one-one with respect to the measure $g dx$. Note that for $g = 1$ we have the classical Hardy-Littlewood maximal operator, which is denoted by Mf .

The paper is organized as follows: in Section 2 we give definitions and characterizations of RH_r^+ , $1 < r < \infty$. In Section 3 we prove two theorems of the best range for the extreme classes A_1^+ and RH_∞^+ . In Section 4 we give a factorization theorem for weights in RH_r^+ , and finally in Section 5 we extend the theorems of Section 3 to A_p^+ and RH_r^+ , using the factorization proved in Section 4. We shall see that the index range depends on the factorization of the weight.

We end this introduction with some notation: for a given interval $I = (a, a + h)$ we denote by I^- the interval $(a - h, a)$, I^+ the interval $(a + h, a + 2h)$, and I^{++} the interval $(a + 2h, a + 3h)$. For any $1 < p < \infty$, p' will be its conjugate exponent, if g is locally integrable and E is a measurable set, $g(E)$ will stand for $\int_E g$ and C will represent a constant that may change from time to time. Finally, we remark that we can change the orientation on the real line obtaining similar results for classes $RH_r^-, A_p^-, 1 < r \leq \infty$ and $1 \leq p \leq \infty$.

2. DEFINITION, CHARACTERIZATION OF RH_r^+ FOR $1 < r < \infty$

We start this section with the definition of RH_r^+ , $1 < r < \infty$.

Definition 2.1. A weight w satisfies the one-sided reverse Hölder RH_r^+ condition, if there exists C such that for any $a < b$,

$$(2.2) \quad \int_a^b w^r \leq C (M(w\chi_{(a,b)})(b))^{(r-1)} \int_a^b w.$$

The smallest such constant will be called the RH_r^+ constant of w and will be denoted by $RH_r^+(w)$.

Definition 2.3. A weight satisfies the one-sided reverse Hölder RH_∞^+ condition, if there exists C such that

$$(2.4) \quad w(x) \leq Cm^+ w(x)$$

for almost all $x \in \mathbb{R}$.

The smallest such constant will be called the RH_∞^+ constant of w and will be denoted by $RH_\infty^+(w)$. It is clear that $C \geq 1$.

The following lemma gives several characterizations of RH_r^+ . The constants are not necessarily the same.

Lemma 2.5. Let $a < b < c < d$, $1 < r < \infty$, and let $w \geq 0$ be locally integrable. Then the following statements are equivalent.

- (i) $\int_a^b w^r \leq C (M(w\chi_{(a,b)})(b))^{(r-1)} \int_a^b w.$
- (ii) $\frac{1}{b-a} \int_a^b w^r \leq C \left(\frac{1}{c-b} \int_b^c w \right)^r$ with $b-a = 2(c-b).$
- (iii) $\frac{1}{b-a} \int_a^b w^r \leq C \left(\frac{1}{d-c} \int_c^d w \right)^r$ with $b-a = d-b = 2(d-c).$

- (iv) $\frac{1}{b-a} \int_a^b w^r \leq C \left(\frac{1}{c-b} \int_b^c w \right)^r$ with $b-a = c-b$.
- (v) $\frac{1}{b-a} \int_a^b w^r \leq C \left(\frac{1}{d-c} \int_c^d w \right)^r$ with $b-a = d-c = \gamma(d-a)$, $0 < \gamma \leq \frac{1}{2}$.

Proof. To see $i) \implies ii)$, we fix $a < b < c$, $b-a = 2(c-b)$ and take any $x \in (b, c)$. Then

$$\int_a^b w^r \leq \int_a^x w^r \leq C (M(w\chi_{(a,x)})(x))^{r-1} \int_a^x w \leq C (M(w\chi_{(a,c)})(x))^{r-1} \int_a^c w.$$

Therefore $(b, c) \subset \{x: (M(w\chi_{(a,c)})(x))^{r-1} \geq \frac{1}{C} \int_a^b w^r\}$. The weak type $(1, 1)$ of the Hardy-Littlewood maximal operator yields

$$(c-b) \left(\int_a^b w^r \right)^{\frac{1}{r-1}} \leq C \left(\int_a^c w \right)^{\frac{r}{r-1}},$$

which implies

$$\frac{1}{b-a} \int_a^b w^r \leq C \left(\frac{1}{c-b} \int_a^c w \right)^r \leq C \left(\frac{1}{c-b} \int_b^c w \right)^r;$$

the last inequality follows from the fact, proved in [7], that a weight satisfying $i)$ satisfies A_p^+ for some p and thus it satisfies the one-sided doubling condition.

We will prove now that $ii) \implies i)$. Let us fix $a < b$ and define a sequence (x_k) as follows: $x_0 = a$ and $b-x_k = 2(b-x_{k+1})$. In particular, $x_{k+1} - x_k = 2(x_{k+2} - x_{k+1}) = (b-x_{k+1})$. Using condition $ii)$ for the points x_k, x_{k+1}, x_{k+2} , we have

$$\begin{aligned} \int_a^b w^r &= \sum_0^\infty \int_{x_k}^{x_{k+1}} w^r \leq C \sum_0^\infty (x_{k+1} - x_k)^{1-r} \left(\int_{x_{k+1}}^{x_{k+2}} w \right)^r \\ &\leq C \sum_0^\infty \int_{x_{k+1}}^{x_{k+2}} w \left(\frac{1}{b-x_{k+1}} \int_{x_{k+1}}^b w \right)^{r-1} \leq (M(w\chi_{(a,b)})(b))^{r-1} C \int_a^b w. \end{aligned}$$

To see $ii) \implies iii)$ let $a < b < c < d$ with $b-a = d-b = 2(d-c)$. Using that w satisfies the one-sided doubling condition, we have

$$\begin{aligned} \frac{1}{b-a} \int_a^b w^r &\leq C \left(\frac{1}{c-b} \int_b^c w \right)^r \leq C \left(\frac{d-c}{c-b} \frac{1}{d-c} \int_b^d w \right)^r \\ &\leq C \left(\frac{1}{d-c} \int_c^d w \right)^r. \end{aligned}$$

$iii) \implies iv)$ is immediate.

First of all we observe that iv) easily implies that the weight w satisfies the one-sided doubling condition. To see that iv) \implies v), let $0 < \gamma \leq \frac{1}{2}$ and $a < b < c < d$, $b - a = d - c = \gamma(d - a)$. Then if x is the midpoint between a and d we have

$$\frac{1}{b-a} \int_a^b w^r \leq \frac{1}{2\gamma} \frac{1}{x-a} \int_a^x w^r \leq \frac{C}{2\gamma} \left(\frac{1}{d-x} \int_x^d w \right)^r,$$

but it follows from the one-sided doubling condition that $\int_x^d w \leq C_\gamma \int_c^d w$.

Suppose v) holds, let $a < b < c$, $b - a = c - b = h$ and let us define for $k = 0, 1, \dots, N$ $x_k = a + ksh$ and $y_k = b + ksh$ where $s = \frac{\gamma}{1-\gamma}$ and N is the first integer such that $(N+1)s > 1$. We observe that the choice of x_k, y_k has been made so that for any $0 \leq k \leq (N-1)$ we have $x_{k+1} - x_k = y_{k+1} - y_k = \gamma(y_{k+1} - x_k)$. Applying v), using that $r > 1$ and the fact that the intervals (y_k, y_{k+1}) are disjoint, we have

$$\begin{aligned} \int_a^b w^r &\leq \sum_{k=0}^{N-1} \int_{x_k}^{x_{k+1}} w^r + \int_{b-sh}^b w^r \\ &\leq C(sh)^{1-r} \sum_{k=0}^{N-1} \left(\int_{y_k}^{y_{k+1}} w \right)^r + C(sh)^{1-r} \left(\int_{c-sh}^c w \right)^r \\ &\leq C_\gamma (c-a)^{1-r} \left(\int_b^c w \right)^r. \end{aligned}$$

So we have proved that v) \implies iv).

Finally, we will show that iv) \implies ii). Let $a < b < c$ with $b - a = 2(c - b)$. Let x be the midpoint between a, b . Using the one-sided doubling property we have

$$\begin{aligned} \frac{1}{b-a} \int_a^b w^r &= \frac{1}{b-a} \left(\int_a^x w^r + \int_x^b w^r \right) \\ &= \frac{1}{2} \left(\frac{1}{x-a} \int_a^x w^r + \frac{1}{b-x} \int_x^b w^r \right) \\ &\leq \frac{C}{2} \left(\left(\frac{1}{b-x} \int_x^b w \right)^r + \left(\frac{1}{c-b} \int_b^c w \right)^r \right) \\ &\leq \frac{C}{2} \left(\left(\frac{1}{c-b} \int_x^c w \right)^r + \left(\frac{1}{c-b} \int_b^c w \right)^r \right) \\ &\leq C \left(\frac{1}{c-b} \int_b^c w \right)^r. \end{aligned}$$

□

Remark. The equivalence of i) and iv) was first proved in [3].

The following lemma tells us that in the definition of A_p^+ we can take two intervals that are not contiguous. Note that in the case of RH_r^+ we have seen this in the previous lemma.

Lemma 2.6. *A weight w belongs to A_p^+ , $p > 1$ if, and only if, there exist $0 < \gamma \leq \frac{1}{2}$ and a constant C_γ such that $b - a = d - c = \gamma(d - a)$ for any $a < b < c < d$, then*

$$(2.7) \quad \int_a^b w \left(\int_c^d w^{1-p'} \right)^{p-1} \leq C_\gamma (b - a)^p.$$

Proof. If $w \in A_p^+$, $0 < \gamma \leq \frac{1}{2}$ and $a < b < c < d$, $b - a = d - c = \gamma(d - a)$ then

$$\int_a^b w \left(\int_c^d w^{1-p'} \right)^{p-1} \leq \int_a^c w \left(\int_c^d w^{1-p'} \right)^{p-1} \leq C(d - a)^p = C_\gamma (b - a)^p.$$

To prove that (2.7) implies A_p^+ we will show that (2.7) implies that for γ and a, b, c, d as above we have

$$\frac{1}{b - a} \int_a^b w \exp \left(\frac{1}{d - c} \int_c^d -\log(w) \right) \leq C.$$

Indeed,

$$(2.8) \quad \begin{aligned} \frac{1}{b - a} \int_a^b w \exp \left(\frac{1}{d - c} \int_c^d -\log(w) \right) \\ = \frac{1}{b - a} \int_a^b \left[w \exp \left(\frac{1}{d - c} \int_c^d \log(w)^{1-p'} \right) \right]^{p-1} \\ \leq \frac{1}{b - a} \int_a^b w \left(\frac{1}{d - c} \int_c^d w^{1-p'} \right)^{p-1} \leq C. \end{aligned}$$

In the same way we prove that $w^{1-p'}$ satisfies

$$(2.9) \quad \exp \left(\frac{1}{b - a} \int_a^b \log(w)^{p'-1} \right) \frac{1}{d - c} \int_c^d w^{1-p'} \leq C.$$

But, according to part *j*) of Theorem 1 in [7], (2.8) is equivalent to saying that $w \in A_\infty^+$ while (2.9) means that $w^{1-p'} \in A_\infty^-$, and according to Theorem 2 in [7] these two conditions imply $w \in A_1^+$. \square

Remark 2.10. We can easily see that $w \in A_1^+$ if, and only if, there exists $C > 0$ such that $\frac{1}{h} \int_{a-h}^a w \leq Cw(a + h)$ for almost every $a \in \mathbb{R}$ and $h > 0$.

3. THE EXTREME CASES: A_1^+ AND RH_∞^+

Theorem 3.1. *Let $w \in A_1^+$ with A_1^+ constant $C > 1$. Then $w \in RH_r^+$ for any $1 < r < \frac{C}{C-1}$, and this is the best possible range.*

Proof. Let us fix the interval $I = (a, b)$. We consider the truncation of w at height N defined by $w_N = \min(w, N)$, which also satisfies A_1^+ with a constant $C_N \leq C$. We claim that if $\lambda_I = M(w_N \chi_I)(b)$ and $E_\lambda = \{x \in I : w_N(x) > \lambda\}$ then

$$(3.2) \quad \int_{E_\lambda} w_N \leq C_N \lambda |E_\lambda| \quad \forall \lambda \geq \lambda_I.$$

Indeed, if $E_\lambda = I$ we do not even need the A_1^+ condition, since

$$w_N(E_\lambda) = \int_a^b w_N \leq M(w_N \chi_I)(b)(b-a) = \lambda_I(b-a) \leq C_N \lambda |E_\lambda|.$$

If $E_\lambda \neq I$ we fix $\varepsilon > 0$ and an open set O such that $E_\lambda \subset O \subset I$ and $|O| \leq \varepsilon + |E_\lambda|$. Let $J_k = (c, d)$, be one of the connected components of O . There are two cases:

- (1) $a \leq c < d < b$,
- (2) $a \leq c < d = b$.

In the first case $d \notin E_\lambda$ and then $w_N(d) \leq \lambda$. Now A_1^+ gives $\int_c^d w_N \leq C_N w_N(d)(d-c) \leq C_N \lambda(d-c)$. The second case is handled as the case $E_\lambda = I$, since $\int_c^b w_N \leq M(w_N \chi_I)(b)(b-c) \leq C \lambda(b-c)$. In any case $w_N(J_k) \leq C_N \lambda |J_k|$. Adding up we get

$$w_N(E_\lambda) \leq w_N(O) \leq C_N \lambda |O| \leq C_N \lambda (\varepsilon + |E_\lambda|).$$

Since ε was arbitrary we are done. Now we proceed in the standard way, i.e., we fix $s > -1$, multiply both sides of (3.2) by λ^s and integrate from λ_I to infinity to obtain,

$$\frac{1}{s+1} \int_I (w_N^{s+2} - \lambda_I^{s+1} w_N) \leq \frac{C_N}{s+2} \int_I w_N^{s+2}.$$

Now if $r = s+2 < \frac{C_N}{C_N-1}$ then $\frac{1}{s+1} - \frac{C_N}{s+2} > 0$, and we get

$$\int w_N^r \leq C_N \lambda_I^{r-1} \int_I w_N = C_N (M(w_N \chi_I)(b))^{r-1} \int_I w_N.$$

Now $C_N \leq C$ implies $\frac{C_N}{C_N-1} \geq \frac{C}{C-1}$, and therefore if $r \leq \frac{C}{C-1}$ then

$$\int_a^b w_N^r \leq C_N (M(w_N \chi_{(a,b)})(b))^{r-1} \int_a^b w_N \leq C (M(w \chi_{(a,b)})(b))^{r-1} \int_a^b w$$

and the monotone convergence theorem gives $w \in RH_r^+$. To see that this is the best possible range we consider the function

$$w(x) = x^{\frac{1}{c}-1} \chi_{(0,\infty)}(x).$$

It is clear that w does not satisfy $RH_{\frac{C}{C-1}}^+$ because $w^{\frac{C}{C-1}}(x) = \frac{1}{x}$ for $x > 0$. To see that it satisfies A_1^+ with the constant C , we consider three cases:

- (1) $a < b \leq 0$,
- (2) $a \leq 0 < b$,
- (3) $0 < a < b$.

In the first case there is nothing to check. In the second case $\frac{1}{b-a} \int_a^b w < \frac{1}{b} \int_0^b w(x) = \frac{C}{b} b^{\frac{1}{c}} = Cw(b)$. Finally, if $0 < a < b$, then $\int_a^b w = C(b^{\frac{1}{c}} - a^{\frac{1}{c}}) \leq C(b-a)w(b)$. \square

Remark. Note that if $C = 1$, then $w(x) = M^-w(x)$, and this implies that w is non-decreasing. This tells us that $w \in RH_\infty^+$.

Theorem 3.3. *If w satisfies RH_∞^+ with a constant $C > 1$, then $w \in A_p^+$ for all $p > C$, and this is the best possible range.*

Proof. A truncation argument as in Theorem 3.1 allows us to suppose that w is bounded away from zero, i.e. there exists $\beta > 0$ such that $w(x) \geq \beta$ for all x . Let us fix $I = (a, b)$ and consider $\lambda_I = m^+(w \frac{1}{\chi_I})(a)$. We claim that if $\lambda < \lambda_I$ and $E_\lambda = \{x \in I : w(x) < \lambda\}$, then

$$(3.4) \quad \lambda |E_\lambda| \leq C \int_{E_\lambda} w.$$

As before, if $E_\lambda = I$ then $\lambda |E_\lambda| = \lambda(b-a) < \lambda_I(b-a) = \int_a^b w \leq w(E_\lambda)$. If $E_\lambda \neq I$ then we approximate it by an open set $O = \bigcup J_k$ where $E_\lambda \subset O \subset I$ and $w(O) < \varepsilon + w(E_\lambda)$. Let us fix $J_k = (c, d)$. There are two cases:

- (1) $a < c$,
- (2) $a = c$.

In the first case $c \notin E_\lambda$ and then $\lambda(d-c) \leq w(c)(d-c) \leq Cm^+w(c)(d-c) \leq C \int_c^d w$. In the second case $\lambda(d-c) \leq \lambda_I(d-a) \leq \int_a^d w$, and (3.4) follows. If we multiply both sides of (3.4) by λ^{-r} with $r > 2$ and integrate we have

$$\int_0^{\lambda_I} \lambda^{1-r} \int \chi_{E_\lambda}(x) dx d\lambda \leq C \int_0^\infty \lambda^{-r} \int_{E_\lambda} w(x) dx d\lambda.$$

For the left hand side we obtain

$$\begin{aligned} \int_{\beta}^{\lambda_I} \lambda^{1-r} \int \chi_{E_{\lambda}}(x) \, dx \, d\lambda &= \frac{1}{2-r} \int_{\{x \in I: w(x) < \lambda_I\}} \lambda_I^{2-r} - w^{2-r} \, dx \\ &\geq \frac{1}{2-r} \int_I \lambda_I^{2-r} - w^{2-r} \, dx = \frac{1}{r-2} \int_I w^{2-r} - \frac{|I|}{r-2} \lambda_I^{2-r}, \end{aligned}$$

while the right hand side is equal to $\frac{C}{r-1} \int_I w^{2-r}$. Therefore

$$\frac{1}{r-2} \int_I w^{2-r} \leq \frac{C}{r-1} \int_I w^{2-r} + \frac{|I|}{r-2} \lambda_I^{2-r}.$$

If we choose $r > 2$ such that $C(r-2) < (r-1)$, we obtain that there exists C such that

$$(3.5) \quad \frac{1}{|I|} \int_I w^{2-r} \leq C \left(m^+ \left(\frac{w}{\chi_I} \right) (a) \right)^{2-r}.$$

We now claim that (3.5) implies that $w \in A_p^+$ with $p = \frac{r-1}{r-2}$. Let us fix $a < b < c$ and choose $x \in (a, b)$. If we keep in mind that $1 - p' = 2 - r$ we may write

$$\left(\frac{1}{c-a} \int_b^c w^{1-p'} \right)^{p-1} \leq \left(\frac{1}{c-x} \int_x^c w^{1-p'} \right)^{p-1} \leq C \left(m^+ \left(\frac{w}{\chi(x,c)} \right) (x) \right)^{-1},$$

but

$$\left(m^+ \left(\frac{w}{\chi(x,c)} \right) (x) \right)^{-1} = \left(\inf_{x < d < c} \frac{1}{d-x} \int_x^d w \right)^{-1} = \sup_{x < d < c} \frac{d-x}{\int_x^d w} = M_w \left(\frac{\chi(a,c)}{w} \right) (x).$$

We have thus proved that if $\lambda = \left(\frac{1}{c-a} \int_b^c w^{1-p'} \right)^{p-1}$ then

$$(a, b) \subset \left\{ x : CM_w \left(\frac{\chi(a,c)}{w} \right) (x) > \lambda \right\},$$

and the weak type of M_w with respect to the measure $w \, dx$ yields $\int_a^b w \leq C(c-a)^p \left(\int_b^c w^{1-p'} \right)^{1-p}$ which is A_p^+ . Finally, it can be checked that the function $w(x)$ which is 0 for $x < -1$, identically one for $x > 0$ and $|x|^{C-1}$ between -1 and 0 , satisfies RH_{∞}^+ with a constant C , but is not in A_C^+ . \square

Remark. Note that if $C = 1$, then $w(x) = m^+ w(x)$, and this implies that w is non-decreasing. This tells us that $w \in A_1^+$.

We have had several different characterizations of RH_r^+ , one involved the maximal operator, but dealt with one interval, and the others involved two intervals but no operator. We can now prove that for RH_∞^+ the situation is the same, we can characterize RH_∞^+ using two intervals instead of the minimal operator.

Corollary 3.6. *We have $w \in RH_\infty^+$ if, and only if, there exists C such that for any interval I ,*

$$(3.7) \quad \operatorname{ess\,sup}_I w \leq C \frac{1}{|I^+|} \int_{I^+} w.$$

Proof. It is immediate that (3.7) implies RH_∞^+ . Assume now that $w \in RH_\infty^+$. The preceding theorem tells us that $w \in A_p$ for some p , and therefore it satisfies the one-sided doubling condition. Therefore if $I = (a, b)$ is any interval, $I^+ = (b, c)$ and $x \in I$, we have

$$w(x) \leq \frac{C}{c-x} \int_x^c w \leq \frac{C}{c-b} \int_b^c w,$$

which is (3.7). □

Remark. Note that with this definition, we have $RH_\infty^+ \subset \bigcap_{r>1} RH_r^+$.

4. FACTORIZATION OF WEIGHTS IN RH_r^+ , $1 < r \leq \infty$

The theorems on the best range for weights in A_p^+ ($p > 1$) or in RH_r^+ , $r < \infty$ will be stated in terms of factorizations of the given weight. Therefore this section will be devoted to proving a factorization of functions in RH_r^+ . The bilateral case was studied in [2].

Definition 4.1. A function w is said to be essentially increasing if there exists C such that $w(x) \leq Cw(y)$ for any $x < y$.

Lemma 4.2. *A function belongs to $RH_\infty^+ \cap A_1^+$ if, and only if, it is essentially increasing.*

Proof. Assume that $w \in RH_\infty^+ \cap A_1^+$ and $x < y$, then $w(x) \leq C \frac{1}{y-x} \int_x^y w \leq Cw(y)$ and w is essentially increasing. Conversely, if w is essentially increasing then for any x and $h > 0$ we have $w(x) \leq \frac{C}{h} \int_x^{x+h} w$, hence $w \in RH_\infty^+$. On the other hand, $\frac{1}{h} \int_{x-h}^x w \leq Cw(x)$, so $w \in A_1^+$ □

Lemma 4.3. Let $1 < r \leq \infty$ and $1 \leq p < \infty$.

- (1) If u is essentially increasing and $v \in RH_r^+$ then $w \in RH_r^+$.
- (2) If u is essentially increasing and $v \in A_p^+$ then $w \in A_p^+$.

Proof. This proof follows immediately from Definition 4.1. □

Lemma 4.4. Let $1 < r \leq \infty$ and $1 \leq p < \infty$. We have $w \in RH_r^+ \cap A_p^+$ if, and only if, $w^r \in A_q^+$, with $q = r(p-1) + 1$.

Proof. Let $C_1 = RH_r^+(w)$ and $C_2 = A_p^+(w)$, $w \in RH_r^+ \cap A_p^+$ and $q = r(p-1) + 1$. Also note that $1 - q' = 1 - \frac{r(p-1)+1}{r(p-1)} = \frac{1}{r(1-p)}$,

$$\begin{aligned} & \left(\frac{1}{|I^-|} \int_{I^-} w^r \right) \left(\frac{1}{|I^+|} \int_{I^+} w^{r(1-q')} \right)^{q-1} \\ & \leq C_1 \left(\frac{1}{|I|} \int_I w \right)^r \left(\frac{1}{|I^+|} \int_{I^+} w^{1-p'} \right)^{r(p-1)} \\ & \leq C_1 C_2^r, \end{aligned}$$

and by Lemma 2.6 we have that $w^r \in A_q^+$.

If $w^r \in A_q^+$, then by Hölder's inequality

$$\begin{aligned} & \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I^+|} \int_{I^+} w^{-1/(p-1)} \right)^{p-1} \\ & \leq \left(\frac{1}{|I|} \int_I w^r \right)^{1/r} \left(\frac{1}{|I^+|} \int_{I^+} w^{-r/(q-1)} \right)^{(q-1)/r} \\ & \leq C^{1/r}, \end{aligned}$$

and we obtain in this way that $w \in A_p^+$. Now again by Hölder's inequality

$$1 = \frac{1}{|I^+|} \int_{I^+} w^{-1/p} w^{1/p} \leq \left(\frac{1}{|I^+|} \int_{I^+} w \right)^{1/p} \left(\frac{1}{|I^+|} \int_{I^+} w^{-p'/p} \right)^{1/p'},$$

so

$$\left(\frac{1}{|I^+|} \int_{I^+} w^{-1/(p-1)} \right)^{1-p} \leq \frac{1}{|I^+|} \int_{I^+} w,$$

and we get

$$\begin{aligned} \left(\frac{1}{|I|} \int_I w^r \right)^{1/r} & \leq C \left(\frac{1}{|I^+|} \int_{I^+} w^{-r/(q-1)} \right)^{-(q-1)/r} = C \left(\frac{1}{|I^+|} \int_{I^+} w^{-1/(p-1)} \right)^{1-p} \\ & \leq C \frac{1}{|I^+|} \int_{I^+} w, \end{aligned}$$

proving that $w \in RH_r^+$. □

Factorization Theorem for weights in $RH_r^+ \cap A_p^+$. A weight w satisfies $w \in RH_r^+ \cap A_p^+$ with $1 \leq p < \infty$, $1 < r \leq \infty$ if, and only if, there exist weights w_0 and w_1 such that $w_0 \in RH_r^+ \cap A_1^+$, $w_1 \in RH_\infty^+ \cap A_p^+$ and $w = w_0 w_1$.

Observe that since $\bigcup_{p < \infty} A_p^+ = \cap_{1 < r} RH_r^*$, every weight in RH_r^+ is in some A_p^+ . See [7].

P r o o f. Let us first consider the cases $p = 1$ or $r = \infty$.

If $p = 1$ and $r \leq \infty$, we put $w_1 = 1$ and $w_0 = w$, then obviously $w_0 \in RH_r^+ \cap A_1^+$ and $w_1 \in RH_\infty^+ \cap A_1^+$.

If $p \geq 1$ and $r = \infty$, we put $w_0 = 1$ and $w_1 = w$, obtaining $w_0 \in RH_\infty^+ \cap A_1^+$, $w_1 \in RH_\infty^+ \cap A_p^+$.

Conversely, given w_0 and w_1 , at least one of them belongs to $RH_\infty^+ \cap A_1^+$ (because $p = 1$ or $r = \infty$), so one of them is essentially increasing, therefore $w_0 w_1 \in RH_r^+ \cap A_p^+$ (Lemma 4.3).

Let us now suppose $p > 1$ and $r < \infty$. Let $w = w_0 w_1$ with $w_0 \in RH_r^+ \cap A_1^+$, and $w_1 \in RH_\infty^+ \cap A_p^+$. We want to see that $w \in RH_r^+ \cap A_p^+$. Note that for w_1 we have

$$\frac{1}{|I|} \int_I w_1^{1-p'} \leq C \left(\frac{1}{|I^-|} \int_{I^-} w_1 \right)^{1-p'} \leq C w_1 (a-h)^{1-p'},$$

which implies $w_1^{1-p'} \in A_1^-$ (Remark 2.10). Let $v = w_1^{1-p'}$, then $w_1 = v^{1-p}$ with $v \in A_1^-$, so $w = w_0 w_1 = w_0 v^{1-p}$ with $w_0 \in A_1^+$ and $v \in A_1^-$ (see [7]), and this implies $w \in A_p^+$.

Now

$$\begin{aligned} \frac{1}{|I|} \int_I w^r &= \frac{1}{|I|} \int_I w_0^r w_1^r \leq (\sup_I w_1)^r C \left(\frac{1}{|I^+|} \int_{I^+} w_0 \right)^r \\ &\leq C \left(\frac{1}{|I^+|} \int_{I^+} w_1 \right)^r \left(\inf_{I^+} w_0 \right)^r \\ &\leq C \left(\frac{1}{|I^+|} \int_{I^+} w_0 w_1 \right)^r, \end{aligned}$$

and by Lemma 2.5 we have $w \in RH_r^+$. Conversely, let $w \in RH_r^+ \cap A_p^+$, then by Lemma 4.4 $w^r \in A_q^+$ with $q = r(p-1) + 1$, there exists $v_0 \in A_1^+$ and $v_1 \in A_1^-$ such that $w^r = v_0 v_1^{1-q}$ (see [7]), or equivalently $w = v_0^{1/r} v_1^{(1-q)/r} = v_0^{1/r} v_1^{1-p}$. Let $w_0 = v_0^{1/r}$ and $w_1 = v_1^{1-p}$. We will see that $w_0 \in RH_r^+ \cap A_1^+$. We note,

$$\begin{aligned} \frac{1}{|I|} \int_I w_0^r &= \frac{1}{|I|} \int_I v_0 \leq C \inf_{I^+} v_0 \\ &\leq C \left(\frac{1}{|I^+|} \int_{I^+} v_0^{1/r} \right)^r = C \left(\frac{1}{|I^+|} \int_{I^+} w_0 \right)^r, \end{aligned}$$

and also

$$\begin{aligned} \frac{1}{|I|} \int_I w_0 &= \frac{1}{|I|} \int_I v_0^{1/r} \leq \left(\frac{1}{|I|} \int_I v_0 \right)^{1/r} \\ &\leq C \inf_{I^+} v_0^{1/r} = C \inf_{I^+} w_0. \end{aligned}$$

We only have to see now that $w_1 \in RH_\infty^+ \cap A_p^+$ and we are done.

First we claim

$$(4.5) \quad \text{if } w \in A_1^- \text{ then } w^{-\gamma} \in RH_\infty^+ \text{ for all } \gamma > 0.$$

In fact, by Hölder's inequality we have $\left(\frac{1}{|I|} \int_I w\right)^{-\gamma} \leq \frac{1}{|I|} \int_I w^{-\gamma}$ for any interval $I = (a, b)$, and as $w \in A_1^-$ we have that $Cw(x) \geq \frac{1}{|I|} \int_I w$ for almost every $x \in I^-$, and therefore

$$w(x)^{-\gamma} \leq C \left(\frac{1}{|I|} \int_I w \right)^{-\gamma} \leq \frac{1}{|I|} \int_I w^{-\gamma} \leq C \frac{1}{b-x} \int_x^b w^{-\gamma}.$$

Let $w_1 = v_1^{1-p}$. As $v_1 \in A_1^-$, then $w_1 \in RH_\infty^+$. Moreover,

$$\begin{aligned} \frac{1}{|I|} \int_I w_1 \left(\frac{1}{|I^+|} \int_{I^+} w_1^{1-p'} \right)^{p-1} &= \frac{1}{|I|} \int_I v_1^{1-p} \left(\frac{1}{|I^+|} \int_{I^+} v_1 \right)^{p-1} \\ &\leq \frac{1}{|I|} \int_I v_1^{1-p} (C \inf_I v_1)^{p-1} \\ &\leq \frac{C}{|I|} \int_I v_1^{1-p} v_1^{p-1} \leq C, \end{aligned}$$

i.e. $w_1 \in A_p^+$. □

Factorization Theorem for weights in A_∞^+ . A weight w satisfies $w \in A_\infty^+$ if, and only if, there exist $w_1 \in RH_\infty^+$ and $w_0 \in A_1^+$ such that $w = w_0 w_1$.

Proof. If $w \in A_\infty^+$ then $w \in A_q^+$ for some $1 < q < \infty$, so there exist $v_0 \in A_1^+$ and $v_1 \in A_1^-$ such that $w = v_0 v_1^{1-q}$. Let $w_0 = v_0$ and $w_1 = v_1^{1-q}$. By (4.5), $w_1 \in RH_\infty^+$. So we are done. Conversely, if $w_1 \in RH_\infty^+$, then $w_1 \in A_q^+$ for some $1 < q$, i.e., there exists C such that

$$\left(\frac{1}{|I|} \int_I w_1 \right)^{q'-1} \frac{1}{|I|} \int_{I^+} w_1^{1-q'} \leq C,$$

but then

$$(\sup_{I^-} w_1)^{q'-1} \frac{1}{|I|} \int_{I^+} w_1^{1-q'} \leq \left(\frac{1}{|I|} \int_I w_1 \right)^{q'-1} \frac{1}{|I|} \int_{I^+} w_1^{1-q'} \leq C,$$

and we get

$$\frac{1}{|I|} \int_{I^+} w_1^{1-q'} \leq C \inf_{I^-} w_1^{1-q'},$$

and it is easy to see that this inequality implies $w_1^{1-q'} \in A_1^-$. Then $v_1 = w_1^{1-q'} \in A_1^-$, so $w = w_0 w_1 = w_0 v_1^{1-q} \in A_q^+ \subset A_\infty^+$. \square

5. CLASSES A_p^+ AND RH_r^+

In this section we will use Theorems 3.1 and 3.3 and the factorization theorems to obtain the best ranges for the classes A_p^+ and RH_r^+ . As we shall see, the range of the index will depend on the factorization of the weights.

The following theorem gives us the precise range in A_p^+ for weights in RH_r^+ .

Theorem 5.1. *Let $w \in RH_r^+$, $w = w_0 w_1^{\frac{1}{r}}$ with $w_0 \in RH_\infty^+$ and $w_1 \in A_1^+$. Then $w \in A_p^+$ for all $p > C$ where $C = RH_\infty^+(w_0)$, and this is the best possible range.*

Proof. Let $w_0 \in RH_\infty^+$ and $w_1 \in A_1^+$. By Theorem 3.3, $w_0 \in A_p^+$ for all $p > C$. Let $p > C$, then there exists $\varepsilon > 0$ such that $w_0 \in A_{p-\varepsilon}^+$, so we choose $s > 1$ satisfying $1 - (p - \varepsilon)' = s(1 - p')$, and by Hölder's inequality

$$\begin{aligned} & \frac{1}{|I^-|} \int_{I^-} w_0 w_1 \left(\frac{1}{|I^+|} \int_{I^+} (w_0 w_1)^{1-p'} \right)^{p-1} \\ & \leq \left(\frac{1}{|I|} \int_I w_0 \right) \left(\frac{1}{|I^-|} \int_{I^-} w_1 \right) \left(\frac{1}{|I^+|} \int_{I^+} w_0^{s(1-p')} \right)^{\frac{(p-1)}{s}} \left(\frac{1}{|I^+|} \int_{I^+} w_1^{s'(1-p')} \right)^{\frac{(p-1)}{s'}} \\ & \leq C. \end{aligned}$$

To see that this is the best range, we consider w_0 as in Theorem 3.3 and $w_1 = 1$. \square

Remark 5.2. Given $w \in RH_r^+$ there exist $u \in RH_\infty^+$, and $v \in A_1^+$ such that $w = uv^{\frac{1}{r}}$. We only have to consider the factorization theorem and choose $u = w_1$ and $v = w_0^r$. We have to prove that $v \in A_1^+$. Keeping in mind that $w_0 \in RH_r^+ \cap A_1^+$ we have

$$\frac{1}{|I^-|} \int_{|I^-|} v = \frac{1}{|I^-|} \int_{|I^-|} w_0^r \leq C \left(\frac{1}{|I|} \int_{|I|} w_0 \right)^r \leq C w_0^r(x) = C v(x)$$

for almost every $x \in I^+$, i.e., $v \in A_1^+$.

The next theorem shows us the precise range of the higher integrability of $w \in RH_r^+$.

Theorem 5.3. *Let $w \in RH_r^+$, $w = uv^{1/r}$ with $u \in RH_\infty^+$ and $v \in A_1^+$. If $C = A_1^+(v)$ then $w \in RH_s^+$ for all $r \leq s < \frac{Cr}{C-1}$. The range of s is the best possible.*

Proof. Let $r < s < \frac{Cr}{C-1}$, let us choose $q > 1$ such that $s < \frac{Cr}{q(C-1)}$. As $1 < \frac{qs}{r} < \frac{C}{C-1}$, by Theorem 3.1 we have $v \in RH_{\frac{qs}{r}}^+$, and using Hölder's inequality we obtain that $u^s \in RH_\infty^+$ and $v \in A_1^+$ which yields

$$\begin{aligned} \frac{1}{|I|} \int_I w^s &= \frac{1}{|I|} \int_I u^s v^{s/r} \leq \left(\frac{1}{|I|} \int_I u^{q's} \right)^{1/q'} \left(\frac{1}{|I|} \int_I v^{qs/r} \right)^{1/q} \\ &\leq \sup_I u^s C \left(\frac{1}{|I^+|} \int_{I^+} v \right)^{s/r} \leq \frac{C}{|I^+|} \int_{I^+} u^s \left(\inf_{I^{++}} v \right)^{s/r} \\ &\leq C \sup_{I^+} u^s \inf_{I^{++}} v^{s/r} \leq C \left(\frac{1}{|I^{++}|} \int_{I^{++}} u \right)^s \inf_{I^{++}} v^{s/r} \\ &\leq C \left(\frac{1}{|I^{++}|} \int_{I^{++}} uv^{1/r} \right)^s = C \left(\frac{1}{|I^{++}|} \int_{I^{++}} w \right)^s, \end{aligned}$$

and we get that $w \in RH_s^+$ (Lemma 2.5).

To see this is the best range possible, we choose $v \in A_1^+$ as in Theorem 3.1 and $u = 1$, then $w = v^{1/r} \in RH_s^+$ for all $r \leq s < \frac{Cr}{C-1}$ ($C = A_1^+(v)$). If $s = \frac{Cr}{C-1}$ and $w \in RH_s^+$ then $v \in RH_{\frac{C}{C-1}}^+$, but we have seen (Theorem 3.1) that this can not happen. \square

The next theorem shows us which is the best range in RH_r^+ for a given weight in A_p^+ .

Theorem 5.4. *Let $w \in A_p^+$, $w = uv^{1-p}$ with $u \in A_1^+$, $v \in A_1^-$ and $C = A_1^+(u)$, then $w \in RH_r^+$ for all $1 < r < \frac{C}{C-1}$, this range being the best possible.*

Proof. By Theorem 3.1 we have $u \in RH_r^+$ for all $1 < r < \frac{C}{C-1}$ and we know that $v^{1-p} \in RH_\infty^+$, hence

$$\begin{aligned} \frac{1}{|I|} \int_I w^r &\leq \frac{1}{|I|} \int_I u^r \sup_I (v^{-r(p-1)}) \\ &\leq C \left(\frac{1}{|I^+|} \int_{I^+} u \right)^r \left(\frac{1}{|I^+|} \int_{I^+} v^{1-p} \right)^r \leq C \left(\inf_{I^{++}} u \right)^r \left(\sup_{I^+} v^{1-p} \right)^r \\ &\leq C \left(\inf_{I^{++}} u \right)^r \left(\frac{1}{|I^{++}|} \int_{I^{++}} v^{1-p} \right)^r \leq C \left(\frac{1}{|I^{++}|} \int_{I^{++}} w \right)^r. \end{aligned}$$

By Lemma 2.5 we conclude $w \in RH_r^+$.

To see this is the best range we take u as in Theorem 3.1 and $v = 1$. So we have $w = u \in A_p^+$ and $w \notin RH_{\frac{C}{C-1}}^+$. \square

Corollary 5.5. *Let $w = uv^{1-p} \in A_p^+$ with $u \in A_1^+$, $v \in A_1^-$ and $C = \max\{A_1^+(u), A_1^-(v)\}$. Then $w^\tau \in A_p^+$ for all $1 \leq \tau < \frac{C}{C-1}$ and the range is the best possible.*

Proof. By Theorem 5.4 we have that $w \in RH_\tau^+$ for all $1 \leq \tau < \frac{C}{C-1}$ and $w^{1-p'} \in RH_\tau^-$ for all $1 \leq \tau < \frac{C}{C-1}$. Let $a < d$, let us choose b, c such that $b - a = d - c = \frac{1}{4}(d - a)$, and we also consider the point $\frac{c+b}{2}$. Then we have four intervals, namely, $I^- = (a, b)$, $I = (b, \frac{b+c}{2})$, $I^+ = (\frac{b+c}{2}, c)$, and $I^{++} = (c, d)$. Now

$$\begin{aligned} \frac{1}{|I^-|} \int_{I^-} w^\tau \left(\frac{1}{|I^{++}|} \int_{I^{++}} w^{\tau(1-p')} \right)^{p-1} &\leq \left(\frac{1}{|I|} \int_I w \right)^\tau \left(\frac{1}{|I^+|} \int_{I^+} w^{1-p'} \right)^{\tau(p-1)}, \\ &\leq C^\tau, \end{aligned}$$

thus $w^\tau \in A_p^+$ (Lemma 2.6). Considering u as in Theorem 3.1, we see this is the best possible range. \square

Using Theorem 5.4 we will show the exact range of $q < p$ such that $w \in A_p^+$ implies $w \in A_q^+$.

Theorem 5.6. *Let $w = uv^{1-p} \in A_p^+$ with $u \in A_1^+$, $v \in A_1^-$ and $C = A_1^-(v)$. Then $w \in A_q^+$ for all $1 + \frac{(p-1)(C-1)}{C} < q < \infty$ and this is the best range for q .*

Proof. Note that $w^{1-p'} = vu^{1-p'} \in A_{p'}^-$, by Theorem 5.4 we have $w^{1-p'} \in RH_r^-$ for all $1 < r < \frac{C}{C-1}$. For the classes RH_r^- and A_p^- we have from Lemma 4.4 that $w^{(1-p')r} \in A_{q'}^-$ where $q' = r(p'-1)+1 = \frac{r}{p-1}+1$. But this is the same as $w^{1-q'} \in A_{q'}^-$, i.e., $w \in A_q^+$ for all $1 + (p-1)\frac{C-1}{C} < q$.

To see this is the best range, let $v(x) = x^{\frac{1-C}{C}}$ if $x \leq 0$ and equal to 0 if $x > 0$ and $u = 1$ for all x . Note that $v \in A_1^-$ and $A_1^-(v) = C$. Then $w = v^{1-p} \in A_p^+$ and $w \in A_q^+$ for all $q > 1 + (p-1)\frac{C-1}{C}$. Observe that $w \notin A_{1+(p-1)\frac{C-1}{C}}^+$. \square

Finally, the last theorem gives us the best possible range for a weight in A_∞^+ .

Theorem 5.7. *Let $w \in A_\infty^+$, $w = w_0 w_1$, $w_0 \in A_1^+$, $w_1 \in RH_\infty^+$ and $C = RH_\infty^+(w_1)$. Then $w \in A_p^+$ for all $p > C$. The range of p 's is the best possible.*

Proof. Note that $w_1 \in RH_\infty^+$ implies $w_1 \in A_p^+$ for all $p > C$, hence

$$\begin{aligned} & \frac{1}{|I|} \int_I w_0 w_1 \left(\frac{1}{|I|^{++}} \int_{I^{++}} (w_0 w_1)^{1-p'} \right)^{p-1} \\ & \leq \sup_I (w_1) \frac{1}{|I|} \int_I w_0 \left(\sup_{I^{++}} w_0^{1-p'} \right)^{p-1} \left(\frac{1}{|I|^{++}} \int_{I^{++}} w_1^{1-p'} \right)^{p-1} \\ & \leq C \frac{1}{|I|^+} \int_{I^+} w_1 \inf_{I^+} (w_0) \sup_{I^{++}} (w_0^{-1}) \left(\frac{1}{|I|^{++}} \int_{I^{++}} w_1^{1-p'} \right)^{p-1} \\ & \leq C \sup_{I^{++}} (w_0^{-1}) \frac{1}{|I^+|} \int_{I^+} w_0 \\ & \leq C \frac{1}{(\inf_{I^{++}} w_0)} \inf_{I^{++}} w_0 \leq C, \end{aligned}$$

and by Lemma 2.6 we have $w \in A_p^+$ for all $p > C$.

To see this is the best range, we consider $w(x) = 0$ if $x \leq -1$, $|x|^{C-1}$ if $-1 < x \leq 0$ and 1 if $x \geq 0$. \square

References

- [1] *H. Aimar, L. Forzani and F. J. Martín-Reyes*: On weighted inequalities for one-sided singular integrals. Proc. Amer. Math. Soc. *125* (1997), 2057–2064.
- [2] *D. Cruz-Uribe and C. J. Neugebauer*: The structure of reverse Hölder classes. Trans. Amer. Math. Soc. *347* (1995), 2941–2960.
- [3] *D. Cruz-Uribe, C. J. Neugebauer and V. Olesen*: The one-sided minimal operator and the one-sided reverse Hölder inequality. Stud. Math *116* (1995), 255–270.
- [4] *P. Guan and E. Sawyer*: Regularity estimates for oblique derivative problem. Anal. of Mathematics, Second Series *157* (1) (1993), 1–70.
- [5] *F. J. Martín-Reyes*: New proofs of weighted inequalities for the one-sided Hardy-Littlewood maximal functions. Proc. Amer. Math. Soc. *117*, 691–698.
- [6] *F. J. Martín-Reyes, P. Ortega and A. de la Torre*: Weighted inequalities for one-sided maximal functions. Trans. Amer. Math. Soc. *319* (2) (1990), 517–534.
- [7] *F. J. Martín-Reyes, L. Pick and A. de la Torre*: A_∞^+ condition. Canad. J. Math. *45* (6) (1993), 1231–1244.
- [8] *C. J. Neugebauer*: The precise range of indices for the RH_r and A_p weight classes. Preprint.
- [9] *E. Sawyer*: Weighted inequalities for the one-sided Hardy-Littlewood maximal function. Trans. Amer. Math. Soc. *297* (1986), 53–61.

Author's address: M. S. Riveros, FaMAF, Universidad Nacional de Córdoba, (5000) Córdoba, Argentina, e-mail: sriveros@mate.uncor.edu; A. de la Torre, Análisis Matemático, Facultad de Ciencias, Universidad de Málaga, (29071) Málaga, Spain, e-mail: torre@anamat.cie.uma.es.